arXiv:q-alg/9503012v1 21 Mar 1995

ACKNOWLEDGEMENTS

I would like to express my deep gratitude to my advisor Igor Frenkel for his guidance and encouragement throughout my study at Yale. Most of what I know about representation theory, mathematical physics and special functions comes from him.

Also, I want to thank my friend and co-author Pavel Etingof; most of the results in this dissertation are based on our joint work, and would never be written without him; the idea of systematic study of the weighted traces of intertwiners belongs to him. His enthusiasm and new ideas often helped to find a new way when I was ready to give up.

My understanding of the subjects discussed in this dissertation benefited a lot from fruitful discussions with other people, and among them Ivan Cherednik, Jintai Ding, Howard Garland, Ian Grojnowski, David Kazhdan, Ian Macdonald, Masatoshi Noumi, Gregg Zuckerman and many others.

Financial support during my first three years of study was provided by Yale and during the final year by Alfred P. Sloan graduate dissertation fellowship.

Finally – and most important of all – I want to thank my wife Варя for her love and patience – even when I started discussing mathematics with Pavel while skiing in New Hampshire...

INTRODUCTION

This dissertation is devoted to study of some structures appearing in representation theory of simple Lie algebras (quantum groups, affine Lie algebras) and their relations with the theory of special functions and mathematical physics. This subject, of course, has a long history. A representation-theoretic approach to special functions was developed in the 40-s and 50-s in the works of I.M.Gelfand, M.A.Naimark, N.Ya.Vilenkin, and their collaborators (see [V], [VK]). The essence of this approach is the fact that most classical special functions can be obtained as suitable specializations of matrix elements or characters of representations of groups. In geometrical terms, interesting special functions appear as spherical functions on symmetric spaces associated with G, which was studied by Harish-Chandra and Helgason (see [HC], [W]).

However, this approach does not cover all interesting cases. For example, it was shown in recent works on representations of (quantum) affine Lie algebras that matrix elements of intertwining operators between certain representations of these algebras are interesting special functions – (q-)hypergeometric functions and their generalizations [TK, FR].

In this dissertation we present a more general scheme, which includes both the classical theory of characters and matrix elements and intertwining operators. This scheme was suggested in the joint paper of the author and Pavel Etingof [EK1] and developed in papers [EK2– EK5, E1, K1, K2]. Briefly it can be described as "theory of characters for intertwining operators", which we call generalized characters. This approach allows to get in a natural way many interesting special functions (for example, Lamé functions, Macdonald's polynomials) and easily prove a number of their properties.

Here are the main ideas of this approach. Let $\mathfrak{g}$ be a simple finite-dimensional Lie algebra over $\mathbb{C}$, and let G be the corresponding compact real Lie group. An important role in the representation theory of G play the characters of finite-dimensional representations; in particular, they form a distinguished basis in the space of functions on G invariant with respect to conjugation; in other words, they are zonal spherical functions on the symmetric space $G \times G/G_{diag}$. We can generalize this considering *equivariant* functions, i.e. functions on G with values in a representation U which satisfy

$$f(gxg^{-1}) = gf(x).$$

This is a particular example of so-called μ-spherical functions which were studied in papers of Harish-Chandra (see [HC, W]).

It turns out that this space has equally natural description. Namely, let V be any finite-dimensional representation of G, and $\Phi : V \to V \otimes U$ be an intertwining operator. Define the corresponding generalized character χ_Φ by

$$\chi_\Phi(g) = \mathrm{Tr}_V(\Phi g).$$

This is a natural generalization of usual characters; however, it takes values not in $\mathbb{C}$ but in the representation U. It is easy to see that χ_Φ is equivariant; moreover, every equivariant function is a linear combination of generalized characters. Note that due to the equivariance, the generalized character is completely determined by its values on the elements of the form $g = e^h, h$ from Cartan subalgebra of $\mathfrak{g}$. For this reason, from now on we consider χ_Φ as a function on Cartan subalgebra.

In the classical case many properties of representations could be expressed in terms of their characters. The same holds here: many properties of intertwining operators can be expressed in terms of corresponding generalized characters. For example, the natural inner product on the space of intertwining operators gives an inner product on generalized characters and an analogue of the orthogonality theorem for them. Similarly, the fact that Φ commutes with the center of $U\mathfrak{g}$ yields differential equations satisfied by the χ_Φ – each element of the center gives a differential equation. Thus, the generalized characters are common eigenfunctions of a family of commuting differential operators, obtained from the center of $U\mathfrak{g}$.

These generalized characters are a rich source of special functions. In general, these functions have not been studied before. However, in some special cases they coincide with well-known objects. For example, if U is a trivial module then this is nothing but the usual characters and thus we recover the classical theory.

More generally, the generalized characters (considered as functions on the Cartan subalgebra) take values in the zero-weight subspace $U[0]$; thus, if we consider the case of Lie algebra $\mathfrak{sl}_n$ and take U to be the space of homogeneous polynomials in n variables of degree $(k-1)n$ then $U[0]$ is one-dimensional and thus the generalized characters can be considered as complex-valued. In this case we show that the ratio χ_Φ/δ^{k-1}, where δ is the Weyl denominator, is the Jack symmetric polynomial. The simplest way to describe Jack polynomials – or more generally, Jacobi polynomials, which are generalization of Jack polynomials to the case of arbitrary root systems introduced by Heckman and Opdam – is to say that they are eigenfunctions of (conjugated by certain function) Sutherland operator:

$$L_k = \Delta - 2k(k-1) \sum_{\alpha \in R^+} \frac{1}{(2 \sinh \frac{(\alpha,h)}{2})^2},$$

where Δ is the Laplace operator on $\mathfrak{h}$. For special values of k this operator is just the radial part of the Laplace operator on certain symmetric spaces associated with the group G, and the eigenfunctions are zonal spherical functions. Heckman and Opdam have showed that for any k, these polynomials have a number of remarkable properties. In particular, they showed that the Sutherland operator can be included in a commutative family of differential operators, isomorphic to the algebra $S(\mathfrak{h})^W \simeq \mathcal{Z}(U\mathfrak{g})$ of Weyl group invariant polynomials on $\mathfrak{h}$; also, these polynomials are orthogonal with respect to a certain inner product. All of these results are highly non-trivial and required a good deal of ingenuity. We will show that our representation-theoretic approach allows to obtain both these properties immediately as a corollary of the general results about the generalized characters. This new interpretation also suggests new generalizations, such as vector-valued analogue of these polynomials or affine analogue (see below).

In a similar way, the generalized characters can be defined for the quantum group $U_q\mathfrak{g}$, corresponding to $\mathfrak{g}$. All the constructions above can be generalized to this case with some minor changes; most importantly, differential operators should be replaced by difference operators. We will show that for the case $\mathfrak{g} = \mathfrak{sl}_n$ and U chosen as above we get the famous Macdonald's polynomials – a family of symmetric polynomials which was recently introduced by I. Macdonald and has been the object of intensive study since that time. Again, this approach allows us to reprove many properties of Macdonald's polynomials (inner product identity, symmetry identity) in a very simple way.

In a similar way, we can define generalized characters for affine Lie algebras. It turns out that if we take U to be tensor product of evaluation representations then these characters are precisely the correlation functions on the torus of the Wess-Zumino-Witten (WZW) model of conformal filed theory. We deduce differential equations, describing dependence of these generalized characters on the modular parameter of the torus and the parameters of the evaluation representations; we call these equations elliptic Knizhnik-Zamolodchikov equations (this was first done, in a different language, by Bernard [Be]) and study their monodromies.

Again, in the case $\mathfrak{g} = \mathfrak{sl}_n$, U – evaluation representation corresponding to the representation U of $\mathfrak{sl}_n$ described above we get an affine analogue of the theory of Jacobi polynomials. This is closely related with the elliptic analogue of the Sutherland operator. We can generalize to this case some results from the theory of usual Jacobi polynomials; there are also arise new phenomena, such as modular properties of these polynomials (as functions of the modular parameter of the torus).

CHAPTER I

BASIC DEFINITIONS

1.1 Simple Lie algebras and their representations

In this whole work, all the objects are always defined over the ground field $\mathbb{C}$ of complex numbers or its extensions. In this section we briefly list the main facts on the simple Lie algebras and their representations which we are going to use; all of these facts are quite standard and can be found in any textbook on Lie algebras (see, for example, [Hu, B]).

Let $\mathfrak{g}$ be a simple Lie algebra over $\mathbb{C}$ of rank r, $\mathfrak{h} \subset \mathfrak{g}$ – its Cartan subalgebra, $U\mathfrak{g}$ – its universal enveloping algebra. Then $\mathfrak{g}$ has the root decomposition:

$$\mathfrak{g} = \mathfrak{h} \oplus \bigoplus_{\alpha \in R} \mathfrak{g}_\alpha,$$

where $R \subset \mathfrak{h}^*$ is the corresponding root system. We fix a polarization of R: $R = R^+ \sqcup -R^+$, where R^+ is the subset of positive roots. We denote by $\alpha_1, \dots, \alpha_r \in R^+$ the basis of simple roots. The polarization of roots gives rise to polarization of $\mathfrak{g} : \mathfrak{g} = \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}^+$, where $\mathfrak{n}^\pm = \bigoplus_{\alpha \in R^\pm} \mathfrak{g}_\alpha$, and corresponding polarization of $U\mathfrak{g} : U\mathfrak{g} = U\mathfrak{n}^- \otimes U\mathfrak{h} \otimes U\mathfrak{n}^+$.

We fix an invariant symmetric bilinear form $(\, , \,)$ on $\mathfrak{g}$ by the condition that for the associated bilinear form on $\mathfrak{h}^*$ we have $(\alpha, \alpha) = 2$ for long roots; this form allows us to identify $\mathfrak{h}^* \simeq \mathfrak{h} : \lambda \mapsto h_\lambda$. Abusing the language, we will also use the notation $(\, , \,)$ for the associated bilinear form on $\mathfrak{h}^*$. We denote by $\langle \, , \, \rangle$ the canonical pairing $\mathfrak{h} \otimes \mathfrak{h}^* \to \mathbb{C}$.

As usual, for every $\alpha \in R$ we define the dual root $\alpha^\vee = \frac{2h_\alpha}{(\alpha,\alpha)} \in \mathfrak{h}$ and introduce the following notions:

$Q = \bigoplus \mathbb{Z}\alpha_i$ – root lattice;

$Q^+ = \bigoplus \mathbb{Z}_+\alpha_i$;

$Q^\vee = \bigoplus \mathbb{Z}\alpha_i^\vee \subset \mathfrak{h}$ – coroot lattice;

$P = \{\lambda \in \mathfrak{h}^* | \langle \lambda, \alpha^\vee \rangle \in \mathbb{Z}\}$ – weight lattice;

$P^+ = \{\lambda \in \mathfrak{h}^* | \langle \lambda, \alpha_i^\vee \rangle \in \mathbb{Z}_+\}$ – cone of integer dominant weights;

ω_i – fundamental weights: $\langle \omega_i, \alpha_j^\vee \rangle = \delta_{ij}$;

$P^\vee = \{h \in \mathfrak{h} | \langle h, \alpha \rangle \in \mathbb{Z}\}$ – coweight lattice;
$\rho = \frac{1}{2}\sum_{\alpha \in R^+} \alpha = \sum \omega_i$;
$\theta \in R$ – the highest root: $\theta - \alpha \in Q^+$ for every $\alpha \in R$;
$h^\vee = \langle \rho, \theta^\vee \rangle + 1$ – dual Coxeter number for $\mathfrak{g}$;
We have a (partial) order on P defined as follows: $\lambda \le \mu$ if $\mu - \lambda \in Q^+$.

We denote by $\mathbb{C}[P]$ the group algebra of the abelian group P, i.e. the algebra over $\mathbb{C}$ spanned by formal exponents $e^\lambda, \lambda \in P$ with relations $e^0 = 1, e^{\lambda+\mu} = e^\lambda e^\mu$. If an element $f \in \mathbb{C}[P]$ can be written in the form $f = \sum_{\mu \le \lambda} a_\mu e^\mu, a_\lambda \ne 0$ then we say that $a_\lambda e^\lambda$ is the highest term of f and write $f = a_\lambda e^\lambda +$ *lower order terms.*

As usual, we denote by W the Weyl group of R and by $l(w)$ length of an element $w \in W$ with respect to the generators $s_i = s_{\alpha_i}$. This group acts naturally on $\mathfrak{h}^*$, preserving R; thus, it acts on P and $\mathbb{C}[P]$. Of special interest for us will be the algebra $\mathbb{C}[P]^W$ of W-invariant elements in $\mathbb{C}[P]$. We will often use the fact that the orbitsums

$$m_\lambda = \sum_{\mu \in W\lambda} e^\mu, \quad \lambda \in P^+ \tag{1.1.1}$$

form a basis in $\mathbb{C}[P]^W$, which follows from the fact that for every $\lambda \in P$ the orbit $W\lambda$ contains precisely one element from P^+.

We choose a basis $e_\alpha \in \mathfrak{g}_\alpha, f_\alpha \in \mathfrak{g}_{-\alpha}$ for $\alpha \in R^+$ such that $(e_\alpha, f_\alpha) = 1$; then $[e_\alpha, f_\alpha] = h_\alpha$. In particular, if $\alpha = \alpha_i$ is a simple root then the elements

$$e_i = \frac{2}{(\alpha_i, \alpha_i)} e_{\alpha_i}, \quad f_i = \frac{2}{(\alpha_i, \alpha_i)} f_{\alpha_i},$$
$$h_i = \alpha_i^\vee = \frac{2}{(\alpha_i, \alpha_i)} h_{\alpha_i}$$

satisfy the usual relations of the Lie algebra $\mathfrak{sl}_2$. Moreover, $e_i, f_i, h_i, i = 1 \dots r$ generate $\mathfrak{g}$.

Let a_i be any orthonormal basis in $\mathfrak{g}$. Define the Casimir element $C \in U\mathfrak{g}$ by $C = \sum a_i^2$. This element is central and does not depend on the choice of orthonormal basis. Therefore, it is easy to check that it can be written as follows: let $x_l, l = 1 \dots r$ be an orthonormal basis in $\mathfrak{h}$; then $C = \sum_{\alpha \in R^+} e_\alpha f_\alpha + f_\alpha e_\alpha + \sum x_l^2$.

We will also use the following useful identity:

$$\Delta(C) = C \otimes 1 + 1 \otimes C + 2\Omega, \tag{1.1.2}$$

where

$$\Omega = \sum a_i \otimes a_i = \sum_{\alpha \in R^+} e_\alpha \otimes f_\alpha + f_\alpha \otimes e_\alpha + \sum x_l \otimes x_l \tag{1.1.3}$$

is the $\mathfrak{g}$-invariant element in $\mathfrak{g} \otimes \mathfrak{g}$, and $\Delta : U\mathfrak{g} \to U\mathfrak{g} \otimes U\mathfrak{g}$ is the comultiplication: $\Delta(x) = x \otimes 1 + 1 \otimes x, x \in \mathfrak{g}$ and $\Delta(ab) = \Delta(a)\Delta(b)$.

In general, there is no explicit construction of the whole center of $U\mathfrak{g}$. However, its structure as a graded algebra (and more over, as a module over $U\mathfrak{g}$) is known.

Theorem 1.1.1.

$$\mathcal{Z}(U\mathfrak{g}) \simeq (S\mathfrak{h})^W.$$

We will construct the isomorphism (Harish-Chandra isomorphism) later; now we only say that under this isomorphism the Casimir element $C \mapsto p_2 : p_2(\lambda) = (\lambda, \lambda) - (\rho, \rho)$. Note that since it is known that $(S\mathfrak{h})^W$ is a free polynomial algebra, the same is true for $\mathcal{Z}(U\mathfrak{g})$.

Also, there exists a unique involutive algebra automorphism (Chevalley involution) $\omega : \mathfrak{g} \to \mathfrak{g}$ such that

$$\begin{aligned} \omega(e_i) &= -f_i, \\ \omega(f_i) &= -e_i, \\ \omega(h) &= -h, \quad h \in \mathfrak{h}. \end{aligned} \tag{1.1.4}$$

Let us now consider the representation theory of $\mathfrak{g}$. If V is a $\mathfrak{g}$-module then we denote an action of $x \in \mathfrak{g}$ in V by $\pi_V(x)$. We will always consider modules with a weight decomposition: $V = \bigoplus_{\lambda \in \mathfrak{h}^*} V[\lambda]$, where $V[\lambda]$ are finite-dimensional spaces such that $hv = \langle \lambda, h \rangle v$ if $v \in V[\lambda], h \in \mathfrak{h}$. We will call $v \in V[\lambda]$ vectors of weight λ. For every $\lambda \in \mathfrak{h}^*$ we define the Verma module M_λ by the following conditions:

1. M_λ is spanned over $U\mathfrak{g}$ by a single vector v_λ (highest weight vector) such that $e_\alpha v_\lambda = 0$, v_λ has weight λ.
2. M_λ is a free module over $U\mathfrak{n}^-$.

These modules are simplest examples of modules from category $\mathcal{O}$. By definition, a $\mathfrak{g}$-module V is said to be from category $\mathcal{O}$, if it has weight decomposition, is finitely generated and satisfies the following condition: for every $v \in V$, the space $U\mathfrak{n}^+ v$ is finite-dimensional.

Let I_λ is the maximal ideal in M_λ which does not contain v_λ (it exists and is unique); denote by $L_\lambda = M_\lambda / I_\lambda$ the corresponding irreducible highest-weight module. For generic λ (that is, for all λ except a subvariety of codimension 1), $M_\lambda = L_\lambda$. Moreover, it is known that there is a unique bilinear form on M_λ (Shapovalov form) such that $(v_\lambda, v_\lambda) = 1$ and $(v_1, xv_2) = -(\omega(x)v_1, v_2)$ for all $x \in \mathfrak{g}$. This form is symmetric and its kernel is I_λ; thus, it descends to a non-degenerate form on L_λ.

It is known that L_λ is finite-dimensional iff $\lambda \in P^+$, and every finite-dimensional irreducible representation of $\mathfrak{g}$ is obtained in this way. Moreover, if $\lambda \in P^+$ then the dual representation L_λ^* is also finite-dimensional irreducible: $L_\lambda^* = L_{\lambda^*}$, where

$$\lambda^* = -w_0(\lambda),$$

w_0 being the longest element in the Weyl group.

If $c \in \mathcal{Z}(U\mathfrak{g})$ is a central element, then $c|_{M_\lambda} = \chi(c)(\lambda + \rho) \operatorname{Id}_{M_\lambda}$, where $\chi : \mathcal{Z}(U\mathfrak{g}) \to (S\mathfrak{h})^W$ is the Harish-Chandra isomorphism mentioned before. The same holds for any subquotient of M_λ, in particular, for L_λ. Note that for the Casimir element we have $C|_{M_\lambda} = (\lambda, \lambda+2\rho)\,\mathrm{Id}_{M_\lambda}$; thus, in the adjoint representation C acts by multiplication by $2h^\vee$.

For a finite-dimensional representation V of $\mathfrak{g}$ let $\operatorname{ch} V = \sum_{\lambda\in\mathfrak{h}^*} \dim V[\lambda] e^\lambda \in \mathbb{C}[P]$. For an irreducible representation, the character is given by the Weyl formula:

$$\operatorname{ch} L_\lambda = \frac{\sum_{w\in W} (-1)^{l(w)} e^{w(\lambda+\rho)}}{\delta}, \quad \lambda \in P^+, \tag{1.1.5}$$

where

$$\delta = \prod_{\alpha\in R^+} (e^{\alpha/2} - e^{-\alpha/2}) = e^\rho \prod_{\alpha\in R^+} (1 - e^{-\alpha}) \tag{1.1.6}$$

is the Weyl denominator. Obviously, $\operatorname{ch} L_\lambda = e^\lambda +$ *lower order terms.*

More generally, if V is a module from category $\mathcal{O}$ then we can define its character by the same formula $\operatorname{ch} V = \sum_{\lambda\in\mathfrak{h}^*} \dim V[\lambda] e^\lambda$. However, in this case it lies not in $\mathbb{C}[P]$ but in its completion $\overline{\mathbb{C}[P]}$, which is defined as follows:

$$\overline{\mathbb{C}[P]} = \{\sum_{n=0}^{\infty} a_n e^{\lambda_n} \,|\, \lim (\rho, \lambda_n) = +\infty\}. \tag{1.1.7}$$

In the following we will be especially interested in the case $\mathfrak{g} = \mathfrak{sl}_n$, i.e. the root system of type A_{n-1}. In this case the root system and related objects admit the following explicit realization:

$\mathfrak{h}^* = \{(\lambda_1, \dots, \lambda_n) | \sum \lambda_i = 0\} \subset \mathbb{C}^n$;

$R = \{\varepsilon_i - \varepsilon_j\}_{i\neq j}$, where ε_i is the standard basis in $\mathbb{C}^n$: $\varepsilon_i = (0, \dots, 0, 1, 0, \dots, 0) \in \mathbb{Z}^n$ (1 in the i-th place);

$(\lambda, \mu) = \sum \lambda_i \mu_i$;

$R^+ = \{\varepsilon_i - \varepsilon_j\}_{i<j}$,;

$\alpha_i = \varepsilon_i - \varepsilon_{i+1}$;

$Q = Q^\vee = \{(\lambda_1, \dots, \lambda_n) \subset \mathbb{Z}^n | \sum \lambda_i = 0\}$;

$P = \{(\lambda_1, \dots, \lambda_n) \subset \frac{1}{n}\mathbb{Z}^n | \sum \lambda_i = 0, \lambda_i - \lambda_j \in \mathbb{Z}\}$;

$P^+ = \{(\lambda_1, \dots, \lambda_n) \subset \frac{1}{n}\mathbb{Z}^n | \sum \lambda_i = 0, \lambda_i - \lambda_{i+1} \in \mathbb{Z}_+\}$;

$\rho = (\frac{n-1}{2}, \frac{n-3}{2}, \dots, \frac{1-n}{2})$;

$W = S_n$;

$\mathbb{C}[P] \simeq \{$homogeneous polynomials in $x_1^{\pm 1}, \dots, x_n^{\pm 1}$ of degree zero$\}$ (this isomorphism is given by letting $e^{\varepsilon_i} = x_i$).

Note also in this case $\lambda, \mu \in P, \lambda < \mu$ with respect to the order on P we defined before implies that $\lambda \prec \mu$, where $\prec$ is the lexicographic order on $\mathbb{R}^n$.

Many results for the Lie algebra $\mathfrak{sl}_n$ are also valid for (reductive) Lie algebra $\mathfrak{gl}_n$ with suitable changes. Namely, the group algebra $\mathbb{C}[P]$ should be replaced by the algebra $\mathbb{C}[x_1, \dots, x_n]$, P^+ should be replaced by the set of all partitions in n parts, i.e. sequences $(\lambda_1, \dots, \lambda_n)$ such that $\lambda_i \in Z_+, \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0$, ρ should be replaced by $(n-1, n-2, \dots, 0)$ and Weyl denominator should be replaced by Vandermonde determinant $\prod_{i<j}(x_i - x_j)$.

1.2 Quantum groups

In this section we briefly list the main definitions from the theory of quantum groups (see [Dr1, J1,J2]). In this section all the objects are considered over the ground field $\mathbb{C}_q = \mathbb{C}(q^{1/2N})$, where $q^{1/2N}$ is a formal variable, and $N = |P/Q|$ (this will be justified below).

Let $R, R^+, ..$ be as defined in Section 1.1. Let us renormalize the inner product in $\mathfrak{h}$, introducing $(x,y)' = m(x,y)$, where m is chosen so that $(\alpha,\alpha)' = 2$ for **short** roots. Thus, m can take values 1 (for simply-laced root systems), 2 (for root systems of type B, C, F) or 3 (for G_2). This renormalization is quite standard in the theory of quantum groups. In this case $d_i = \frac{(\alpha_i,\alpha_i)'}{2}$ are positive integers such that $d_i a_{ij} = d_j a_{ji}$, where $a_{ij} = 2(\alpha_i,\alpha_j)/(\alpha_i,\alpha_i)$ is the Cartan matrix, and g.c.d.$(d_i) = 1$.

In this section we will identify $\mathfrak{h}$ and $\mathfrak{h}^*$ using the form $(\ ,\ )'$; (this differs by a factor of m from the identification of the previous chapter!); then $\alpha_i = d_i \alpha_i^\vee$, and therefore $Q \subset Q^\vee, P \subset P^\vee$. Since, by definition, for every $\lambda \in P$ we have $N\lambda \in Q$, this implies $(\lambda,\mu)' \in \frac{1}{N}\mathbb{Z}$ for every $\lambda \in P, \mu \in P$, which is the reason why have chosen the ground field to be $\mathbb{C}(q^{1/2N})$.

Define the quantum group $U_q\mathfrak{g}$ as an associative algebra over $\mathbb{C}_q$ with generators $e_i, f_i, q^h, h \in \frac{1}{2}P^\vee \subset \mathfrak{h}, i = 1\dots r$ and relations

$$\begin{gathered}
q^0 = 1, \quad q^{a+b} = q^a q^b,\\
q^h f_j q^{-h} = q^{-\langle h,\alpha_j\rangle} f_j,\\
q^h e_j q^{-h} = q^{\langle h,\alpha_j\rangle} e_j,\\
[e_i, f_j] = \delta_{ij} \frac{q^{d_i h_i} - q^{-d_i h_i}}{q^{d_i} - q^{-d_i}},
\end{gathered} \tag{1.2.1}$$

$$\begin{gathered}
\sum_{n=0}^{1-a_{ij}} \frac{(-1)^n}{[n]_i! [1 - a_{ij} - n]_i!} e_i^n e_j e_i^{1-a_{ij}-n} = 0,\\
\sum_{n=0}^{1-a_{ij}} \frac{(-1)^n}{[n]_i! [1 - a_{ij} - n]_i!} f_i^n f_j f_i^{1-a_{ij}-n} = 0,
\end{gathered} \tag{1.2.2}$$

where, as before, $h_i = \alpha_i^\vee$ and

$$[n]_i = \frac{q^{nd_i} - q^{-nd_i}}{q^{d_i} - q^{-d_i}}, \quad [n]_i! = [1]_i \dots [n]_i. \tag{1.2.3}$$

This is a Hopf algebra with the following comultiplication, counit and antipode:

(1.2.4)

$$\begin{gathered}
\Delta e_i = e_i \otimes q^{-d_i h_i/2} + q^{d_i h_i/2} \otimes e_i, \ \Delta f_i = f_i \otimes q^{-d_i h_i/2} + q^{d_i h_i/2} \otimes f_i,\\
\Delta q^h = q^h \otimes q^h,\\
\epsilon(q^h) = 1, \epsilon(e_i) = \epsilon(f_i) = 0, S(e_i) = -q^{-d_i} e_i, S(f_i) = -q^{d_i} f_i, S(q^h) = q^{-h}.
\end{gathered}$$

Often in the literature a smaller algebra is considered, which only contains q^h for $h \in \frac{1}{2}Q^\vee$, in which case the definition is given in terms of generators $K_i = q^{h_i}$. Also, one can get many different forms by replacing our e_i by $q^{\pm d_i h_i/2} e_i$ or replacing q by q^{-1}; however, essentially all these definitions are equivalent (as long as they are over rational functions of q and not over formal power series).

Many of the facts about $U\mathfrak{g}$ can be generalized to $U_q\mathfrak{g}$ without difficulties. For example, $U_q\mathfrak{g}$ has a polarization: $U_q\mathfrak{g} = U^+ \cdot U^0 \cdot U^-$. It also admits a ($\mathbb{C}_q$-linear) Chevalley involution ω which is defined by $e_i \mapsto -f_i, f_i \mapsto -e_i, q^h \mapsto q^{-h}$; this is an algebra automorphism and coalgebra antiautomorphism. We will also need the following property of this involution: $S\omega = \omega S^{-1}$.

Similarly, the representation theory of $U_q\mathfrak{g}$ is quite parallel to the classical case (see [L1]), the only difference being that all the representations must be considered as linear spaces over $\mathbb{C}_q$. Namely, for every $\lambda \in \mathfrak{h}^*$ we can define Verma module M_λ in a way similar to that of Section 1.1. If λ is not an integer weight, then this module should be considered as a linear space over the field $\mathbb{C}_{q,q^\lambda}$ of rational functions in $q^{1/2N}, t_i = q^{\langle \lambda, \alpha_i^\vee \rangle/2N}$. Again, each M_λ has a unique (non-trivial) irreducible quotient L_λ, and L_λ is finite-dimensional iff $\lambda \in P^+$. Verma modules and all its subfactors (in particular, L_λ) have weight decomposition, and character (and therefore, dimension) of L_λ is the same as in the classical case. More generally, the same is true for the whole category $\mathcal{O}$: every $\mathfrak{g}$-module from category $\mathcal{O}$ has a q-analogue.

On each M_λ one can define Shapovalov form $(\ ,\ )$ such that $(xv, v') = (v, \omega S(x)v')$; this form is defined uniquely up to a factor, it is symmetric and its restriction to L_λ is non-degenerate.

Though it is not literally true that every finite-dimensional irreducible module is of the form L_λ, it is almost so: if we restrict ourselves to consideration of the modules with weight decomposition: $V = \bigoplus V[\lambda], q^h|_{V[\lambda]} = q^{\langle h,\lambda \rangle} \operatorname{Id}_{V[\lambda]}$ then the category of such finite-dimensional representations of $U_q\mathfrak{g}$ is equivalent to that of representations of $\mathfrak{g}$ as a tensor category; in particular, every finite-dimensional representation is completely reducible, the only irreducible finite-dimensional representations are $L_\lambda, \lambda \in P^+$, and $L_\lambda \otimes L_\mu = \sum N^\nu_{\lambda\mu} L_\nu$, where the multiplicities $N^\nu_{\lambda\mu}$ are the same as in classical case. In this whole work we only consider representations with weight decomposition.

However, there are several very important differences between the representations of $U_q\mathfrak{g}$ and of $\mathfrak{g}$. First of all, we have a different notion of dual representation. Namely, for every finite-dimensional representation V of $U_q\mathfrak{g}$ we define the action of $U_q\mathfrak{g}$ on the space V^* of linear functionals on V by the rule $\langle xv^*, v \rangle = \langle v^*, S(x)v \rangle$ for $v \in V, v^* \in V^*, x \in U_q\mathfrak{g}$, where S is the antipode in $U_q\mathfrak{g}$. This endows V^* with the structure of $U_q\mathfrak{g}$ representation which we will call right dual to V. In a similar way, the left dual ${}^*V$ is the representation of $U_q\mathfrak{g}$ in the space of linear functionals on V defined by $\langle xv^*, v \rangle = \langle v^*, S^{-1}(x)v \rangle$. Then the following natural pairings and embeddings are $U_q\mathfrak{g}$-homomorphisms (warning: the order is important here!):

$$\begin{aligned} &V \otimes {}^*V \to \mathbb{C}_q, \quad V^* \otimes V \to \mathbb{C}_q \\ &\mathbb{C}_q \to V \otimes V^*, \quad \mathbb{C}_q \to {}^*V \otimes V \end{aligned} \tag{1.2.5}$$

Note that V^* and ${}^*V$, considered as two structures of representation of $U_q\mathfrak{g}$ on the same vector space, don't coincide, but they are isomorphic. Namely, $q^{-2\rho} : {}^*V \to V^*$ is an isomorphism of $U_q\mathfrak{g}$-modules (recall that we identified $\mathfrak{h}$ and $\mathfrak{h}^*$ using the form $(\, , \,)'$, so $2\rho \in Q \subset Q^\vee$). This follows from the relation $S^2(x) = q^{-2\rho} x q^{2\rho}$, which can be easily checked (it suffices to check it for the generators e_i, f_i).

If $V = L_\lambda$ is an irreducible finite-dimensional representation, then so is V^*: $L_\lambda^* \simeq L_{\lambda^*}$.

Second, unlike the classical case, if V, W are finite-dimensional representations of $U_q\mathfrak{g}$ then $V \otimes W$ and $W \otimes V$ are isomorphic, but the isomorphism is non-trivial. More precisely (see [Dr1]), there exists a universal R-matrix $\mathcal{R}$ which is an element of a certain completion of $U_q\mathfrak{g} \otimes U_q\mathfrak{g}$ such that

$$\check{R}_{V,W} = P \circ \pi_V \otimes \pi_W(\mathcal{R}) \colon V \otimes W \to W \otimes V \tag{1.2.6}$$

is an isomorphism of representations. Here P is the transposition: $Pv \otimes w = w \otimes v$. Also, it is known that $\mathcal{R}$ has the following form:

$$\begin{aligned} &\mathcal{R} = q^{-\sum a_i \otimes a_i} \mathcal{R}^*, \quad \mathcal{R}^* \in U^+ \hat{\otimes} U^- \\ &(\epsilon \otimes 1)(\mathcal{R}^*) = (1 \otimes \epsilon)(\mathcal{R}^*) = 1 \otimes 1, \end{aligned} \tag{1.2.7}$$

where a_i is an orthonormal basis in $\mathfrak{h}$ with respect to $(\, , \,)'$. As we said, $\mathcal{R}$ does not lie in the tensor square of $U_q\mathfrak{g}$ but in its certain completion; however, for any pair of finite-dimensional representations V, W of $U_q\mathfrak{g}$ the operator $\pi_V \otimes \pi_W(\mathcal{R})$ is well-defined (this is where we need fractional powers of q in the definition of $\mathbb{C}_q$).

Finally, if V is a representation of $U_q\mathfrak{g}$ let us consider the action of $U_q\mathfrak{g}$ in V given by $\pi_{V^\omega}(x) = \pi(\omega x)$, where ω is the Chevalley involution defined above. We denote V endowed with this action by V^ω. One can easily check that if V is finite-dimensional then $V^\omega \simeq {}^*V$ (which is, of course, equivalent to saying that $V^\omega \simeq V^*$): if $(\, , \,)$ is Shapovalov form on V then $x \otimes y \mapsto (x, y)$ is a homomorphism of $U_q\mathfrak{g}$-modules $V \otimes V^\omega \to \mathbb{C}_q$. Note also that $(V \otimes W)^\omega = W^\omega \otimes V^\omega$ and that if $\Phi \colon V \to W$ is an intertwiner then Φ is also an intertwiner considered as a map $V^\omega \to W^\omega$.

In certain instances, the structure of $U_q\mathfrak{g}$ is even simpler than that of $U\mathfrak{g}$. A good example is the following description of the center of $U_q\mathfrak{g}$, due to Drinfeld ([Dr2]) (a similar construction was independently proposed by N. Reshetikhin [R]).

Theorem 1.2.1.

(1) *Let V be a finite-dimensional representation of $U_q\mathfrak{g}$. Define*

$$c_V = (\mathrm{Id} \otimes \mathrm{Tr}_V) \left(\mathcal{R}^{21} \mathcal{R} (1 \otimes q^{-2\rho}) \right), \tag{1.2.8}$$

where $\mathcal{R}^{21} = P(\mathcal{R}), P(x \otimes y) = y \otimes x$. *Then* c_V *is a central element in* $U_q\mathfrak{g}$.

(2)

$$c_V|_{M_\lambda} = \operatorname{ch} V(q^{-2(\lambda+\rho)}) \operatorname{Id}_{M_\lambda},$$

where $f(q^{2\lambda})$ *for* $f \in \mathbb{C}[P]$ *denotes the polynomial in* $q^{1/2N}$ *obtained by replacing each* e^μ *by* $q^{2(\lambda,\mu)'}$.

(3) *The elements* $c_{L_\lambda}, \lambda \in P^+$ *form a basis in* $\mathcal{Z}(U_q\mathfrak{g})$.

Proof. (1) This is based on the following statement (see [Dr2]): if $\theta: U_q\mathfrak{g} \to \mathbb{C}_q$ is such that $\theta(xy) = \theta(yS^2(x))$ then the element $c_\theta = (\operatorname{Id} \otimes \theta)(\mathcal{R}^{21}\mathcal{R})$ is central. On the other hand, we know that $S^2(x) = q^{-2\rho}xq^{2\rho}$, so $\theta(x) = \operatorname{Tr}|_V(xq^{-2\rho})$, where V is any finite-dimensional representation of $U_q\mathfrak{g}$, satisfies $\theta(xy) = \theta(yS^2(x))$.

(2) Let v_λ be a highest-weight vector in M_λ; let us calculate $c_V v_\lambda$. Let $w \in V[\mu]$. Then (1.2.7) implies

$$\mathcal{R}^{21}\mathcal{R}(v_\lambda \otimes w) = q^{-2(\lambda,\mu)'} v_\lambda \otimes w + \sum v'_i \otimes w'_i,$$

where wt $w'_i < \mu$. Thus, $c_r v_\lambda = (\sum_\mu (\dim V[\mu]) q^{-2(\lambda,\mu)'} q^{-2(\rho,\mu)'}) v_\lambda$, where the sum is taken over all the weights of V.

(3) This is based on the quantum version of Harish-Chandra isomorphism. Namely, if C is a central element then it acts in Verma module M_λ by some constant; denote it $\gamma(C)(\lambda+\rho)$. It is easy to see that $\gamma(C)(\lambda+\rho)$ can be uniquely written in the form $f(q^{\lambda+\rho})$ for some $f \in \mathbb{C}_q[\frac{1}{2}P^\vee]^W$. Moreover, it is known (see [T]) that γ is an isomorphism $\mathcal{Z}(U_q\mathfrak{g}) \simeq \mathbb{C}_q[2P]^W$. Since characters of irreducible representations form a basis in $\mathbb{C}_q[P]^W$, we obtain the statement (3). □

Remark. Note that though $\mathcal{R}$ is defined only in some completion of $U_q\mathfrak{g}^{\otimes 2}$, the central elements we constructed belong to $U_q\mathfrak{g}$ itself.

CHAPTER II

GENERALIZED CHARACTERS

This chapter is the heart of the whole dissertation. In this chapter we define for every $\mathfrak{g}$-intertwining operator Φ a certain weighted trace χ_Φ, which we call *generalized character*, which is justified by the fact that if $\Phi : V \to V$ is the identity operator then χ_Φ is the usual character of V. We also study basic properties of these characters: namely, we prove an orthogonality theorem for these characters, and show that they are eigenfunctions of a large family of commuting differential operators. We also describe the q-analogue of these generalized characters, which is defined in a similar way, but $\mathfrak{g}$ must be replaced with the quantum group $U_q\mathfrak{g}$.

All of constructions of this chapter belong to the author, P. Etingof and in some part I. Frenkel, see [EK1, EK2, EK4, EFK].

2.1 Weighted traces of intertwiners and orthogonality theorem

In this section we define the basic object of our study: the weighted traces of intertwining operators, which we call generalized characters. **All the definitions and theorems of this section apply equally to the quantum group $U_q\mathfrak{g}$ and the usual universal enveloping algebra $U\mathfrak{g}$.** To avoid unnecessary repetitions, we formulate all the results for $U_q\mathfrak{g}$; however, the reader should keep in mind that all of the statements also hold if one replaces $U_q\mathfrak{g}$ by $U\mathfrak{g}$, and $\mathbb{C}_q$ by $\mathbb{C}$.

Definition 2.1.1. Let V be a $U_q\mathfrak{g}$-module from category $\mathcal{O}$, U – an arbitrary $U_q\mathfrak{g}$-module with weight decomposition (not necessarily finite-dimensional), and let $\Phi \colon V \to V \otimes U$ be a non-zero intertwining operator.

A generalized character $\chi_\Phi \in \overline{\mathbb{C}[P]} \otimes U$ is defined by

$$\chi_\Phi = \sum_{\lambda \in P} e^\lambda \operatorname{Tr}|_{V[\lambda]}(\Phi). \tag{2.1.1}$$

In particular, if V is finite-dimensional then $\chi_\Phi \in \mathbb{C}[P] \otimes U$.

Example. Let $U = \mathbb{C}$ be the trivial representation of $U_q\mathfrak{g}$, and let $\Phi : V \to V$ be the identity operator. Then χ_{Id} is the usual character of V, which justifies the name "generalized character".

For finite-dimensional V, we can consider these traces as functions on $\mathfrak{h}$ with values in U by the rule $e^\lambda(h) = e^{2\pi \mathrm{i}\langle\lambda,h\rangle}$. This is equivalent to letting

$$\chi_\Phi(h) = \mathrm{Tr}_V(\Phi e^{2\pi \mathrm{i} h}), \tag{2.1.2}$$

where $e^{2\pi \mathrm{i} h}$ is an operator in V defined by $e^{2\pi \mathrm{i} h}|_{V[\lambda]} = e^{2\pi \mathrm{i}\langle h,\lambda\rangle}\,\mathrm{Id}_{V[\lambda]}$.

Using the fact that Φ preserves weight, it is easy to see that in fact χ_Φ takes values in the zero-weight subspace $U[0]$. Also, for $V = L_\lambda, \lambda \in P^+$ the highest term of χ_Φ is ue^λ and the lowest term is $u'e^{-\lambda^*}$ for some $u, u' \in U$.

Note that in the quantum case (i.e., for $U_q\mathfrak{g}$ rather than $U\mathfrak{g}$) χ_Φ has no Weyl group symmetry if U is a nontrivial representation (see a counterexample in Theorem 4.2.2).

Let us recall the well-known results on the orthogonality of (usual) characters. Let us introduce the (q-linear) bar involution in $\mathbb{C}_q[P]$ by $\overline{e^\lambda} = e^{-\lambda}$, and let $[\ \ ]_0 : \mathbb{C}[P] \to \mathbb{C}$ be the constant term: $[\sum c_\lambda e^\lambda]_0 = c_0$. Then it is known that characters of irreducible finite-dimensional representations of $U_q\mathfrak{g}$ (which are the same as for $\mathfrak{g}$) are orthonormal with respect to the following inner product in $\mathbb{C}[P]$:

$$\langle f, g\rangle_1 = \frac{1}{|W|}[f\bar{g}\Delta]_0, \tag{2.1.3}$$

where

$$\Delta = \delta\bar{\delta} = \prod_{\alpha\in R}(1 - e^\alpha). \tag{2.1.4}$$

It turns out that a similar statement holds for generalized characters:

Theorem 2.1.2. (Orthogonality theorem for generalized characters) *Let $\lambda, \mu \in P^+$, and let U be a finite-dimensional representation of $U_q\mathfrak{g}$. Let $\Phi_\lambda : L_\lambda \to L_\lambda \otimes U, \Phi_\mu : L_\mu \to L_\mu \otimes U$ be $U_q\mathfrak{g}$-intertwiners, and $\lambda \neq \mu$. Then the generalized characters $\chi_1 = \chi_{\Phi_\lambda}, \chi_2 = \chi_{\Phi_\mu}$ are orthogonal with respect to the following inner product: $\langle f, g\rangle_1 = \frac{1}{|W|}[(f, \bar{g})_U\Delta]_0$, where $(\cdot,\cdot)_U$ is the Shapovalov form in U and all the other notations are as before.*

Proof. As was explained above, we can as well consider Φ_μ as an intertwiner $L^\omega_\mu \to U^\omega \otimes L^\omega_\mu$. This implies that $(\chi_1, \overline{\chi_2})_U = \chi_\Psi$ (note the change of sign of h in the second factor!), where the intertwiner $\Psi : L_\lambda \otimes L^\omega_\mu \to L_\lambda \otimes L^\omega_\mu$ is defined as the following composition

$$(2.1.5) \qquad L_\lambda \otimes L_\mu^\omega \xrightarrow{\Phi_\lambda \otimes \Phi_\mu^\omega} L_\lambda \otimes U \otimes U^\omega \otimes L_\mu^\omega \xrightarrow{\mathrm{Id} \otimes (\cdot,\cdot)_U \otimes \mathrm{Id}} L_\lambda \otimes L_\mu^\omega.$$

Since $L_\lambda \otimes L_\mu^\omega = \bigoplus N_\nu L_\nu$, we see that $(\chi_1, \overline{\chi_2})_U$ is a linear combination of usual characters $\operatorname{ch} L_\nu$. But since these characters are the same as for $\mathfrak{g}$, we know that $\frac{1}{|W|}[\chi_\nu \Delta]_0 = \delta_{\nu,0}$. On the other hand, it is known that if $\lambda \neq \mu$ then the decomposition of $L_\lambda \otimes L_\mu^\omega$ does not contain the trivial representation (i.e. $N_0 = 0$); thus, in this case χ_1 and χ_2 are orthogonal. $\square$

REMARK. Note that the calculation of the norm $\langle \chi_\Phi, \chi_\Phi \rangle_1$ (which is trivial for $U = \mathbb{C}$) in general case is very complicated, since it requires calculation of restriction of the intertwiner Ψ defined above to the trivial subrepresentation in $L_\lambda \otimes L_\lambda^\omega$. In some simplest cases (where the space of intertwiners is one-dimensional) it can be calculated explicitly (we will do it in Chapter 5), and the answer is quite non-trivial; it coincides with so-called Macdonald inner product identities for root system A_n, see [M2].

2.2 Center of $U\mathfrak{g}$ and commuting differential operators

In this section we consider some properties of generalized characters for $U\mathfrak{g}$, following the paper of Etingof [E1]. In this section, the words "representation" and "intertwiner" stand for representation of $\mathfrak{g}$ and $\mathfrak{g}$-intertwiner. In this section we always consider the case when V, U are finite-dimensional, though in fact all the results can be easily generalized (see remark in the end of this section).

Let $\mathbb{C}[P](e^\alpha - 1)^{-1}$ denote the ring of fractions $\frac{f}{g}$, where $f, g \in \mathbb{C}[P]$ and g is a product of linear terms of the form $e^\alpha - 1, \alpha \in R$. Let us introduce the following ring of differential operators:

$$DO = \{\text{differential operators on } \mathfrak{h} \text{ with coefficients from } \mathbb{C}[P](e^\alpha - 1)^{-1}\}.$$

These operators can be also considered formally, as derivations of the ring of fractions $\mathbb{C}[P](e^\alpha - 1)^{-1}$. In particular, we will use the following differential operators:

$\Delta_\mathfrak{h}$ – Laplace operator in $\mathfrak{h}$ normalized so that $\Delta_\mathfrak{h} e^\lambda = (\lambda, \lambda) e^\lambda$;

$\partial_\alpha, \alpha \in \mathfrak{h}^*$ – differentiation along the vector $\frac{1}{2\pi \mathrm{i}} h_\alpha$, so that $\partial_\alpha e^\lambda = (\alpha, \lambda) e^\lambda$.

THEOREM 2.2.1. *Let us denote by I the left ideal in $U\mathfrak{g}$ generated by $\mathfrak{h}$. Then for any $u \in U\mathfrak{g}$ there exists a unique differential operator $D_u \in DO \otimes U\mathfrak{g}/I$, such that for any intertwiner $\Phi : V \to V \otimes U$ we have*

$$(2.2.1) \qquad \operatorname{Tr}_V(\Phi u e^{2\pi \mathrm{i} h}) = D_u \operatorname{Tr}_V(\Phi e^{2\pi \mathrm{i} h}).$$

Note that $D_u f$ is well defined for any function f with values in $U[0]$.

We do not give the proof of this theorem (which can be found in [E1]) here, since the proof is quite similar to the proof in the q-case, which we give later (see Theorem 2.3.1). The main idea of the proof will be quite clear from the example of Casimir element, which we discuss below.

COROLLARY 2.2.2. *Let c be a central element: $c \in \mathcal{Z}(U\mathfrak{g})$, V, U be finite-dimensional representations of $\mathfrak{g}$ such that c acts in V by multiplication by a constant $c(V)$, and $\Phi : V \to V \otimes U$ be a non-zero intertwiner. Then the generalized character χ_Φ satisfies the following differential equation*

$$D_c \chi_\Phi = c(V)\chi_\Phi, \tag{2.2.2}$$

where the differential operator D_c is defined in Theorem 2.2.1.

PROOF. Let us consider $f(h) = \mathrm{Tr}_V(\Phi c e^{2\pi i h})$. On one hand, since c acts by multiplication by $c(V)$, we see that $f(h) = c(V)\chi_\Phi(h)$. On the other hand, by Theorem 2.2.1, $f(h) = D_c\chi_\Phi$. □

In general, $u \mapsto D_u$ is not an algebra homomorphism. However, it becomes a homomorphism when restricted to the center:

THEOREM 2.2.3. *Let us keep the notations of Theorem 2.2.1. Let $c, c' \in \mathcal{Z}(U\mathfrak{g})$. Then $D_{cc'} = D_c D_{c'}$; thus, $c \mapsto D_c$ is a homomorphism $\mathcal{Z}(U\mathfrak{g}) \to DO \otimes U\mathfrak{g}/I$.*

PROOF. By definition, $D_{cc'}\chi_\Phi = \mathrm{Tr}(\Phi c c' e^{2\pi i h})$. On the other hand, since c is central, Φc is also an intertwiner, and therefore we can write $\mathrm{Tr}(\Phi c c' e^{2\pi i h}) = D_{c'}\chi_{\Phi c} = D_{c'} D_c \chi_\Phi$.

Let us apply the construction of Theorem 2.2.1 to the Casimir element C defined in Section 1.1. In this case the answer can be written down explicitly:

PROPOSITION 2.2.4.

1. *Let C be the Casimir element. Then*

$$\begin{aligned} D_C &= \Delta_{\mathfrak{h}} - 2\sum_{\alpha\in R^+} \frac{e_\alpha f_\alpha}{(e^{\alpha/2} - e^{-\alpha/2})^2} - \sum_{\alpha\in R^+} \frac{1+e^\alpha}{1-e^\alpha}\partial_\alpha \\ &= \delta^{-1} \circ \left(\Delta_{\mathfrak{h}} - 2\sum_{\alpha\in R^+} \frac{e_\alpha f_\alpha}{(e^{\alpha/2} - e^{-\alpha/2})^2} - (\rho,\rho)\right) \circ \delta, \end{aligned} \tag{2.2.3}$$

   where δ is Weyl denominator.

2. *Let $\Phi : L_\lambda \to L_\lambda \otimes U$ be an intertwiner. Then the corresponding generalized character χ_Φ satisfies the following equation:*

$$\left(\Delta_{\mathfrak{h}} - 2\sum_{\alpha\in R^+} \frac{e_\alpha f_\alpha}{(e^{\alpha/2} - e^{-\alpha/2})^2}\right)(\delta\chi_\Phi) = (\lambda+\rho, \lambda+\rho)\delta\chi_\Phi. \tag{2.2.4}$$

PROOF. Recall that the Casimir element was defined by: $C=\sum_{\alpha\in R^+}(e_\alpha f_\alpha+f_\alpha e_\alpha)+\sum x_l^2$, where $e_\alpha\in\mathfrak{g}_\alpha, f_\alpha\in\mathfrak{g}_{-\alpha}$ are such that $(e_\alpha,f_\alpha)=1$, and x_l is an orthonormal basis in $\mathfrak{h}$ with respect to (,). Therefore, (1) follows from the following identities:

$$\operatorname{Tr}_V(\Phi(\sum x_l^2)e^{2\pi\mathrm{i}h})=\Delta_{\mathfrak{h}}\chi_\Phi, \tag{2.2.5}$$

$$\operatorname{Tr}_V(\Phi e_\alpha f_\alpha e^{2\pi\mathrm{i}h})=-\frac{e_\alpha f_\alpha}{(e^{\alpha/2}-e^{-\alpha/2})^2}\chi_\Phi-\frac{e^\alpha}{1-e^\alpha}\partial_\alpha\chi_\Phi, \tag{2.2.6}$$

$$\operatorname{Tr}_V(\Phi f_\alpha e_\alpha e^{2\pi\mathrm{i}h})=-\frac{f_\alpha e_\alpha}{(e^{\alpha/2}-e^{-\alpha/2})^2}\chi_\Phi-\frac{1}{1-e^\alpha}\partial_\alpha\chi_\Phi. \tag{2.2.7}$$

The first of these identities is immediate corollary of the definition of χ_Φ. Let us prove the second one. Denote $X=\operatorname{Tr}_V(\Phi e_\alpha f_\alpha e^{2\pi\mathrm{i}h})$. Then, using the identities $\Phi e_\alpha=(1\otimes e_\alpha+e_\alpha\otimes 1)\Phi$, $e^{2\pi\mathrm{i}h}e_\alpha=e^{2\pi\mathrm{i}\langle\alpha,h\rangle}e_\alpha e^{2\pi\mathrm{i}h}$ and the cyclic property of the trace, we can write:

$$\begin{aligned}X=&\operatorname{Tr}_V(\Phi e_\alpha f_\alpha e^{2\pi\mathrm{i}h})=e_\alpha\operatorname{Tr}(\Phi f_\alpha e^{2\pi\mathrm{i}h})+\operatorname{Tr}(\Phi f_\alpha e^{2\pi\mathrm{i}h}e_\alpha)\\ =&e_\alpha\operatorname{Tr}(\Phi f_\alpha e^{2\pi\mathrm{i}h})+e^{2\pi\mathrm{i}\langle\alpha,h\rangle}\operatorname{Tr}(\Phi f_\alpha e_\alpha e^{2\pi\mathrm{i}h})\\ =&e_\alpha\operatorname{Tr}(\Phi f_\alpha e^{2\pi\mathrm{i}h})+e^\alpha\operatorname{Tr}(\Phi(e_\alpha f_\alpha-h_\alpha)e^{2\pi\mathrm{i}h})\\ =&e_\alpha\operatorname{Tr}(\Phi f_\alpha e^{2\pi\mathrm{i}h})+e^\alpha(X-\partial_\alpha\chi_\Phi),\end{aligned}$$

(recall that e^α denotes both the element of $\mathbb{C}[P]$ and a function on $\mathfrak{h}$ given by $e^\alpha(h)=e^{2\pi\mathrm{i}\langle\alpha,h\rangle}$). Therefore,

$$X=\frac{1}{1-e^\alpha}\left(e_\alpha\operatorname{Tr}(\Phi f_\alpha e^{2\pi\mathrm{i}h})-e^\alpha\partial_\alpha\chi_\Phi\right).$$

Similar arguments show that

$$\operatorname{Tr}(\Phi f_\alpha e^{2\pi\mathrm{i}h})=\frac{f_\alpha}{1-e^{-\alpha}}\chi_\Phi,$$

which gives us (2.2.6). Proof of (2.2.7) is quite similar.

This proves the first identity in (1); the second identity can be checked by direct calculation or deduced from Proposition 3.1.1 (see below).

(2) obviously follows from (1) and Corollary 2.2.2 along with the fact that $C|_{L_\lambda}=(\lambda,\lambda+2\rho)$.

EXAMPLE. Let $U=\mathbb{C}$ be the trivial representation, $\Phi=\mathrm{Id}_{L_\lambda}$. Then Proposition 2.2.4 implies that the character $\operatorname{ch}L_\lambda$ satisfies the following equation:

$$\Delta(\delta\operatorname{ch}L_\lambda)=(\lambda+\rho,\lambda+\rho)\delta\operatorname{ch}L_\lambda; \tag{2.2.8}$$

in particular,

$$\Delta\delta = (\rho, \rho)\delta. \tag{2.2.9}$$

These equations are well-known and have a nice geometric interpretation – as the heat equation on the compact Lie group G associated with $\mathfrak{g}$.

Remarks.

(1) Note that the theorems of this section can be easily generalized to the case when V is from category $\mathcal{O}$. In this case, we must consider the differential operators with coefficients from the ring $\overline{\mathbb{C}[P]}$: since for $\alpha \in R^-$, $(1-e^\alpha)^{-1} = 1 + e^\alpha + e^{2\alpha} + \dots$ converges in $\overline{\mathbb{C}[P]}$, there is no need to extend this ring.

(2) The equations (2.2.3), (2.2.4) are special cases of the formulas for radial part of Laplace operator in the theory of μ-special functions, due to Harish-Chandra (see [W, Chapter 9]). We discuss the relation between generalized charcters and spherical functions in Section 2.4 below.

2.3 Center of $U_q\mathfrak{g}$ and difference operators

In this section we discuss the analogue of the construction of Section 2.2 in the q-case. The construction is quite parallel to the classical case, but the differential operators should be replaced by difference operators. In this section, the words "representation" and "intertwiner" stand for representation of $U_q\mathfrak{g}$ and $U_q\mathfrak{g}$-intertwiner, and the ground field is the field $\mathbb{C}_q$ of rational functions in $q^{1/2N}$ (see Section 1.2)

Introduce the following ring of difference operators, acting on the functions $f \in \mathbb{C}_q[P]$:

$$DO_q = \{D = \sum_{\alpha\in\frac{1}{2}P^\vee} a_\alpha T_\alpha |\ \text{almost all } a_\alpha = 0\},$$

where $T_\alpha e^\lambda = q^{\langle\alpha,\lambda\rangle} e^\lambda$, and $a_\alpha \in \mathbb{C}_q[P](q^m e^\mu - 1)^{-1}$ are fractions of the form $\frac{f}{g}, f, g \in \mathbb{C}_q[P]$ such that g is a product of factors of the form $q^m e^\mu - 1$ for some $\mu \in Q, m \in \mathbb{Z}$.

Note that if we consider elements of $\mathbb{C}_q[P]$ as functions on $\mathfrak{h}$ then T_α can be rewritten as follows: $(T_\alpha f)(h) = f(h + \alpha\frac{\log q}{2\pi \mathrm{i}})$; however, to avoid difficulties with proper definition of $\log q$ when q is a formal variable, we prefer not to use this language and by "function" we always mean an element of $\mathbb{C}_q[P]$ (or a its field of fractions). However, to simplify the notations we sometimes will write the generalized character χ_Φ as $\mathrm{Tr}(\Phi e^{2\pi \mathrm{i} h})$ meaning by this the element of $\mathbb{C}_q[P]$ defined by (2.1.1). One can check that in fact all our arguments can be carried in this formal language.

As in the previous section, let V, U be finite-dimensional representations of $U_q\mathfrak{g}$.

Theorem 2.3.1. *Let I be the left ideal in $U_q\mathfrak{g}$ generated by $q^h - 1$. Then for any $u \in U_q\mathfrak{g}$ there exists a unique difference operator $D_u \in DO_q \otimes U_q\mathfrak{g}/I$ such that for any intertwiner $\Phi : V \to V \otimes U$ we have*

$$\mathrm{Tr}_V(\Phi u e^{2\pi \mathrm{i} h}) = D_u \,\mathrm{Tr}_V(\Phi e^{2\pi \mathrm{i} h}). \tag{2.3.1}$$

Again, $D_u f$ is well defined for any function f with values in $U[0]$.

Proof. Without loss of generality we can assume that u is a monomial in the generators e_i, f_i, q^h of the form $u = u^- u^0 u^+$, $u^\pm \in U^\pm, u^0 \in U^0$. Define sdeg $u =$ deg $u^+ -$ deg u^-, where deg $e_i = -$deg $f_i = 1$. We prove the theorem by induction in sdeg u.

If sdeg $u = 0$ then $u = u^0 = q^\alpha$ for some $\alpha \in \frac{1}{2}P^\vee$. Then it follows immediately from the definition that $D_u = T_\alpha$, so the theorem holds.

Let us make the induction step. Assume that sdeg $u > 0$; then either deg $u^+ \neq 0$ or deg $u^- \neq 0$. We can assume that $u^- \in U[\mu], \mu \in -Q^+, \mu \neq 0$.

Since Φ is an intertwiner, $\mathrm{Tr}(\Phi u^- u^0 u^+ e^{2\pi \mathrm{i} h}) = \mathrm{Tr}(\Delta(u^-)\Phi u^0 u^+ e^{2\pi \mathrm{i} h})$. From the definition of comultiplication one easily sees that $\Delta(u^-) = u^- \otimes q^{\mu/2} + \sum u_j \otimes v_j$ for some $u_j, v_j \in U^0 U^-$ such that sdeg $(u_j u_0 u^+) <$ sdeg u. Thus,

$$\mathrm{Tr}(\Phi u e^{2\pi \mathrm{i} h}) = q^{\mu/2}\,\mathrm{Tr}(\Phi u^0 u^+ e^{2\pi \mathrm{i} h} u^-) + \sum v_j \,\mathrm{Tr}(\Phi u^0 u^+ e^{2\pi \mathrm{i} h} u_j).$$

Since commuting with $e^{2\pi \mathrm{i} h}$ does not change sdeg u_j, by the induction assumption we can write

$$\begin{aligned}\mathrm{Tr}(\Phi u e^{2\pi \mathrm{i} h}) =& q^{\mu/2}\,\mathrm{Tr}(\Phi u^0 u^+ e^{2\pi \mathrm{i} h} u^-) + D'\,\mathrm{Tr}(\Phi e^{2\pi \mathrm{i} h})\\ =& q^{(\mu,\mu)'/2} e^\mu \,\mathrm{Tr}(\Phi u^0 u^+ u^- e^{2\pi \mathrm{i} h}) + D'\,\mathrm{Tr}(\Phi e^{2\pi \mathrm{i} h})\\ =& q^{(\mu,\mu)'/2} e^\mu \,\mathrm{Tr}(\Phi (u + [u^0 u^+, u^-]) e^{2\pi \mathrm{i} h}) + D'\,\mathrm{Tr}(\Phi e^{2\pi \mathrm{i} h})\end{aligned}$$

for some $D' \in DO_q \otimes U_q\mathfrak{g}$. Since sdeg of all terms in $[u^0 u^+, u^-]$ is less than sdeg $u^- u^0 u^+$, we can again apply induction assumption and get

$$\mathrm{Tr}(\Phi u e^{2\pi \mathrm{i} h}) = \frac{1}{1 - q^{(\mu,\mu)'/2} e^\mu} D''\,\mathrm{Tr}(\Phi e^{2\pi \mathrm{i} h}).$$

This proves the existence part of the theorem. Uniqueness follows from the following lemma:

Lemma 2.3.2. *Let us fix a finite-dimensional $U_q\mathfrak{g}$-module U. Suppose that $D \in DO \otimes \mathrm{Hom}(U[0], U[\mu])$ is such that for any $U_q\mathfrak{g}$-intertwiner $\Phi : V \to V \otimes U, V$ – finite-dimensional $U_q\mathfrak{g}$-module we have $D\chi_\Phi = 0$. Then $D = 0$.*

Proof of the lemma. Let us assume that $D \neq 0$. Multiplying D by a suitable element from $\mathbb{C}_q[P]$ we can assume that D has polynomial coefficients:

$D = \sum_{\lambda \in P} e^{\lambda} D_{(\lambda)}$, $D_{(\lambda)}$ being difference operators with constant matrix-valued coefficients. Let us take the maximal (with respect to the standard order in P) λ such that $D_{(\lambda)} \neq 0$. Then if we have a generalized character χ such that $\chi = e^{\mu} u +$ *lower order terms* then, taking the highest term of $D\chi$, we see that $D_{(\lambda)}(e^{\mu} u) = 0$. On the other hand, if we take μ such that $\langle \mu, \alpha_i^{\vee} \rangle \gg 0$ then for every $u \in U[0]$ there exists a non-zero intertwiner $\Phi \colon L_{\mu} \to L_{\mu} \otimes U$ such that the corresponding generalized character has the form $\chi_{\Phi} = e^{\mu} u +$ lower order terms. Thus $D_{(\lambda)}(e^{\mu} u) = 0$ for all $\mu \gg 0, u \in U[0]$. It is easy to show that it is only possible if $D_{(\lambda)} = 0$, which contradicts the assumption $D_{(\lambda)} \neq 0$. $\square$

PROPOSITION 2.3.3. *Let us keep the notations of Theorem* 2.3.1. *Then* $c \mapsto D_c$ *is an algebra homomorphism of* $\mathcal{Z}(U_q\mathfrak{g})$ *to* $DO_q \otimes U_q\mathfrak{g}[0]/I$.

PROOF. The same as in the classical case (Theorem 2.2.3)

Unlike the classical case, in the quantum case we have an explicit construction of central elements (see Theorem 1.2.1). Applying the previous construction to those central elements, we get the following theorem:

THEOREM 2.3.4. *Let* V *be a finite-dimensional representation of* $U_q\mathfrak{g}$, c_V *– the corresponding central element in* $U_q\mathfrak{g}$ *(see Theorem* 1.2.1*), and* D_{c_V} *– the corresponding difference operator constructed in Theorem* 2.3.1. *Let* $\Phi : L_{\lambda} \to L_{\lambda} \otimes U$ *be a* $U_q\mathfrak{g}$*-intertwiner, and* χ_{Φ}*– the corresponding generalized character. Then* χ_{Φ} *satisfies the following difference equation:*

$$D_{c_V} \chi_{\Phi} = \operatorname{ch} V(q^{-2(\lambda+\rho)}) \chi_{\Phi}. \tag{2.3.2}$$

PROOF. The same as in classical case (see Corollary 2.2.2); the value of c_V in L_{λ} is taken from Theorem 1.2.1.

REMARK. As in the classical case, all the constructions of this section can be easily generalized to the case where V is an arbitrary module from category $\mathcal{O}$.

2.4 Generalized characters as spherical functions

In this section we show that generalized characters for $\mathfrak{g}$ can be interpreted as spherical functions on the space $G \times G/G$. The constructions of this section are due to the author, I. Frenkel and P. Etingof (see [EFK]).

As before, let $\mathfrak{g}$ be a simple Lie algebra over $\mathbb{C}$, and let G be the compact real simply connected Lie group corresponding to $\mathfrak{g}$. It is known that every finite-dimensional complex representation of $\mathfrak{g}$ can be lifted to G. Also, it is known that every finite-dimensional representation V of G is unitary: there exists a positive definite hermitian form $(\, , \,)_V$ on V such that $(gv_1, gv_2)_V = (v_1, v_2)_V$.

Let us fix some finite-dimensional representation U of $\mathfrak{g}$. Let $C^{\infty}(G, U)$ be the space of all smooth functions on G with values in U. We can define a (hermitian) inner product in $C^{\infty}(G, U)$ by

$$(f_1, f_2) = \int_G (f_1(g), f_2(g))_U \, dg, \tag{2.4.1}$$

where $(\ ,\ )_U$ is the inner product in U and dg is the Haar measure on G. It is easy to see that the inner product defined by (2.4.1) is positive-definite. We denote by $L^2(G, U)$ the closure of the space $C^\infty(G, U)$ with respect to the norm $\|f\| = \sqrt{(f, f)}$.

Definition 2.4.1. A function $f \in C^\infty(G, U)$ is called *equivariant* (notation: $f \in C^\infty(G, U)^G$) if for every $g, x \in G$ we have

$$f(gxg^{-1}) = gf(x). \tag{2.4.2}$$

The same applies to $f \in L^2(G, U)$.

This definition can be rewritten as follows. Let us consider the group $G \times G$, and let $G_d \subset G \times G$ be the diagonal subgroup. Consider the functions $f : G \times G \to U$ such that for every $k \in G_d, x \in G \times G$ we have

$$\begin{aligned} f(xk) &= f(x), \\ f(kx) &= kf(x). \end{aligned} \tag{2.4.3}$$

This is a special case of what is called μ-spherical functions (see [W]) on the group $G \times G$ with respect to the subgroup G_d. On the other hand, it is easy to see that

$$f(x, y) \mapsto f(xy^{-1})$$

establishes isomorphism between the space of spherical functions on $G \times G/G_d$ defined by (2.4.3) and the space of equivariant functions in the sense of Definition 2.4.1.

Here is yet another description of the same space. Let $f \in C^\infty(G, U)^G$, and let $u \in U^*$. Define a complex-valued function on G by $f_u(g) = \langle u, f(g)\rangle$. Define the action of G on scalar functions on G by $(T_g f)(x) = f(g^{-1}xg)$. Then it is easy to see that $T_g f_u = f_{gu}$, and thus f_u satisfies the following condition:

(2.4.4)
Under the action of G defined above, f_u spans a finite-dimensional subspace in $C^\infty(G)$, and as a representation of G, this space is isomorphic to U^*.

Vice versa, it is easy to see that every scalar function on G satisfying the condition (2.4.4) can be obtained as $f_u(g)$ for some $f \in C^\infty(G, U)^G, u \in U^*$.

Now, let V be a finite-dimensional representation of G and let $\Phi : V \to V \otimes U$ be a G-intertwiner. Define coresponding generalized character $\chi_\Phi \in C^\infty(G, U)$ by

$$\chi_\Phi(g) = \mathrm{Tr}_V(\Phi g). \tag{2.4.5}$$

Note that for $g = e^{2\pi i h}$ this coincides with the previously given definition of the generalized character (see (2.1.2)).

LEMMA 2.4.2.

1. *For every intertwiner* $\Phi : V \to V \otimes U$, *the generalized character* χ_Φ *defined by* (2.4.5) *is equivariant:* $\chi_\Phi \in C^\infty(G, U)^G$.
2. *Assume that* V *is irreducible. Then* $\chi_\Phi = 0$ *iff* $\Phi = 0$.

PROOF. (1) is quite trivial:

$$\mathrm{Tr}_V(\Phi g x g^{-1}) = \mathrm{Tr}((g \otimes g)\Phi x g^{-1}) = g\,\mathrm{Tr}(\Phi x).$$

(2) Let us consider χ_Φ on the maximal torus, i.e. on the points of the form $e^{2\pi \mathrm{i} h}, h \in \mathfrak{h}_\mathbb{R}$. As was explained in Section 2.1, we can as well consider it as an element of $\mathbb{C}[P] \otimes U$, and $\chi_\Phi = 0$ as a function on $\mathfrak{h}$ iff $\chi_\Phi = 0$ as an element of $\mathbb{C}[P] \otimes U$. Let v_λ be a highest-weight vector in V. Then $\Phi v_\lambda = v_\lambda \otimes u_0 +$ *lower order terms* for some $u_0 \in U[0]$. It is known that $u_0 = 0$ iff $\Phi = 0$. On the other hand, $\chi_\Phi = e^\lambda \otimes u_0 +$ *lower order terms*, which proves the theorem. □

Let $\lambda \in P^+$. Recall that L_λ is the irreducible highest-weight module over $\mathfrak{g}$ with highest weight λ, which in this case is finite-dimensional and thus can be considered as a module over G. Let $H_\lambda = \mathrm{Hom}_G(L_\lambda, L_\lambda \otimes U)$ (note that this space is finite-dimensional). Due to Lemma 2.4.2, we have an injective map

$$H_\lambda \to C^\infty(G, U)^G : \Phi \mapsto \chi_\Phi.$$

We will denote the image of H_λ in $C^\infty(G, U)^G$ also by H_λ.

THEOREM 2.4.3.

1. *For* $\lambda \neq \mu$, H_λ *and* H_μ *are orthogonal with respect to the inner product in* $C^\infty(G, U)$ (*see* (2.4.1)).
2. $$L^2(G, U)^G = \bigoplus_{\lambda \in P^+} H_\lambda.$$

   (*direct sum should be understood in the sense of Hilbert spaces*).

We refer the reader to [EFK] for the proof of this theorem.

EXAMPLE. Let $U = \mathbb{C}$ be the trvial representation. Then Theorem 2.4.3 coincides with well-known Peter-Weyl theorem.

Finally, since the conjugacy classes in G are the same as W-orbits in the maximal torus

$$T = \exp(\mathrm{i}\mathfrak{h}_\mathbb{R}) \simeq \mathfrak{h}_\mathbb{R}/Q^\vee, \quad \mathfrak{h}_\mathbb{R} = \bigoplus \mathbb{R}\alpha_i^\vee, \tag{2.4.6}$$

it is easy to see that every equivariant function on G is uniquely defined by its values on T. More precisely, we have the following isomorphism:

$$C^\infty(G, U)^G \simeq C^\infty(T, U[0])^W, \tag{2.4.7}$$

i.e. functions on T with values in $U[0]$ satisfying $f(wt) = wf(t)$ for every element from the Weyl group.

Thus, restriction of the inner product defined by (2.4.1) to the space of equivariant functions can be rewritten in terms of integral over T. The answer is given by the following theorem:

THEOREM 2.4.4. *If* $f_1, f_2 \in C^\infty(G,U)^G$ *then*

$$(f_1, f_2) = \frac{1}{|W|}\int_T (f_1(t), f_2(t))_U \delta\bar{\delta}\, dt, \tag{2.4.8}$$

where, as before, dt *is the Haar measure on* T*,* $(\ ,\ )_U$ *is the inner product in* U*, and* δ *is the Weyl denominator:*

$$\delta(e^{2\pi \mathrm{i} h}) = \prod_{\alpha\in R^+}\left(e^{\pi\mathrm{i}\langle\alpha,h\rangle} - e^{-\pi\mathrm{i}\langle\alpha,h\rangle}\right),$$

and $\bar{\delta}$ *is the complex conjugate of* δ*.*

REMARK. Again, in the case $U = \mathbb{C}$ this theorem is well known; it was first proved by H. Weyl.

PROOF. It is very easy to check that the scalar function $\psi(g) = (f_1(g), f_2(g))_U$ is conjugation invariant. Thus, the theorem is reduced to the following statement: for every central function ψ on G, we have

$$\int_G \psi(g)\, dg = \frac{1}{|W|}\int_T \psi(t)\delta\bar{\delta}\, dt,$$

which is well-known (essentially, this is the result of Weyl we referred to above). □

Note that this theorem along with the orthogonality statement of Theorem 2.4.3 gives a new proof of the hermitian version of the orthogonality theorem for generalized characters for $\mathfrak{g}$ (Theorem 2.1.2); of course, these arguments wouldn't help in the q-case.

Also, using the description of the space of equivariant functions as scalar-valued functions on G transforming under the conjugation as U^* (see (2.4.4)), it is easy to see that every conjugation-invariant scalar differential operator D on G defines an operator in $C^\infty(G,U)^G$, acting by a scalar in every H_λ (see details in [EFK]). Since the space of such operators is isomorphic to $(U\mathfrak{g})^{\mathfrak{g}} \simeq \mathcal{Z}(U\mathfrak{g})$, this gives a natural action of $\mathcal{Z}(U\mathfrak{g})$ by differential operators on $C^\infty(G,U)^G$. Using the isomorphism (2.4.7), we can rewrite this action in terms of differential operators on T with coefficients from End $U[0]$ (this operation is often referred to as "taking the radial part of the Laplacian"). It can be easily shown that the differential operators appearing in this way coincide with those constructed in Theorem 2.2.1. In particular, the operator D_C given by (2.2.3) is exactly the radial part of the second order Laplace-Beltrami operator Δ_G. We refer the reader to [EFK] for more detailed information.

Note that it would be quite difficult to calculate this radial part by geometrical arguments, whereas the algebraic approach makes it very simple.

Again, we note that the radial part of the Laplace operator has been calculated in more general situation by Harish-Chandra (see [W, Chapter 9]), so (2.2.3) can be deduced from his results.

CHAPTER III

JACOBI POLYNOMIALS

In this chapter we show that in some special cases the generalized characters for the Lie algebra $\mathfrak{sl}_n$ coincide with so-called Jack polynomials, which were studied by Heckman and Opdam (see [HO, O1, H1, H2]). These results are due to the author and Pavel Etingof (see [EK2, EK4]). In fact, Heckman and Opdam defined and studied analogues of these polynomials for arbitrary root systems; they called them Jacobi polynomials associated with a root system, the reason being that for $\mathfrak{g} = \mathfrak{sl}_2$ these polynomials coincide with so-called ultraspherical (Gegenbauer) polynomials, which are a special case of Jacobi polynomials. In this chapter we only consider classical case, i.e. representations, intertwiners and generalized characters for $\mathfrak{g}$ rather than for $U_q\mathfrak{g}$.

3.1 Jacobi polynomials and corresponding differential operators

In thie section we give definition and main properties of Jacobi polynomials associated with an arbitrary reduced irreducible root system R, following papers of Heckman and Opdam.

Recall (see Chapter 2) that $\mathbb{C}[P]$ denotes the group algebra of the weight lattice, and $\mathbb{C}[P](e^\alpha - 1)^{-1}$ is the ring obtained by adjoining to $\mathbb{C}[P]$ the expressions of the form $(e^\alpha - 1)^{-1}, \alpha \in R$. Note that the elements of $\mathbb{C}[P]$ may be considered as functions on $\mathfrak{h}$ by the rule: $e^\alpha(h) = e^{2\pi \mathrm{i}\langle \alpha, h\rangle}$. Under this identification elements of $\mathbb{C}[P](e^\alpha - 1)^{-1}$ become functions on the real torus $T = \mathfrak{h}_\mathbb{R}/Q^\vee, \mathfrak{h}_\mathbb{R} = \bigoplus \mathbb{R}\alpha_i^\vee$ with singularities on the hypersurfaces $e^\alpha(h) = 1, \alpha \in R$. However, we will use the formal language as far as possible. Abusing the language, we will call elements of $\mathbb{C}[P]$ polynomials, and elements of $\mathbb{C}[P]^W$ symmetric polynomials. Similarly, we will talk of divisibility of polynomials meaning divisibility in the ring $\mathbb{C}[P]$.

Recall also (see Section 2.2) that we denote by DO the ring of differential operators in $\mathfrak{h}$ with coefficients from $\mathbb{C}[P](e^\alpha - 1)^{-1}$; again, they can be treated formally as derivations of the ring $\mathbb{C}[P](e^\alpha - 1)^{-1}$. In particular, for every $\alpha \in \mathfrak{h}^*$ we defined ∂_α so that $\partial_\alpha e^\lambda = (\lambda, \alpha)e^\lambda$ and the Laplace operator $\Delta_\mathfrak{h}$ so that $\Delta_\mathfrak{h} e^\lambda = (\lambda, \lambda)e^\lambda$.

Through this whole chapter, we fix some positive integer k.

Consider the following differential operator

(3.1.1)
$$L = L_k = \Delta_{\mathfrak{h}} - k(k-1)\sum_{\alpha\in R^+}\frac{(\alpha,\alpha)}{(e^{\alpha/2}-e^{-\alpha/2})^2}.$$

This operator for the root system A_n was introduced by Sutherland ([Su]) and for an arbitrary root system by Olshanetsky and Perelomov ([OP]) as a Hamiltonian of an integrable quantum system. We will call L the Sutherland operator.

As before, let δ be the Weyl denominator defined by (1.1.6).

Define the following version of the Sutherland operator:

(3.1.2)
$$M_k = \delta^{-k}(L_k - k^2(\rho,\rho))\delta^k.$$

Proposition 3.1.1. ([HO])

(1)

(3.1.3)
$$M_k = \Delta_{\mathfrak{h}} - k\sum_{\alpha\in R^+}\frac{1+e^{\alpha}}{1-e^{\alpha}}\partial_\alpha = \Delta_{\mathfrak{h}} - 2k\sum_{\alpha\in R^+}\frac{1}{1-e^{\alpha}}\partial_\alpha + 2k\partial_\rho.$$

(2) *Both L_k, M_k commute with the action of the Weyl group.*

(3) *M_k preserves the algebra of symmetric polynomials $\mathbb{C}[P]^W \subset \mathbb{C}[P](e^\alpha-1)^{-1}$.*

Proof. We do not give the proof of (1) here, referring the reader to [HO]. Note that the proof involves some non-trivial statement about the root systems, which we will discuss later in the proof of the affine analogue of this statement (Theorem 7.1.2).

(2) is obvious, and (3) immediately follows from (1).

Recall (see 1.1.1) that we denoted by m_λ the basis of orbitsums in $\mathbb{C}[P]^W$: $m_\lambda = \sum\limits_{\mu\in W\lambda} e^\mu, \lambda \in P^+$.

Lemma 3.1.2.

(3.1.4)
$$M_k m_\lambda = (\lambda, \lambda + 2k\rho)m_\lambda + \sum_{\substack{\mu<\lambda\\ \mu\in P^+}} c_{\lambda\mu}m_\mu.$$

Proof. Explicit calculation.

Now we can consider the eigenfunction problem for M_k. Let us consider the action of M_k in the finite-dimensional space spanned by m_μ with $\mu \le \lambda$. Then the eigenvalue $(\lambda, \lambda + 2k\rho)$ has multiplicity one in this space due to the following trivial but very useful fact:

Lemma 3.1.3. *Let* $\lambda, \mu \in P^+$, $\mu < \lambda$. *Then* $(\mu+\rho, \mu+\rho) < (\lambda+\rho, \lambda+\rho)$.

Thus, we can give the following definition:

Definition 3.1.4. *Jacobi polynomials* $J_\lambda, \lambda \in P^+$ *are the elements of* $\mathbb{C}[P]^W$ *defined by the following conditions:*

(1) $J_\lambda = m_\lambda + \sum\limits_{\mu<\lambda} c_{\lambda\mu} m_\mu$.

(2) $M_k J_\lambda = (\lambda, \lambda + 2k\rho) J_\lambda$.

Due to Lemma 3.1.3, these properties determine J_λ uniquely. Note that this definition is valid for any complex k, and components of J_λ are rational functions of k.

Let us introduce an inner product in $\mathbb{C}[P]^W$. Let

$$\langle f, g\rangle_0 = \frac{1}{|W|}[f\bar{g}]_0, \tag{3.1.5}$$

where, as in Section 2.1, $[\ ]_0$ is the constant term of a polynomial, and the bar involution is defined by $\overline{e^\lambda} = e^{-\lambda}$. More generally, let

$$\langle f, g\rangle_k = \langle f\delta^k, g\delta^k\rangle_0. \tag{3.1.6}$$

Note that for $k = 1$ this definition coincides with previously given (2.1.3).

Lemma 3.1.5. M_k *is self-adjoint with respect to the inner product* $\langle\cdot,\cdot\rangle_k$.

Proof. This is equivalent to L_k being self-adjoint with respect to the inner product $\langle\cdot,\cdot\rangle_0$, which is obvious.

Corollary 3.1.6. $\langle J_\lambda, J_\mu\rangle_k = 0$ *if* $\lambda < \mu$.

In fact, one has a stronger result:

Theorem 3.1.7. ([O1]) $\langle J_\lambda, J_\mu\rangle_k = 0$ *if* $\lambda \neq \mu$.

We will prove this theorem for the root system A_n in the next section – see Theorem 3.2.4.

In fact, the operator M_k can be included in a large commutative family of differential operators. Define

$$\mathbb{D} = \{D \in DO | D \text{ is } W\text{-invariant}, [D, M_k] = 0\}. \tag{3.1.7}$$

Then we have the following theorem:

THEOREM 3.1.8.

(1) *Every operator $D \in \mathbb{D}$ preserves the space $\mathbb{C}[P]^W$, and J_λ is a common eigenbasis for the action of $\mathbb{D}$ in $\mathbb{C}[P]^W$: there exists a map $\gamma : \mathbb{D} \to (S\mathfrak{h})^W$ such that*

$$DJ_\lambda = \gamma(D)(\lambda + k\rho)J_\lambda. \tag{3.1.8}$$

(2) *γ is an isomorphism $\mathbb{D} \simeq (S\mathfrak{h})^W$.*

We do not give the proof of this theorem here, referring the reader to the above mentioned papers of Heckman and Opdam (in fact, for the root system A_n this theorem was known before). However, we note here that part (1) is relatively easy, and so is injectivity of γ; the difficult part is to prove surjectivity, or to construct γ^{-1}. Again, we will reprove it by representation-theoretic methods for the root system A_n in the next section.

3.2 Jack polynomials as generalized characters

In this section we show that one can get Jacobi polynomials for the root system A_{n-1}, in which case they are also known under the name "Jack polynomials", as generalized characters. This construction is due to the author and Pavel Etingof ([EK2, EK3]). **In this section, we only consider the case $\mathfrak{g} = \mathfrak{sl}_n$.**

Recall that we have fixed a positive integer k. Define the representation $U = U_{k-1}$ (which we will later use to define the generalized characters) as the irreducible representation of $\mathfrak{sl}_n$ with the highest weight $(k-1)n\omega_1$; it can be described as the symmetric power of the fundamental representation:

$$U = S^{(k-1)n}\mathbb{C}^n. \tag{3.2.1}$$

In other words, U can be identified with the space of homogeneous polynomials in $x_1, \dots, x_n$ of degree $(k-1)n$. The action of $\mathfrak{sl}_n$ is given by the following formulas:

$$\begin{gathered} e_i \mapsto x_i\partial_{i+1}, \quad f_i \mapsto x_{i+1}\partial_i \\ h_i \mapsto x_i\partial_i - x_{i+1}\partial_{i+1}, \end{gathered} \tag{3.2.2}$$

where $\partial_i = \frac{\partial}{\partial x_i}$.

It is very important for us that all weight subspaces of U are one-dimensional. In particular, the same is true for $U[0]$; we fix an element $u_0 = (x_1 \dots x_n)^{k-1} \in U[0]$, which allows us to identify

$$U[0] \simeq \mathbb{C} : \quad u_0 \mapsto 1. \tag{3.2.3}$$

Also, we will use the fact that

$$e_\alpha f_\alpha|_{U[0]} = k(k-1)\,\mathrm{Id}_{U[0]}\,.$$

Lemma 3.2.1. *Let* $\lambda \in P^+$. *A non-zero* $\mathfrak{sl}_n$*-homomorphism* $\Phi\colon L_\lambda \to L_\lambda \otimes U$ *exists iff* $\lambda - (k-1)\rho \in P^+$*; if it exists, it is unique up to a factor.*

Proof. Standard exercise.

For brevity, from now on we use the following notation:

$$\lambda^k = \lambda + (k-1)\rho. \tag{3.2.4}$$

For every $\lambda \in P^+$ fix an intertwiner

$$\Phi_\lambda : L_{\lambda^k} \to L_{\lambda^k} \otimes U \tag{3.2.5}$$

by the condition $\Phi_\lambda(v_{\lambda^k}) = v_{\lambda^k} \otimes u_0 +$ *lower order terms*, where v_{λ^k} is the highest weight vector in L_{λ^k}.

Define by φ_λ the corresponding generalized character:

$$\varphi_\lambda = \chi_{\Phi_\lambda} = \mathrm{Tr}_{L_{\lambda^k}}(\Phi_\lambda e^{2\pi i h}). \tag{3.2.6}$$

It takes values in the space $U[0]$; since this space is one-dimensional, we can identify it with $\mathbb{C}$ using (3.2.3) and consider φ_λ as scalar-valued function. Under this identification, φ_λ has the following form: $\varphi_\lambda = e^{\lambda+(k-1)\rho} +$ *lower order terms.*

Proposition 3.2.2.

$$\varphi_0 = \delta^{k-1}. \tag{3.2.7}$$

The proof of this proposition will be given later in a more general case (see Proposition 4.2.2).

Now we can prove the main theorem of this section:

Theorem 3.2.3. *Let* $\varphi_\lambda, \lambda \in P^+$ *be the generalized characters for* $\mathfrak{sl}_n$ *defined by formulas* (3.2.6), (3.2.7)*. Then* φ_λ *is divisible by* φ_0*, and the ratio* $\varphi_\lambda/\varphi_0$ *is the Jack polynomial* J_λ (*see Definition* 3.1.4).

Proof. Let us first prove that φ_λ is divisible by φ_0, and the ratio is a symmetric polynomial with highest term e^λ. Consider the tensor product $V = L_\lambda \otimes L_{(k-1)\rho}$. It decomposes as follows: $V = L_{\lambda+(k-1)\rho} + \sum_{\mu<\lambda} N_{\lambda\mu} L_{\mu+(k-1)\rho}$. Consider the intertwiner $\Phi = \mathrm{Id}_{L_\lambda} \otimes \Phi_0 : V \to V \otimes U$. On one hand, it follows from the definition that $\chi_\Phi = \mathrm{ch}\, L_\lambda \cdot \varphi_0$, where $\mathrm{ch}\, L_\lambda$ is the (usual) character of L_λ. On the other hand, the decomposition of V implies that $\chi_\Phi = \varphi_\lambda + \sum_{\mu<\lambda} a_{\lambda\mu}\varphi_\mu$, and thus $\varphi_\lambda/\varphi_0 = \mathrm{ch}\, L_\lambda + \sum_{\mu<\lambda} a_{\lambda\mu}\varphi_\mu/\varphi_0$. Since $\mathrm{ch}\, L_\lambda$ is a symmetric polynomial, it follows by induction in λ that $\varphi_\lambda/\varphi_0$ is also a symmetric polynomial.

Let us prove that $\varphi_\lambda/\varphi_0$ satisfies the equation $M_k(\varphi_\lambda/\varphi_0) = (\lambda, \lambda+2k\rho)(\varphi_\lambda/\varphi_0)$. Since $\varphi_0 = \delta^{k-1}$ (see Proposition 3.2.2), this equation is equivalent to

$$L_k \delta\varphi_\lambda = (\lambda + k\rho, \lambda + k\rho)\delta\varphi_\lambda. \tag{3.2.8}$$

On the other hand, recall that we have proved in Section 2.2 that the generalized characters are eigenfunctions of a certain differential operator D_C which was obtained from the Casimir element C. Comparing the expressions (2.2.3) for D_C and (3.1.1) for L_k, we see that

$$L_k = \delta \circ \pi_U(D_C) \circ \delta^{-1}, \tag{3.2.9}$$

since $e_\alpha f_\alpha u_0 = k(k-1)u_0$. Therefore, (3.2.8) immediately follows from the differential equation for generalized characters derived in Proposition 2.2.4. □

This immediately implies the orthogonality of Jack polynomials:

THEOREM 3.2.4. *If J_λ are Jack polynomials then*

$$\langle J_\lambda, J_\mu \rangle_k = 0 \quad \textit{if } \lambda \neq \mu. \tag{3.2.9}$$

PROOF. It follows immediately from Theorem 3.2.3 and the orthogonality theorem for generalized characters (Theorem 2.1.2).

Thus we have reproved by representation-theoretic arguments the orthogonality theorem 3.1.7 for Jacobi polynomials for the root system A_{n-1}. Note that the proof given by Heckman and Opdam uses transcendental (i.e., not algebraic) methods; the representation-theoretic approach outlined above gives probably the simplest proof of this fact. On the other hand, assuming (3.2.9) we could give an alternative proof of Theorem 3.2.3 – using the orthogonality theorem for generalized characters rather than the differential equation satisfied by them. We will use this approach in the next chapter, where we discuss q-analogue of Jacobi polynomials – Macdonald's polynomials.

Our construction also allows one to construct differential operators commuting with M_k. Namely, it follows from Theorem 2.2.3 that the operators of the form $D = \delta^{-(k-1)}\pi_U(D_c)\delta^{k-1}$, where $c \in \mathcal{Z}(U\mathfrak{g})$, commute with each other (and thus, with M_k, which can be obtained from the Casimir element) and are W-invariant. This means that the map

$$\mathcal{Z}(U\mathfrak{g}) \simeq (S\mathfrak{h})^W \to \mathbb{D} : c \mapsto \delta^{-(k-1)}\pi_U(D_c)\delta^{k-1}$$

is the inverse map to the map γ defined by (3.1.8). This proves (for the root system A_{n-1}) surjectivity of γ, i.e. the difficult part of Theorem 3.1.8 on the structure of $\mathbb{D}$.

CHAPTER IV

MACDONALD'S POLYNOMIALS AND GENERALIZED CHARACTERS

In this chapter we develop q-analogue of the constructions of Chapter 3. We define the q-analogue of Jacobi polynomials, called Macdonald polynomials, and show that in some special cases the generalized characters for the quantum group $U_q\mathfrak{sl}_n$ coincide with Macdonald's polynomials for the root system A_{n-1}. This construction is due to the author and Pavel Etingof [EK2].

In this chapter, the words "representation" etc. always stand for representation of the quantum group $U_q\mathfrak{g}$.

4.1 Definition of Macdonald's polynomials

In this section we give the definition of Macdonald's polynomials for reduced irreducible root system, following [M2].

We preserve the notations of Chapter 1. Consider the algebra of W-invariant polynomials $\mathbb{C}[P]^W$. The main goal of this section is to construct a basis in $\mathbb{C}_{q,t}[P]^W$, where $\mathbb{C}_{q,t} = \mathbb{C}(q,t)$ is the field of rational functions in two independent variables q,t. Unless otherwise stated, in this section "polynomial" will stand for an element of $\mathbb{C}_{q,t}[P]$, and divisibility will stand for divisibility in this ring; similarly, elements of $\mathbb{C}_{q,t}[P]$ will be called symmetric polynomials.

Recall (see (1.1.1)) that we denoted by m_λ the basis of orbitsums in $\mathbb{C}[P]^W$: $m_\lambda = \sum\limits_{\mu\in W\lambda} e^\mu, \lambda \in P^+$.

Theorem 4.1.1. (Macdonald)

There exists a unique family of polynomials $P_\lambda(q,t) \in \mathbb{C}_{q,t}[P]^W, \lambda \in P^+$ such that

(1) $P_\lambda = m_\lambda + \sum_{\mu<\lambda} c_{\lambda\mu} m_\mu$.

(2) *These polynomials are orthogonal with respect to the following inner product on $\mathbb{C}_{q,t}[P]$:*

$$\langle f, g\rangle_{q,t} = \frac{1}{|W|}[f\bar{g}\Delta_{q,t}]_0, \tag{4.1.1}$$

where, as before, the bar involution is defined by $\overline{e^{\lambda}} = e^{-\lambda}$, $[\ \ ]_0$ *is the constant term:* $[\sum c_\lambda e^\lambda]_0 = c_0$*, and*

$$\Delta_{q,t} = \prod_{\alpha\in R}\prod_{m=0}^{\infty} \frac{1-q^{2m}e^{\alpha}}{1-q^{2m}t^2e^{\alpha}}. \tag{4.1.2}$$

These polynomials are called Macdonald's polynomials (our notation slightly differs from that of Macdonald: what we denote by $P_\lambda(q,t)$ in the notations of [M2] would be $P_\lambda(q^2,t^2)$).

Remark. In fact, for non simply-laced systems there this theorem can be generalized, allowing different variables t_α for roots of different lengths; see [M2] for details.

It is often convenient to consider Macdonald's polynomials for $t = q^k, k \in \mathbb{Z}_+$ (see examples below). In this case, Macdonald's polynomials lie in $\mathbb{C}(q)[P]$; we will write $\langle\ ,\ \rangle_k$ instead of $\langle\ ,\ \rangle_{q,q^k}$, etc. (note that it agrees with Definition 3.1.6 for $q = 1$). However, most of the properties of Macdonald's polynomials obtained for $t = q^k$ can be generalized to the case when q, t are independent variables. Abusing the notations, we will sometimes say "$t = q^k$, k is an independent variable" instead of saying that q, t are independent variables.

Example.

1. For $k = 0$, we have $P_\lambda = m_\lambda$ independently of q; for $k = 1$, $P_\lambda = \operatorname{ch} L_\lambda$ – also independently of q.
2. In the limit $q, t \to 1$ so that $t = q^k$ Macdonald's polynomials tend to Jacobi polynomials introduced in Section 3.1, which follows from Theorem 3.1.7.

For the case $\mathfrak{g} = \mathfrak{sl}_n$ (that is, when R is the root system of type A_{n-1}), one can slightly modify the above definition and define Macdonald's polynomials for the $\mathfrak{gl}_n$ as a basis in $\mathbb{C}[x_1,\dots,x_n]^{S_n}$ labeled by the partitions λ. We will keep the same notation $P_\lambda(x;q,t)$ for these polynomials. In this form they were introduced in [M1].

For the root system A_{n-1} the polynomials $P_\lambda(x;q,t)$ can be defined in a different way; namely, they can be defined as the eigenfunctions of a certain family of commuting difference operators. Recall that we have identified $\mathbb{C}[P]^W$ for the root system A_{n-1} the space of with symmetric polynomials in $x_1^{\pm 1},\dots,x_n^{\pm 1}$ of degree zero (see end of Section 1.1). Let us define the following operators acting in $\mathbb{C}[P]^W$:

$$M_r = t^{r(r-n)} \sum_{\substack{I\subset\{1,\dots,n\}\\ |I|=r}} \Bigl(\prod_{\substack{i\in I\\ j\notin I}} \frac{t^2x_i - x_j}{x_i - x_j}\Bigr) T_{q^2,I}, \tag{4.1.3}$$

where $(T_{q^2,i}f)(x_1,\dots,x_n)=f(x_1,\dots,q^2x_i,\dots,x_n)$, $T_{q^2,I}=\prod_{i\in I}T_{q^2,i}$ and $r=1,\dots,n$. It is not too difficult (though it is not quite obvious) to show that these operators preserve the space of symmetric polynomials. Note that since the total degree is zero, $M_nf=f$ for every $f\in\mathbb{C}[P]^W$, and any difference operator acting in polynomials in $x_1,\dots,x_n$ is defined by its action in $\mathbb{C}[P]^W$ uniquely up to a multiple of M_n. We could consider the case of $\mathfrak{gl}_n$ rather then $\mathfrak{sl}_n$ thus replacing $\mathbb{C}[P]^W$ by $\mathbb{C}[x_1,\dots,x_n]^{S_n}$; then all M_i act non-trivially. Let us, however, stick to the $\mathfrak{sl}_n$ case.

Theorem 4.1.2. (Macdonald)

1. $[M_i,M_j]=0$.
2. *M_r is self-adjoint with respect to the inner product $\langle\cdot,\cdot\rangle_{q,t}$.*
3. *$M_rP_\lambda(x;q,t)=c^r_\lambda P_\lambda(x;q,t)$, and*

$$c^r_\lambda=\sum_{|I|=r}\prod_{i\in I}q^{2\lambda_i}t^{2\rho_i},$$

where $\rho_i=\frac{n+1-2i}{2}$ (see end of Section 1.1).

This characterization of Macdonald's polynomials is analogous to the definition of Jacobi polynomials as eigenfunctions of the commuting family $\mathbb{D}$ of differential operators given in Chapter 3. In fact, one can show that any differential operator $D\in\mathbb{D}$ can be obtained as a certain linear combinations of coefficients of expansion of Macdonald's difference operators M_i in powers of $(q-1)$. However, these expressions are rather messy; we will return to this relation later (see remark at the end of the next section).

We also note (though it is not relevant for our purposes) that the same holds for an arbitrary root system: Macdonald's polynomials can be defined as eigenfunctions of a certain family of commuting difference operators (see [C2]); unfortunately, for root systems other than A_n it is very difficult to write these difference operators explicitly.

4.2 Macdonald's polynomials of type A as generalized characters

Through this section, we assume $t=q^k, k\in\mathbb{N}$ and show how one gets Macdonald's polynomials $P_\lambda(x;q,q^k)$ for the root system A_{n-1} as generalized characters. The construction is quite parallel to that of Section 3.2. In this section we only consider $\mathfrak{g}=\mathfrak{sl}_n$; unless otherwise specified, the words "representation", "intertwiner", "generalized character" stand for representation of $U_q\mathfrak{sl}_n$, etc. Also, all the objects are defined over the field $\mathbb{C}_q=\mathbb{C}(q^{1/2n})$.

Let U be the finite-dimensional representation of $U_q\mathfrak{g}$ with the highest weight $(k-1)n\omega_1$; this is a q-analogue of the representation $U=S^{(k-1)n}\mathbb{C}^n$ considered in Section 3.2. As before, this representation can be realized explicitly in homogeneous polynomials of degree $(k-1)n$ of n variables $x_1,\dots,x_n$ with the action of $U_q\mathfrak{sl}_n$ given by

$$\begin{gathered}
h_i \mapsto x_i \frac{\partial}{\partial x_i} - x_{i+1} \frac{\partial}{\partial x_{i+1}}, \\
e_i \mapsto x_i D_{i+1}, \quad f_i \mapsto x_{i+1} D_i, \\
(D_i f)(x_1, \dots, x_n) = \frac{f(x_1, \dots, qx_i, \dots, x_n) - f(x_1, \dots, q^{-1}x_i, \dots, x_n)}{(q - q^{-1})x_i}.
\end{gathered} \tag{4.2.1}$$

Since the multiplicities in tensor products for $U_q\mathfrak{g}$ are the same as for $\mathfrak{g}$, we have the following proposition:

PROPOSITION 4.2.1. *Let* $\lambda \in P^+$. *A non-zero* $U_q\mathfrak{sl}_n$*-homomorphism* $\Phi\colon L_\lambda \to L_\lambda \otimes U$ *exists iff* $\lambda - (k-1)\rho \in P^+$*; if it exists, it is unique up to a factor.*

As in Chapter 3, we denote $\lambda^k = \lambda + (k-1)\rho$ and define the intertwiner Φ_λ and the corresponding generalized character φ_λ:

$$\begin{gathered}
\Phi_\lambda : L_{\lambda^k} \to L_{\lambda^k} \otimes U, \\
\varphi_\lambda = \chi_{\Phi_\lambda}.
\end{gathered} \tag{4.2.2}$$

As before, we consider φ_λ as a scalar-valued function, and choose the identification $U[0] \simeq \mathbb{C}$ so that $\varphi_\lambda = e^{\lambda+(k-1)\rho} +$ *lower order terms.*

PROPOSITION 4.2.2.

$$\varphi_0 = \prod_{i=1}^{k-1} \prod_{\alpha \in R^+} (e^{\alpha/2} - q^{2i} e^{-\alpha/2}). \tag{4.2.3}$$

PROOF. First, we prove the following statement:

LEMMA 1. φ_λ *is divisible by* $(1 - q^{2j} e^{-\alpha})$ *for any positive root* α *and* $1 \le j \le k-1$.

The proof is done in several steps. Let us introduce $F_i = f_i q^{-d_i h_i/2}$; then

$$\Delta(F_i) = F_i \otimes q^{-d_i h_i} + 1 \otimes F_i.$$

Let F be a (non-commutative) polynomial in $F_1, \dots, F_{n-1}$ of weight $-\alpha, \alpha \in Q^+$. Let $\varphi_\lambda^F = \operatorname{Tr}_{L_\lambda}(\Phi_\lambda F e^{2\pi i h})$. Also, let us fix a basis in U: $U[\alpha] = \mathbb{C} u_\alpha, \alpha \in Q$. Then

LEMMA 2. *There exists a polynomial* $P_F \in \mathbb{C}(q)[P]$ *such that*

$$\varphi_\lambda^F = \frac{P_F \varphi_\lambda}{\prod_{\substack{\beta \le \alpha \\ \beta \in Q^+}} (1 - q^{(\beta,\beta)} e^{-\beta})} u_{-\alpha}. \tag{4.2.4}$$

Proof is by induction in $\alpha \in Q^+$. For $\alpha = 0$ the statement is obvious. Now, let $\alpha = \sum m_i\alpha_i, \sum m_i = m$ and assume that the statement is proved for all $\alpha' < \alpha$. Take $F = F_{j_1} \dots F_{j_m}$. Then $\Delta F = \Delta(F_{j_1}) \dots \Delta(F_{j_m}) = \sum_i q^{\sigma_i} F(\gamma_i) \otimes \tilde{F}(\alpha - \gamma_i)q^{-h_{\gamma_i}} + F \otimes q^{-h_\alpha}$, where $\gamma_i \in Q^+, \gamma_i < \alpha, \sigma_i \in \mathbb{Z}$, and $F(\gamma)$ has weight $-\gamma$. Therefore, using the intertwining property of Φ_λ and the cyclic property of the trace, we get

$$\varphi_\lambda^F = \mathrm{Tr}(\Delta(F)\Phi_\lambda e^{2\pi \mathrm{i} h}) = q^{-h_\alpha}\,\mathrm{Tr}(F\Phi_\lambda e^{2\pi \mathrm{i} h}) + A = q^{(\alpha,\alpha)}e^{-\alpha}\varphi_\lambda^F + A,$$

where $A = \sum q^{\sigma_i}\tilde{F}(\alpha - \gamma_i)q^{-h_{\gamma_i}}\,\mathrm{Tr}(F(\gamma_i)\Phi_\lambda e^{2\pi \mathrm{i} h})$. Thus,

$$\varphi_\lambda^F = \frac{1}{1 - q^{(\alpha,\alpha)}e^{-\alpha}}A.$$

On the other hand, it follows from the induction assumption that A is an expression of the form (4.2.4) containing only the factors $1 - q^{(\beta,\beta)}e^{-\beta}$ with $\beta < \alpha$ in the denominator. This completes the proof of Lemma 2.

LEMMA 3. *Let $\alpha \in R^+$, and let F_α be a (non-commutative) polynomial in F_i which in the limit $q = 1$ becomes a root element of $\mathfrak{sl}_n$. Then $P_{F_\alpha^{k-1}}$ is a non-zero polynomial relatively prime to $\prod_{j=1}^{k-1}(1 - q^{2j}e^{-\alpha})$.*

It suffices to prove this lemma for $q = 1$. But for $q = 1$, $\Delta(F_\alpha) = F_\alpha \otimes 1 + 1 \otimes F_\alpha$, and therefore

$$\begin{aligned}\varphi_\lambda^{F_\alpha^{k-1}} &= \mathrm{Tr}(\Phi_\lambda F_\alpha^{k-1}e^{2\pi \mathrm{i} h}) = F_\alpha\,\mathrm{Tr}(\Phi_\lambda F_\alpha^{k-2}e^{2\pi \mathrm{i} h}) + \mathrm{Tr}(\Phi F_\alpha^{k-2}e^{2\pi \mathrm{i} h}F_\alpha) \\ &= F_\alpha\varphi_\lambda^{F_\alpha^{k-2}} + e^{-\alpha}\varphi_\lambda^{F_\alpha^{k-1}},\end{aligned}$$

so

$$\varphi_\lambda^{F_\alpha^{k-1}} = (1 - e^{-\alpha})^{-1}F_\alpha\varphi_\lambda^{F_\alpha^{k-2}} = \dots = (1 - e^{-\alpha})^{1-k}\varphi_\lambda F_\alpha^{k-1}u_0.$$

Since $F_\alpha^{k-1}u_0 = c_\alpha u_{(1-k)\alpha}$ for some $c_\alpha \neq 0$, we see that

$$P_{F_\alpha^{k-1}} = c_\alpha \frac{\prod_{\beta \le (k-1)\alpha}(1 - e^{-\beta})}{(1 - e^{-\alpha})^{k-1}} = c_\alpha \prod_{\substack{\beta < (k-1)\alpha \\ \beta \neq s\alpha}}(1 - e^{-\beta})\prod_{s=1}^{k-1}(1 + e^{-\alpha} + \dots + e^{(1-s)\alpha}).$$

One can easily see that this polynomial is relatively prime to $1 - e^{-\alpha}$. Thus, we have proved Lemma 3.

Now, let us return to the proof of Lemma 1. Let us write

$$\mathrm{Tr}(\Phi_\lambda F_\alpha^{k-1} e^{2\pi i h}) = \frac{P_{F_\alpha^{k-1}} \varphi_\lambda}{\prod\limits_{j=1}^{k-1} (1 - q^{2j} e^{-\alpha})}.$$

Since the left-hand side is a non-zero polynomial, and $P_{F_\alpha^{k-1}}$ is relatively prime to $\prod\limits_{j=1}^{k-1} (1 - q^{2j} e^{-\alpha})$, we see that φ_λ must be divisible by $\prod\limits_{j=1}^{k-1} (1 - q^{2j} e^{-\alpha})$. So, Lemma 1 is proved.

Now it is easy to prove Proposition 4.2.2: Lemma 1 implies that we have the following identity: $\varphi_0 = f \prod\limits_{j=1}^{k-1} \prod\limits_{\alpha \in R^+} (1 - q^{2j} e^{-\alpha})$ for some polynomial f; comparing the highest and the lowest terms on both sides we see that $f = e^{(k-1)\rho}$. This completes the proof of Proposition 4.2.2. □

Now we can prove the main theorem of this section:

THEOREM 4.2.3. *If $\lambda \in P^+$ and $\varphi_\lambda, \varphi_0$ are the generalized characters for $U_q\mathfrak{sl}_n$ defined by* (4.2.2), (4.2.3) *then φ_λ is divisible by φ_0, and the ratio $\varphi_\lambda / \varphi_0$ is the Macdonald's polynomial $P_\lambda(q, q^k)$ for the root system A_{n-1}.*

PROOF. First, the same arguments as in the proof of Theorem 3.2.3 – no changes are needed – show that $\varphi_\lambda / \varphi_0$ is a symmetric polynomial with highest term e^λ. Therefore, to prove the theorem it suffices to prove that these ratios are orthogonal with respect to the inner product $\langle \cdot, \cdot \rangle_k$. This immediately follows from the orthogonality theorem for generalized characters and formula for φ_0 (Proposition 4.2.2). Indeed, we know from the orthogonality theorem that $[\varphi_\lambda \bar{\varphi}_\mu \Delta]_0 = 0$ if $\lambda \neq \mu$. Therefore, $[(\varphi_\lambda/\varphi_0)\overline{(\varphi_\mu/\varphi_0)} \varphi_0 \bar{\varphi}_0 \Delta]_0 = 0$. Due to Proposition 4.2.2, $\varphi_0 \bar{\varphi}_0 \Delta = \prod\limits_{\alpha \in R} \prod\limits_{i=0}^{k-1} (1 - q^{2i} e^\alpha) = \Delta_{q,t}$, which proves the orthogonality of $\{\varphi_\lambda / \varphi_0\}$ with respect to the inner product $\langle \cdot, \cdot \rangle_k$. □

REMARK 4.2.4. Note that the proof only uses the following properties of U:

(1) All weight subspaces are one dimensional.
(2) $e_\alpha^k u_0 = 0, e_\alpha^{k-1} u_0 \neq 0$ for every $\alpha \in R^+$.

(we also used Proposition 4.2.1, which ensures existence of intertwiners; however, it can be deduced from properties (1), (2) above.)

Thus, the same theorem must be true if we replace U by any other representation satisfying these properties. However, one can check that for $U_q\mathfrak{sl}_n$ we have only two possibilities: $U = L_{(k-1)n\omega_1}$ (which we used) or $U = L_{(k-1)n\omega_{n-1}}$, which can be obtained from the previous one by an outer automorphism of $\mathfrak{g}$, i.e. by flip of the Dynkin diagram. For other Lie algebras, such representations do not exist at all (except some small number of exceptional cases, where such representations exist only for finite number of values of k), which explains why this theory can not be generalized to arbitrary root systems in a trivial way.

Remark. This theorem can be generalized for the case of generic k, i.e., the case where q, t are independent variables; see details in [EK2].

4.3 The center of $U_q\mathfrak{sl}_n$ and Macdonald's operators

In this section we show how one can get Macdonald's operators M_r introduced in Section 4.1 from the quantum group $U_q\mathfrak{sl}_n$. This construction is parallel to the one for $q = 1$ (see Section 3.2). As before, in this section we only consider $\mathfrak{g} = \mathfrak{sl}_n$ and $t = q^k, k \in \mathbb{N}$.

Recall (see Section 2.3) that we have denoted by DO_q the ring of difference operators, i.e. operators of the form $\sum\limits_{\alpha \in \frac{1}{2}P^\vee} a_\alpha T_\alpha$, where $T_\alpha e^\lambda = q^{\langle \alpha, \lambda \rangle} e^\lambda$, and $a_\alpha \in \mathbb{C}_q[P](q^m e^\mu - 1)^{-1}$. We have also constructed for every element c of the center of $U_q\mathfrak{sl}_n$ a difference operator $D_c \in DO_q$ such that for every intertwiner Φ we have $\operatorname{Tr}(\Phi c e^{2\pi i h}) = D_c \operatorname{Tr}(\Phi e^{2\pi i h})$. These operators commute, and generalized characters are their common eigenfunctions.

Let us show that applying this construction to the above case (i.e., $\mathfrak{g} = \mathfrak{sl}_n$ and U is chosen as in the beginning of Section 4.2) we can get Macdonald's difference operators M_r defined by (4.1.3).

Theorem 4.3.1. *Define the central elements* $c_r \in \mathcal{Z}(U_q\mathfrak{sl}_n), r = 1 \dots, n-1$ *by*

$$c_r = c_{\Lambda_q^{n-r}}, \tag{4.3.1}$$

(*cf.* *Theorem* 1.2.1), *where* Λ_q^i *is the q-deformation of the representation of* $\mathfrak{sl}_n$ *in the i-th exterior power* $\Lambda^i \mathbb{C}^n$ *of the fundamental representation, and let* $D_{c_r} \in DO_q$ *be the corresponding difference operator* (*see Theorem* 2.3.1). *Then*

$$M_r = \varphi_0^{-1} \circ D_{c_r} \circ \varphi_0, \tag{4.3.2}$$

where M_r *is Macdonald's operator introduced in* (4.1.3), *and* φ_0 *is the operator of multiplication by the function* φ_0 *defined by* (4.2.3).

Proof. Let us first prove that

$$\varphi_0^{-1} D_{c_r}(\varphi_0 P_\lambda) = \sum_I q^{2\sum_{i \in I}(\lambda + k\rho)_i} P_\lambda, \tag{4.3.3}$$

where the sum is taken over all subsets $I \subset \{1, \dots, n\}$ of cardinality r. Indeed, we know from Theorem 2.3.4 that

$$D_r \varphi_\lambda = \operatorname{ch} \Lambda^{n-r}(q^{-2(\lambda + k\rho)}) \varphi_\lambda.$$

Since $\operatorname{ch} \Lambda^{n-r} = e^{\frac{n-r}{r}(1,\dots,1)} \sum e^\mu$, where the sum is taken over all $\mu = (\mu_1, \dots, \mu_n)$ such that $\mu_i = 0$ or 1, $\sum \mu_i = n - r$, we get

$$\operatorname{ch}\Lambda^{n-r}(q^{-2(\lambda+k\rho)}) = \sum_I q^{2\sum_{i\in I}(\lambda+k\rho)_i}.$$

Since $P_\lambda = \varphi_\lambda/\varphi_0$, we get (4.3.3).

Comparing (4.3.3) with the formula (4.1.4) for eigenvalues of Macdonald's operators, we see that M_r and $\varphi_0^{-1}\circ D_{c_r}\circ\varphi_0$ coincide on Macdonald's polynomials, and thus, on all symmetric polynomials. Repeating the uniqueness arguments outlined in the proof of Lemma 2.3.2, we see that it is only possible if they are equal. □

Remarks.

(1) Recently a straightforward proof of Theorem 4.3.1 was found by Mimachi ([Mi]).

(2) The central elements c_r are closely related to those constructed in [FRT]. Essentially, the central elements constructed in [FRT] are traces of the powers of L-matrix, whereas c_r are coefficients of the characteristic polynomial of L.

This theorem allows one to see the relation between the differential operators from $\mathbb{D}$ (see (3.1.7)) and Macdonald's difference operators. This relation is nothing but the relation between the center of $U_q\mathfrak{sl}_n$ (which, as we have seen, is spanned by the elements $c_V = (1\otimes \operatorname{Tr}_V)(\mathcal{R}^{21}\mathcal{R}(1\otimes q^{-2\rho}))$) and the center of $U\mathfrak{sl}_n$, which does not have a nice explicit description, but can be described by means of Harish-Chandra isomorphism $\mathcal{Z}(U\mathfrak{g})\simeq(S\mathfrak{h})^W$.

CHAPTER V

INNER PRODUCT AND SYMMETRY IDENTITIES FOR MACDONALD'S POLYNOMIALS

In this chapter we use the technique of generalized characters developed in the previous sections to prove so-called inner product (norm) and symmetry identities for Macdonald's polynomials of type A_n. These results are due to the author and Pavel Etingof ([EK5]).

Inner product identities express the norm $\langle P_\lambda, P_\lambda\rangle$ as a certain product over the positive roots. They have been conjectured by Macdonald for arbitrary root systems. He also gave a proof for A_n (unpublished); the proof for arbitrary root systems was given in a recent paper of Cherednik [C2].

Symmetry identity relates the values of $P_\lambda(q^{2(\mu+k\rho)})$ and $P_\mu(q^{2(\lambda+k\rho)})$. For A_n it was first proved by Koornwinder (unpublished), so the first published proof is in [EK5]. Again, recently Cherednik proved this identity for arbitrary root systems ([C3]).

In this chapter we only consider root system of type A_{n-1}, i.e. $\mathfrak{g} = \mathfrak{sl}_n$. Our proofs use the realization of Macdonald's polynomials as generalized characters for $U_q\mathfrak{sl}_n$ (see Chapter 4) and the technique of representing identities in the category of representations of a quantum group by ribbon graphs, developed by Reshetikhin and Turaev. We refer the reader to their papers [RT1, RT2] or to recent books of Turaev [T] and Kassel [Kas] for description of this technique; a very brief introduction can be found in the Appendix to [EK5].

5.1 Inner product identities

Let us fix a positive integer k. Recall (see Chapter 4) that we have defined Macdonald's polynomials $P_\lambda \in \mathbb{C}_q[P], \lambda \in P^+$ and an inner product $\langle\, , \rangle_k$ in $\mathbb{C}_q[P]$ such that $\langle P_\lambda, P_\mu\rangle_k = 0$ if $\lambda \neq \mu$. The goal of this section is to calculate $\langle P_\lambda, P_\lambda\rangle_k$; our proof is based on Theorem 4.2.3, which shows that P_λ can be expressed in terms of generalized characters for $U_q\mathfrak{sl}_n$.

Recall the notations $U = U_{k-1} = S^{(k-1)n}\mathbb{C}^n, u_0 = u_0^{k-1} \in U[0], \lambda^k = \lambda + (k-1)\rho, \Phi_\lambda : L_{\lambda^k} \to L_{\lambda^k} \otimes U, \varphi_\lambda = \chi_{\Phi_\lambda} = e^{\lambda^k} \otimes u_0 +$ *lower order terms* introduced in Chapter 4; also, recall the Chevalley involution ω and Shapovalov form $(\, ,)_V$:

$V \otimes V^\omega \to \mathbb{C}$, discussed in Section 1.2. We assume that Shapovalov form in U is normalized so that $(u_0, u_0)_U = 1$.

Lemma 5.1.1. *The inner product $\langle P_\lambda, P_\lambda \rangle_k$ can be calculated from*

$$\Psi \mathbf{1}_\lambda = \langle P_\lambda, P_\lambda \rangle_k \mathbf{1}_\lambda, \tag{5.1.1}$$

where $\mathbf{1}_\lambda$ is an invariant vector in $L_{\lambda^k} \otimes L^\omega_{\lambda^k}$ and $\Psi : L_{\lambda^k} \otimes L^\omega_{\lambda^k} \to L_{\lambda^k} \otimes L^\omega_{\lambda^k}$ is the following operator:

$$L_{\lambda^k} \otimes L^\omega_{\lambda^k} \xrightarrow{\Phi_\lambda \otimes \Phi^\omega_\lambda} L_{\lambda^k} \otimes U \otimes U^\omega \otimes L^\omega_{\lambda^k} \xrightarrow{\mathrm{Id} \otimes (\cdot,\cdot)_U \otimes \mathrm{Id}} L_{\lambda^k} \otimes L^\omega_{\lambda^k}. \tag{5.1.2}$$

Proof. It follows from Theorem 4.2.3 that $\langle P_\lambda, P_\lambda \rangle_k = \langle \varphi_\lambda, \varphi_\lambda \rangle_1$, where the inner product on generalized characters is introduced in Theorem 2.1.2. Now we can repeat the same arguments we used in the proof of Theorem 2.1.2. □

It will be convenient to rewrite this in a slightly different way as follows:

Theorem 5.1.2. *In the notations of previous Lemma, we have:*

$$\langle P_\lambda, P_\lambda \rangle_k = (\langle v^*_{\lambda^k}, \Phi_\lambda \Phi^\circ_\lambda v_{\lambda^k} \rangle)_U, \tag{5.2.3}$$

where the intertwiner $\Phi^\circ_\lambda : L_{\lambda^k} \to L_{\lambda^k} \otimes U^\omega$ is defined by the condition $\Phi^\circ_\lambda(v_{\lambda^k}) = v_{\lambda^k} \otimes (u_0)^\omega +$ lower order terms.

Proof.

The proof is obvious if we use the technique of ribbon graphs. Namely, it follows from Lemma 5.1.1 that the inner product $\langle P_\lambda, P_\lambda \rangle_k = A_\lambda$ can be defined from the following identity of ribbon graphs:



where $\dim_q L = \mathrm{Tr}_L(q^{-2\rho})$, $\phi : L^*_{\lambda^k} \to L_{\lambda^{k*}}$, $\psi : L_{n(k-1)\omega_1^*} \to L^*_{n(k-1)\omega_1} = U^*_{k-1}$ are isomorphisms and ψ is chosen so that $\langle u_0, \psi(u_0^{\omega})\rangle = 1$. It is easy to check that



and thus,



so



□

Theorem 5.1.3. *In the notations of the previous theorem, we have:*

$$(\Phi_\lambda \Phi^\circ_\lambda)_U = \prod_{\alpha\in R^+} \prod_{i=1}^{k-1} \frac{1-q^{2(\alpha,\lambda+k\rho)+2i}}{1-q^{2(\alpha,\lambda+k\rho)-2i}} \operatorname{Id}_{L_{\lambda^k}} . \tag{5.1.4}$$

This theorem will be proved in the next section, using detailed analysis of the poles of the intertwining operators.

This theorem, along with Theorem 5.1.2 immediately gives the main result of this section:

Theorem 5.1.4. (Macdonald)

$$\langle P_\lambda, P_\lambda\rangle_k = \prod_{\alpha\in R^+} \prod_{i=1}^{k-1} \frac{1-q^{2(\alpha,\lambda+k\rho)+2i}}{1-q^{2(\alpha,\lambda+k\rho)-2i}}. \tag{5.1.5}$$

This is precisely the Macdonald's inner product identity for the root system A_{n-1}.

Remark. As before, these identities can be easily generalized to the case of arbitrary k – see [EK5].

5.2 Algebra of intertwiners.

This section is devoted to the proof of Theorem 5.1.3, which expresses product of two intertwining operators in terms of a single intertwiner. The technique we use here is considering the intertwining operators for arbitrary value of λ (which, of course, requires use of Verma modules) and study their behavior (in particular, poles) as functions of λ, which can be done using the Shapovalov determinant formula.

Let us start with more general situation. Consider intertwiners of the form:

$$\Phi^{\mu}_{\lambda} : M_{\lambda} \to M_{\lambda} \otimes L_{\mu}, \tag{5.2.1}$$

where M_λ is Verma module over $U_q\mathfrak{g}$, L_μ is the finite-dimensional irreducible module; we assume that $\mu \in P^+ \cap Q$, so $L_\mu[0] \neq 0$. Let $u \in L_\mu[0]$. We consider all the modules over the field of rational functions $\mathbb{C}_q = \mathbb{C}(q^{1/2N})$, where $N = |P/Q|$; if λ is not an integral weight then we also have to add $q^{\langle \lambda, \alpha_i^\vee \rangle/2N}$ to this field.

It is known that if M_λ is irreducible then there exists a unique intertwiner of the form (5.2.1) such that $\Phi^{\mu}_{\lambda}(v_\lambda) = v_\lambda \otimes u +$ *lower order terms.* We will denote this intertwiner by $\Phi^{\mu,u}_{\lambda}$. The same is true if we consider the weight λ as indeterminate, i.e. if we consider $t_i = q^{\langle \lambda, \alpha_i^\vee \rangle/2N}$ as algebraically independent variables over $\mathbb{C}_q$.

Let us identify M_λ with U^- in a standard way. Then we can say that we have a family of actions of $U_q\mathfrak{g}$ in the same space $M \simeq U^-$, and thus we have a family of intertwiners $\Phi^{\mu,u}_{\lambda} : M \to M \otimes L_\mu$, defined for generic values of λ. This identification also gives rise to Shapovalov form in U^-. We fix a homogeneous basis g_k in U^- and denote the matrix elements of Shapovalov form in this basis by F_{kl}: $F_{kl} = (g_k v_\lambda, g_l v_\lambda)_{M_\lambda} = (v_\lambda, (\omega S(g_k)) g_l v_\lambda)_{M_\lambda}$.

For $\lambda \in \mathfrak{h}^*$, let us call a trigonometric rational function of λ a rational function in $q^{1/2N}, q^{\lambda/2N}$ (that is, in $q^{1/2N}$ and $t_i = q^{\langle \alpha_i^\vee, \lambda \rangle/2N}, i = 1, \dots, n$) and call a trigonometric polynomial in λ a polynomial in $q^{\pm\lambda/2N}$ with coefficients from $\mathbb{C}_q$. Note that the ring of trigonometric polynomials is a unique factorization ring, and invertible elements in this ring are of the form $c(q)q^{\langle \lambda, \alpha \rangle}$, $\alpha \in \frac{1}{2N}Q^\vee$.

Lemma 5.2.1. *For fixed $\mu \in P^+, u \in L_\mu[0], u \neq 0$ we have*

1. *Let $\lambda \in \mathfrak{h}^*$ be such that M_λ is irreducible. Then there exists a unique intertwining operator $\Phi^{\mu,u}_{\lambda} : M_\lambda \to M_\lambda \otimes L_\mu$ such that $\Phi^{\mu,u}_{\lambda}(v_\lambda) = v_\lambda \otimes u +$ lower order terms. Its matrix elements are trigonometric rational functions of λ. Moreover, we have the following formula for Φv_λ:*

$$\Phi^{\mu,u}_{\lambda}(v_\lambda) = \sum_{k,l} (F^{-1})_{kl} g_k v_\lambda \otimes q^{\lambda+2\rho} \omega(g_l) u, \tag{5.2.2}$$

where g_k is a homogeneous basis in U^-, F^{-1} is the inverse matrix to the Shapovalov form in M_λ, and as in Section 1.2, $q^\lambda|_{V[\nu]} = q^{(\lambda,\nu)'} \mathrm{Id}_{V[\nu]}$.

2. *Define the operator $\tilde{\Phi}^{\mu,u}_{\lambda}$ by*

$$\tilde{\Phi}^{\mu,u}_{\lambda} = d_\mu(\lambda) \Phi^{\mu,u}_{\lambda}, \tag{5.2.3}$$

where

$$d_\mu(\lambda) = \prod_{\alpha\in R^+}\prod_{i=1}^{n_\mu^\alpha}\left(1 - q^{(2(\alpha,\lambda+\rho)' - i(\alpha,\alpha)')}\right), \tag{5.2.4}$$
$$n_\mu^\alpha = \max\{i \in \mathbb{Z}_+ | L_\mu[i\alpha] \neq 0\}.$$

Then matrix elements of $\tilde\Phi_\lambda^{\mu,u}$ *are trigonometric polynomials, i.e. have no poles; thus,* $\tilde\Phi_\lambda^{\mu,u}$ *is well defined for all* λ.

(3) *Consider the special case* $\mathfrak{g} = \mathfrak{sl}_n, \mu = kn\omega_1$. *Recall that in this case* $L_\mu[0]$ *is one-dimensional. Let* u *be a non-zero vector in* $L_\mu[0]$. *Then* $d_\mu(\lambda)$ *is the least common denominator of matrix elements of* $\Phi_\lambda^{\mu,u}$*; in other words, in this case matrix elements of* $\tilde\Phi_\lambda^{\mu,u}$ *do not have non-trivial common divisors.*

Proof. The proof is essentially the same as in the classical case, which is given in [ES]; however, we repeat it here marking necessary changes. To prove (1), it suffices to check that the vector in the right-hand side is the unique highest-weight vector of weight λ in $M_\lambda \otimes L_\mu$ of the form $v = v_\lambda \otimes u + \dots$. Suppose $v \in M_\lambda \otimes L_\mu$ is of the form above. Define $E_i = e_i q^{d_i h_i/2}$; then v is highest-weight iff $\Delta E_i v = 0$. On the other hand, explicit calculation shows that $\Delta E_i = E_i \otimes 1 - q^{2d_i}(1 \otimes SE_i) q^{d_i h_i} \otimes q^{d_i h_i}$. Thus,

$$\Delta E_i v = (E_i \otimes 1 - q^{d_i(2+\langle\lambda,\alpha_i^\vee\rangle)} 1 \otimes SE_i)v = (E_i \otimes 1 - q^{(\alpha_i,\lambda+2\rho)'} 1 \otimes SE_i)v.$$

For $x \in M_\lambda, w \in M_\lambda \otimes L_\mu$ define (x, w) by $(x, w_1 \otimes w_2) = (x, w_1)_{M_\lambda} w_2 \in L_\mu$. If M_λ is irreducible, Shapovalov form in M_λ is non-degenerate, and therefore, $w = 0 \iff (x, w) = 0$ for all $x \in M_\lambda$. Therefore, we can rewrite the condition that v is a highest-weight vector as follows:

$$\begin{aligned}&\Delta E_i v = 0 \iff (x, \Delta E_i v) = 0 \quad \text{for all } x \in M_\lambda \iff \\ &(x, (E_i \otimes 1 - q^{(\alpha_i,\lambda+2\rho)'} 1 \otimes SE_i)v) = 0 \iff \\ &(\omega S(E_i)x, v) = q^{(\alpha_i,\lambda+2\rho)'} SE_i(x, v).\end{aligned}$$

It is easy to see that the last condition is equivalent to the following: for any homogeneous $F \in U^-$, we have

$$(Fv_\lambda, v) = q^{-(\mathrm{wt}\ F,\lambda+2\rho)'}\omega(F)(v_\lambda, v) = q^{-(\mathrm{wt}\ F,\lambda+2\rho)'}\omega(F)u.$$

This proves that the highest-weight vector of the desired form exists and is unique. It is easy to check that the vector given by (5.2.2) satisfies the condition above.

To prove (2), note that it follows from (5.2.2) that matrix coefficients of Φ may have poles only at the points where the determinant of Shapovalov form vanishes.

The formula for the determinant of the Shapovalov form in the quantum case can be found in [CK], and the factors occuring there are precisely the factors in formula (5.2.4) (up to invertible factors). One can check in the same way as it is done in [ES] for $q = 1$ – that is, by comparing the order of pole of the matrix of Shapovalov form and its minors – that in fact all the poles of F^{-1} are simple.

The restriction on i in (5.2.4) appears because the coefficients $(F^{-1})_{kl}$ which may have poles of the form (5.2.4) with $i > n_\mu^\alpha$ appear with zero coefficient.

To prove (3), it suffices to check this statement for $q = 1$, which is also done in the paper [ES] (note that it is quite non-trivial!). □

REMARK. It is seen from this proof that $\tilde{\Phi}_\lambda^{\mu,u}$ is actually a trigonometric polynomial in λ with operator coefficients, i.e. the degrees of its matrix coefficients, as trigonometric polynomials, are uniformly bounded (under a suitable definition of degree).

We will discuss what happens to these intertwiners when $\lambda \in P^+$ and how they are related with intertwiners $L_\lambda \to L_\lambda \otimes L_\mu$ later.

We will also need one more technical lemma.

LEMMA 5.2.2. *Let us write* $\tilde{\Phi}_\lambda^{\mu,u} = d_\mu(\lambda)\Phi_\lambda^{\mu,u}$ (*see* (5.2.4)) *in the following form:*

$$\tilde{\Phi}_\lambda^{\mu,u} v_\lambda = d_\mu(\lambda) v_\lambda \otimes u + \ldots + a(\lambda) v_\lambda \otimes u_\mu,$$

where u_μ *is the highest-weight vector in* L_μ, *and* $a(\lambda) \in \mathbb{C}_q[q^{\pm\lambda/2N}] \otimes U^-[-\mu]$ *is a trigonometric polynomial of* λ *with values in the universal enveloping algebra. Then the greatest common divisor of the components of* $a(\lambda)$ *is 1.*

PROOF. It is easy to see, using the irreducibility of L_μ, that if $a(\lambda) = 0$ then $\tilde{\Phi}_\lambda^{\mu,u} = 0$. On the other hand, we have shown before that the coefficients of $\tilde{\Phi}_\lambda^{\mu,u}$ have no nontrivial common divisors, and thus $\tilde{\Phi}_\lambda^{\mu,u}$ could only vanish on a subvariety of codimension more than one. Thus, the same must be true for $a(\lambda)$.

Now we want to define a structure of algebra on these intertwiners. Let $\Phi_1 : M_\lambda \to M_\lambda \otimes L_{\mu_1}, \Phi_2 : M_\lambda \to M_\lambda \otimes L_{\mu_2}$ be non-zero intertwiners. Let us define their product $\Phi_1 * \Phi_2 : M_\lambda \to M_\lambda \otimes L_{\mu_1+\mu_2}$ as the composition

$$M_\lambda \xrightarrow{\Phi_2} M_\lambda \otimes L_{\mu_2} \xrightarrow{\Phi_1 \otimes 1} M_\lambda \otimes L_{\mu_1} \otimes L_{\mu_2} \xrightarrow{1 \otimes \pi} M_\lambda \otimes L_{\mu_1+\mu_2}, \tag{5.2.5}$$

where π is a fixed projection $\pi : L_{\mu_1} \otimes L_{\mu_2} \to L_{\mu_1+\mu_2}$.

Now, let us apply these notions to the situation considered in the previous sections in connection with Macdonald's theory. That is, assume that $\mathfrak{g} = \mathfrak{sl}_n$, $\mu = kn\omega_1$ for some $k \in \mathbb{Z}_+$, so that $L_\mu = U_k = q-$ analogue of $S^{kn}\mathbb{C}^n$. As before, we identify U_k with the space of homogeneous polynomials of degree kn in variables $x_1, \ldots, x_n$ (see (4.2.1)), and let $u_0 = (x_1 \ldots x_n)^k \in L_\mu[0]$; then $L_\mu[0] = \mathbb{C}_q u_0^k$. For brevity, we will write U_k for $L_{kn\omega_1}$, Φ_λ^k for $\Phi_\lambda^{\mu=kn\omega_1, u_0^k}$, etc. Let us fix the projection $\pi : U_k \otimes U_l \to U_{k+l}$ by $\pi(u_0^k \otimes u_0^l) = u_0^{k+l}$. In this case, $n_\mu^\alpha = k$ for all $\alpha \in R^+$ and

$$d_k(\lambda) = \prod_{\alpha\in R^+}\prod_{i=1}^{k}(1-q^{2(\alpha,\lambda+\rho)-2i}). \tag{5.2.6}$$

Here comes the main result of this section:

THEOREM 5.2.3.

$$\tilde{\Phi}^k_\lambda * \tilde{\Phi}^l_\lambda = \tilde{\Phi}^{k+l}_\lambda. \tag{5.2.7}$$

PROOF. Let us denote the left-hand side of (5.2.7) by Ψ. Then Ψ is an intertwiner $M_\lambda \to M_\lambda \otimes U_{k+l}$, whose matrix coefficients are trigonometric polynomials in λ. In particular, we can write $\Psi(v_\lambda) = f(\lambda)v_\lambda \otimes u_0^{k+l} + \text{l.o.t.}$ On the other hand $\tilde{\Phi}^{k+l}_\lambda(v_\lambda) = d_{k+l}(\lambda)v_\lambda \otimes u_0^{k+l} + \text{l.o.t.}$ Since the intertwining operator is unique for generic λ, this implies $\Psi(\lambda) = \frac{f(\lambda)}{d_{k+l}(\lambda)}\tilde{\Phi}^{k+l}_\lambda$. Since the greatest common divisor of the matrix elements of $\tilde{\Phi}^{k+l}$ is 1, this implies that $f(\lambda)$ is divisible by $d_{k+l}(\lambda)$.

Let us now consider the lowest term of Ψ. If we write the lowest term of $\tilde{\Phi}^k_\lambda$ as $a_k(\lambda)v_\lambda \otimes u_k$ (cf. Lemma 5.2.2) and lowest term of Φ^l_λ as $a_l(\lambda)v_\lambda \otimes u_l$ then the lowest term of Ψ will be $a_l(\lambda)a_k(\lambda)v_\lambda \otimes u_{k+l}$ (up to some power of q). Since we know that components of a_k have no common divisors, and the same is true for a_l, it follows that the greatest common divisor of components of $a_k(\lambda)a_l(\lambda)$ is 1. Indeed, suppose that $p(\lambda)$ is a common divisor of components of $a_k(\lambda)a_l(\lambda)$. Passing if necessary to a certain algebraic extension of $\mathbb{C}_q$ we get that $a_k(\lambda)a_l(\lambda)$ vanishes on a certain subvariety of codimension 1. On the other hand, this contradicts the fact that both a_k, a_l could only vanish on subvarieties of codimension more than one, since U^- has no zero divisors. Thus, the greatest common divisor of coefficients of $\Psi(\lambda)$ is one, which implies that $d_{k+l}(\lambda)$ is divisible by $f(\lambda)$.

This proves that $\Psi(\lambda) = c(q)q^{(\lambda,\alpha)}\tilde{\Phi}^{k+l}_\lambda$ for some $\alpha \in \frac{1}{2N}Q$ and rational function $c(q)$, independent of λ. To calculate α, c, let us consider the limit of both sides of (5.2.7) as $\lambda \to +\rho\infty$, i.e. letting $t_i = q^{\langle\lambda,\alpha_i^\vee\rangle/2N} = 0$.

LEMMA 5.2.4.

$$\lim_{\lambda\to\rho\infty}\Phi_k(v_\lambda) = v_\lambda \otimes u_0^k.$$

To prove the lemma, note first that $\lim\tilde{\Phi}^k = \lim\Phi^k$. Due to Lemma 5.2.1, we can write $\Phi(v_\lambda) = \sum_{k,l}(F^{-1})_{kl}g_kv_\lambda \otimes (\omega g_l)u$. It is known that if we choose a basis g_k in such a way that $g_0 = 1, g_k$ has strictly negative weight for $k > 0$, then $\lim_{\lambda\to\rho\infty}(F^{-1})_{kl} = \delta_{k,0}\delta_{l,0}$ (this follows, for example, from [L2, Proposition 19.3.7], which gives much more detailed information about the asymptotic behavior of F; it states that under a suitable normalization the Shapovalov form in the Verma module M_λ formally converges to the Drinfeld's form on U^- as $\lambda \to +\infty\rho$). This proves the lemma.

Using this lemma and the fact that in the identification $M_\lambda \simeq M \simeq U^-$ the action of U^- does not depend on λ, one can show that

$$\lim \Psi(v_\lambda) = v_\lambda \otimes u_{k+l}.$$

Comparing it with the the expression for $\lim \Phi_{k+l}(v_\lambda)$, we get the statement of the theorem. □

Corollary 5.2.5.

$$\Phi_\lambda^{k+l} = \frac{d_k(\lambda)d_l(\lambda)}{d_{k+l}(\lambda)}\Phi_\lambda^k * \Phi_\lambda^l. \tag{5.2.8}$$

So far, we have proved Theorem 5.2.3 only for the case when $k, l \in \mathbb{Z}_+$. However, it can be generalized. Let us consider the space $\tilde{U}_k = \{(x_1 \dots x_n)^k p(x), p(x) \in \mathbb{C}_q[x_1^{\pm 1}, \dots x_n^{\pm 1}], p(x)$ is a homogeneous polynomial of degree $0\}$ where k is an arbitrary complex number. Formula (4.2.1) defines an action of $U_q\mathfrak{sl}_n$ in $\tilde{U}_k$. Also, define $u_0^k = (x_1 \dots x_n)^k \in \tilde{U}_k$.

Lemma 5.2.6.

(1) *The set of weights of* $\tilde{U}_k$ *coincides with the weight lattice* Q, *and each weight subspace is one-dimensional. In particular,* $\tilde{U}_k[0] \simeq \mathbb{C}_q u_0^k$.

(2) *For generic* k, *the mapping*

$$x^\lambda \mapsto \frac{\Gamma_q(\lambda_1+1)\dots\Gamma_q(\lambda_n+1)}{(\Gamma_q(k+1))^n} x^{-1-\lambda}, \tag{5.2.9}$$

defines an isomorphism $\tilde{U}_k^\omega \simeq \tilde{U}_{-1-k}$. *The normalization is chosen so that* $u_0^k \mapsto u_0^{-1-k}$. *Here* $\Gamma_q(\lambda)$ *is q-gamma function:*

$$\Gamma_q(x) = \frac{1}{(1-q^2)^{x-1}} \prod_{n=0}^{\infty} \frac{1-q^{2(n+1)}}{1-q^{2(n+x)}},$$

so $\Gamma(x+1) = q^{x-1}[x]\Gamma(x)$, *where*

$$[x] = \frac{q^x - q^{-x}}{q - q^{-1}}.$$

Note that the factors of the form $(1-q^2)^\lambda$ *in the product in* (5.2.9) *cancel, and thus we can consider this product as a formal power series in* q *with coefficients which are rational functions in* q^λ, q^k (*which we consider as independent variables*).

(3) *If $k \in \mathbb{Z}_+$ then $\tilde{U}_k$ contains a finite-dimensional submodule, isomorphic to the module U_k defined above: $U_k = \tilde{U}_k \cap \mathbb{C}_q[x_1, \dots, x_n]$. Also, in this case $\tilde{U}_{-1-k}$ has a finite-dimensional quotient $U^k = \tilde{U}_{-1-k}/(x^\lambda$ such that at least one $\lambda_i \in \mathbb{Z}_+)$. In particular, U_{-1} can be projected onto $U^0 \simeq \mathbb{C}_q$. Moreover, formula* (5.2.9) *above defines an isomorphism $U_k^\omega \simeq U^k$ for $k \in \mathbb{Z}_+$.*

Proof of this lemma is straightforward.

Now, let us assume that λ is generic and consider an intertwiner $\Phi_\lambda^k : M_\lambda \to M_\lambda \hat{\otimes} \tilde{U}_k$ such that $\Phi_\lambda^k(v_\lambda) = v_\lambda \otimes u_0^k + \dots$, and $\hat{\otimes}$ is a tensor product completed with respect to ρ-grading in M_λ. Note that if $k \in \mathbb{Z}_+$ then image of $\Phi(v_\lambda)$ lies in the submodule $M_\lambda \otimes U_k$ (which follows from the explicit formula (5.2.2) for Φ), so this is consistent with our previous notations. Also, for $k \in \mathbb{C}$ we define

$$d_k(\lambda) = \prod_{\alpha \in R^+} \prod_{i=0}^{\infty} \frac{1 - q^{2(\alpha,\lambda+\rho)+2i} q^{-2k}}{1 - q^{2(\alpha,\lambda+\rho)+2i}}, \tag{5.2.10}$$

which we again consider as a formal power series in q with coefficients which are rational functions in q^λ, q^k. Note that if $k \in \mathbb{Z}_+$, this coincides with previously given definition.

Theorem 5.2.7. *For any $k \in \mathbb{Z}_+, l \in \mathbb{C}$ we have*

$$\Phi_\lambda^k * \Phi_\lambda^l = \frac{d_{k+l}(\lambda)}{d_k(\lambda) d_l(\lambda)} \Phi_\lambda^{k+l} = \Phi_\lambda^l * \Phi_\lambda^k, \tag{5.2.11}$$

where $d_k(\lambda)$ is given by formula (5.2.10).

Proof. Let us fix k. Then the matrix elements of the operators on both sides of (5.2.11) are rational functions in q, q^l, q^λ, which follows from the fact that $\frac{d_{k+l}(\lambda)}{d_k(\lambda) d_l(\lambda)}$ is a rational function in q, q^l, q^λ. Now the statement of the theorem follows from the fact that this is true for $l \in \mathbb{Z}_+$ and the following trivial statement:

If $F(q,t) \in \mathbb{C}(q,t)$ is such that $F(q, q^l) = 0$ for all $l \in \mathbb{Z}_+$ then $F = 0$.

□

Let us apply this to case when $l = -1 - k$. In this case explicit calculation gives the following answer:

Corollary 5.2.8. *For $k \in \mathbb{Z}_+$,*

$$\Phi_\lambda^k * \Phi_\lambda^{-1-k} = \Phi_\lambda^{-1} \prod_{\alpha \in R^+} \prod_{i=1}^{k} \frac{1 - q^{2(\alpha,\lambda+\rho)+2i}}{1 - q^{2(\alpha,\lambda+\rho)-2i}}.$$

Now, let us relate these intertwiners with intertwiners for finite-dimensional representations.

THEOREM 5.2.9. *Let* $\lambda, \mu \in P^+, u \in L_\mu[0], u \neq 0$ (*note that in this case* L_λ *is finite-dimensional*). *Assume that* λ *is such that the intertwiner* $\Phi_\lambda^{\mu,u} : M_\lambda \to M_\lambda \otimes L_\mu$ *defined in Lemma* 5.2.1 *is well-defined at this point, i.e. does not have a pole. Let* I_λ *be the maximal submodule in* M_λ: $L_\lambda = M_\lambda / I_\lambda$. *Then* $\Phi_\lambda^{\mu,u}(I_\lambda) \subset I_\lambda \otimes L_\mu$, *and thus,* $\Phi_\lambda^{\mu,u}$ *can be considered as an intertwiner* $L_\lambda \to L_\lambda \otimes L_\mu$. *Moreover, this is the unique intertwiner* $L_\lambda \to L_\lambda \otimes L_\mu$ *such that* $v_\lambda \mapsto v_\lambda \otimes u +$ *lower order terms.*

REMARK. Note that for $\lambda \in P^+$ the intertwiner $M_\lambda \to M_\lambda \otimes L_\mu$ such that $v_\lambda \mapsto v_\lambda \otimes u +$ *lower order terms* is not unique, so $\Phi_\lambda^{\mu,u}$ is a very special intertwiner of this form.

PROOF. Composing $\Phi_\lambda^{\mu,u}$ with the projection $M_\lambda \to L_\lambda$, we get an intertwiner $M_\lambda \to L_\lambda \otimes L_\mu$. Since the tensor product is finite-dimensional, this intertwiner must annihilate I_λ. Uniqueness can be proved in the same way as for M_λ for generic λ (see the proof of Lemma 5.2.1). $\square$

This shows that we can now apply the results of this section to study of products of intertwiners for finite-dimensional representations. In particular, we can now prove Theorem 5.1.3 form the previous section. Recall that it was formulated as follows:

THEOREM 5.1.3. *Let* $U = U_{k-1}$ *and let* $\Phi_\lambda : L_{\lambda^k} \to L_{\lambda^k} \otimes U, \Phi_\lambda^\circ : L_{\lambda^k} \to L_{\lambda^k} \otimes U^\omega$ *be such that* $\Phi_\lambda(v_{\lambda^k}) = v_{\lambda^k} \otimes u_0^{k-1}, \Phi_\lambda^\circ(v_{\lambda^k}) = v_{\lambda^k} \otimes (u_0^{k-1})^\omega$, *where* $\lambda^k = \lambda + (k-1)\rho, u_0 \in U[0], (u_0^{k-1})^\omega \in U^\omega[0]$. *Let* $(,)_U : U \otimes U^\omega \to \mathbb{C}_q$ *be Shapovalov form normalized so that* $(u_0, u_0^\omega)_U = 1$. *Then*

$$(\Phi_\lambda \Phi_\lambda^\circ)_U = \prod_{\alpha \in R^+} \prod_{i=1}^{k-1} \frac{1 - q^{2(\alpha, \lambda + k\rho) + 2i}}{1 - q^{2(\alpha, \lambda + k\rho) - 2i}} \operatorname{Id}_{L_{\lambda^k}}.$$

PROOF. Recall the notation $\tilde{U}_k$ (see Lemma 5.2.6). Then $U = U_{k-1}$ is a submodule in $\tilde{U}_{k-1}$, and U^ω is a factormodule of $\tilde{U}_{-k}$; moreover, the Shapovalov form coincides with the restriction on $U \otimes U^\omega$ of the map $\tilde{U}_{k-1} \otimes \tilde{U}_{-k} \to U_{-1} \to \mathbb{C}_q$. This together with Corollary 5.2.8 and Theorem 5.2.9 proves the desired statement.

5.3 Symmetry identities

In this section we only consider the case $\mathfrak{g} = \mathfrak{sl}_n$.

The main goal of this section is to prove Theorem 5.3.3, which establishes certain symmetry between the values $P_\lambda(q^{2(\mu+k\rho)})$ and $P_\mu(q^{2(\lambda+k\rho)})$ (notations will be explained later). The proof of this theorem is based on the technique of ribbon graphs.

As before, let $\Phi_\lambda : L_{\lambda^k} \to L_{\lambda^k} \otimes U_{k-1}$ be such that $\Phi(v_{\lambda^k}) = v_{\lambda^k} \otimes u_0^{k-1} + \dots$, $\lambda^k = \lambda + (k-1)\rho$, and let $\varphi_\lambda = \chi_{\Phi_\lambda}$ be the corresponding generalized character.

Lemma 5.3.1.

(5.3.1)



$$= \varphi_\mu(q^{2(\lambda+k\rho)})\Phi_\lambda.$$

where $\varphi_\mu(q^\lambda)$ *stands for polynomial in* q, q^{-1} *which is obtained by replacing each formal exponent* e^α *in the expression for* φ_μ *by* $q^{(\alpha,\lambda)}$.

Proof. Let us consider the operator $F : L_{\lambda^k} \to L_{\lambda^k} \otimes U_{k-1}$ corresponding to the ribbon graph on the left hand side of (5.3.1). It is some $U_q\mathfrak{g}$-homomorphism. Since we know that such a homomorphism is unique up to a constant, it follows that $F = a\Phi_\lambda$ for some constant a. To find a, let us find the image of the highest-weight vector. First, consider the following part of this picture:



The corresponding operator is the product $R^{21}R : L_{\lambda^k} \otimes (L_{\mu^k})^* \to L_{\lambda^k} \otimes (L_{\mu^k})^*$. It follows from the explicit form of R-matrix (1.2.7) that if $x \in L^*_{\mu^k}[\alpha]$ then $R^{21}R(v_{\lambda^k} \otimes x) = q^{-2(\lambda^k,\alpha)}v_{\lambda^k} \otimes x + \dots$. Thus, if x_i is basis in L_{μ^k}, x^i – dual basis in $L^*_{\mu^k}$, x_i has weight α_i then explicit calculation shows that



$$: v_{\lambda^k} \mapsto \sum_i q^{2(\alpha_i,\lambda+k\rho)} v_{\lambda^k} \otimes x^i \otimes \Phi(x_i) + l.o.t.$$

and thus, $F(v_{\lambda^k}) = \varphi_\mu(q^{2(\lambda+k\rho)})v_{\lambda^k} \otimes u_0^{k-1} + \dots$, which completes the proof. □

Remark. In the case $k = 1$, i.e. $U = \mathbb{C}$, (5.3.1) reduces to the formula for the value of the central element $c_{L^*_{\mu^k}}$ in L_{λ^k} (see Theorem 1.2.1)

Corollary 5.3.2. *Let* $\Phi_\lambda : L_{\lambda^k} \to L_{\lambda^k} \otimes U_{k-1}, \Phi^\circ_\lambda : L_{\lambda^k} \to L_{\lambda^k} \otimes U^\omega_{k-1}$ *be as in Theorem* 5.1.2. *Then*

(5.3.2)



$$= \varphi_\mu(q^{2(\lambda+k\rho)})\langle P_\lambda, P_\lambda\rangle_k \dim_q L_{\lambda^k}.$$

Proof. This follows from the previous lemma and the arguments used in the proof of Theorem 5.1.2. □

In a similar way, we can replace everywhere U by U^* (see Remark 4.2.4), and repeating with necessary changes all the steps of previous sections prove the following theorem:

Theorem 5.3.3. *Formula* (5.3.2) *remains valid if we replace in the graph on the left hand side* Φ_μ *by* Φ°_μ*,* Φ°_λ *by* Φ_λ *and interchange* ω_1 *and* ω_1^*.

Theorem 5.3.4.
(5.3.3)

$$\frac{P_\mu(q^{2(\lambda+k\rho)})}{P_\lambda(q^{2(\mu+k\rho)})} = q^{2k(\rho,\lambda-\mu)} \prod_{\alpha\in R^+}\prod_{i=0}^{k-1} \frac{1-q^{2(\alpha,\mu+k\rho)+2i}}{1-q^{2(\alpha,\lambda+k\rho)+2i}} = \prod_{\alpha\in R^+}\prod_{i=0}^{k-1} \frac{[(\alpha,\mu+k\rho)+i]}{[(\alpha,\lambda+k\rho)+i]}.$$

Proof. The proof is based on the following identity of the ribbon graphs:

(5.3.4)



Due to Corollary 5.3.2 and Theorem 5.3.3, this implies:

$$\varphi_\mu(q^{2(\lambda+k\rho)})\langle P_\lambda, P_\lambda\rangle_k \dim_q L_{\lambda^k} = \varphi_\lambda(q^{2(\mu+k\rho)})\langle P_\mu, P_\mu\rangle_k \dim_q L_{\mu^k}. \tag{5.3.5}$$

Substituting in this formula explicit expression for $\langle P_\lambda, P_\lambda\rangle_k$ (Theorem 5.1.4) and using the following expression for $\dim_q L_\lambda$:

$$\dim_q L_\lambda = q^{-2(\lambda,\rho)} \prod_{\alpha\in R^+} \frac{1-q^{2(\alpha,\lambda+\rho)}}{1-q^{2(\alpha,\rho)}} = \prod_{\alpha\in R^+} \frac{[(\alpha,\lambda+\rho)]}{[(\alpha,\rho)]},$$

which can be easily deduced from the Weyl character formula, we get the statement of the theorem. □

Corollary 5.3.5. (Macdonald's special value identity, [M1,M2])
(5.3.6)

$$P_\lambda(q^{2k\rho}) = q^{-2k(\rho,\lambda)} \prod_{\alpha\in R^+}\prod_{i=0}^{k-1} \frac{1-q^{2(\alpha,\lambda+k\rho)+2i}}{1-q^{2(\alpha,k\rho)+2i}} = \prod_{\alpha\in R^+}\prod_{i=0}^{k-1} \frac{[(\alpha,\lambda+k\rho)+i]}{[(\alpha,k\rho)+i]},$$

Proof. Let $\mu=0$. Then $P_\mu=1$, and formula (5.3.3) reduces to (5.3.6). □

We refer the reader to [EK5] for the generalization of this formula for arbitrary k.

CHAPTER VI

GENERALIZED CHARACTERS FOR AFFINE LIE ALGEBRAS

In this chapter we start the study of the generalized characters for the affine Lie algebras. Mostly, the constructions are parallel to the finite-dimensional case, but since the objects we work with are infinite-dimensional, extra care should be taken.

6.1 Affine Lie algebras

Here we review the notations and facts about affine Lie algebras and root systems. All of them can be found in [Ka1]. We keep the notations of Chapter 1; as a rule, we will use hat (̂) in the notations of affine analogues of finite-dimensional objects.

Let $\hat{\mathfrak{g}}$ be the affine Lie algebra corresponding to $\mathfrak{g}$:

$$\hat{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}c \oplus \mathbb{C}d,$$

with the commutation rule given by

$$\begin{gathered} [x \otimes t^n, y \otimes t^m] = [x, y] \otimes t^{m+n} + n\delta_{m,-n}(x, y)c, \\ c \text{ is central}, \\ [d, x \otimes t^n] = nx \otimes t^n. \end{gathered} \tag{6.1.1}$$

Sometimes we will use a smaller algebra $\tilde{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}c$. For brevity, we will use the notation

$$x \otimes t^n = x[n], \quad x \in \mathfrak{g}, n \in \mathbb{Z}.$$

Similarly to the finite-dimensional case, we define Cartan subalgebra $\hat{\mathfrak{h}} = \mathfrak{h} \oplus \mathbb{C}c \oplus \mathbb{C}d$, $\hat{\mathfrak{h}}^* = \hat{\mathfrak{h}}^* \oplus \mathbb{C}\delta \oplus \mathbb{C}\Lambda_0$, where $\langle \Lambda_0, \mathfrak{h} \oplus \mathbb{C}d \rangle = \langle \delta, \mathfrak{h} \oplus \mathbb{C}c \rangle = 0$, $\langle \delta, d \rangle = 1, \langle \Lambda_0, c \rangle = 1$. It will be convenient to consider affine hyperplanes $\hat{\mathfrak{h}}^*_K = \mathfrak{h}^* \oplus \mathbb{C}\delta + K\Lambda_0, K \in \mathbb{C}$; we will refer to the elements of $\hat{\mathfrak{h}}^*_K$ as having level K.

Again, we have a bilinear non-degenerate symmetric form $(\cdot, \cdot)$ on $\hat{\mathfrak{h}}^*$ which coincides with previously defined on $\mathfrak{h}^*$ and $(\Lambda_0, \delta) = 1, (\Lambda_0, \mathfrak{h}^*) = (\delta, \mathfrak{h}^*) = (\Lambda_0, \Lambda_0) =$

$(\delta,\delta)=0$. This gives an identification $\hat{\mathfrak{h}}^*\simeq\hat{\mathfrak{h}}:\lambda\mapsto h_\lambda$, and a bilinear form on $\hat{\mathfrak{h}}$ such that $(c,d)=1$. Note that under this identification, $Q^\vee\subset Q$.

We define the affine root system $\widehat{R}=\{\hat\alpha=\alpha+n\delta|\alpha\in R, n\in\mathbb{Z}$ or $\alpha=0, n\in\mathbb{Z}\setminus\{0\}\}$. Again, we have the notion of positive roots: $\widehat{R}^+=\{\hat\alpha=\alpha+n\delta\in\widehat{R}|n>0$ or $n=0,\alpha\in R^+\}$ and the basis of simple roots $\alpha_0=-\theta+\delta,\alpha_1,\dots,\alpha_r$. For $\hat\alpha=\alpha+n\delta\in\widehat{R}^+$ we define $e_{\hat\alpha}=e_\alpha[n]$ if $\alpha\in R^+$, and $e_\alpha=f_{-\alpha}[n]$ if $\alpha\in R^-$; $f_{\hat\alpha}$ are defined similarly. As before, we use the notations $e_i=2e_{\alpha_i}/(\alpha_i,\alpha_i), f_i=2f_{\alpha_i}/(\alpha_i,\alpha_i), h_i=\alpha_i^\vee=2h_{\alpha_i}/(\alpha_i,\alpha_i), i=0,\dots,r$. This gives the polarization of $\hat{\mathfrak{g}}$: $\hat{\mathfrak{g}}=\hat{\mathfrak{n}}^+\oplus\hat{\mathfrak{h}}\oplus\hat{\mathfrak{n}}^-$.

As usual, we denote $\hat{Q}=\bigoplus\mathbb{Z}\alpha_i, \hat{Q}^+=\bigoplus\mathbb{Z}_+\alpha_i$.

We define the affine Weyl group $\widehat{W}$ as the group of transformations of $\hat{\mathfrak{h}}^*$ generated by the reflections with respect to $\alpha_i, i=0\dots r$. We have notion of sign of an element of $\widehat{W}$: $\varepsilon(w)=(-1)^l$ if w is a product of l reflections. This group preserves the bilinear form; also, it preserves each of the affine hyperplanes $\hat{\mathfrak{h}}^*_K$.

Theorem 6.1.1. (see [Ka1]) *$\widehat{W}\simeq W\ltimes Q^\vee$, where the action of W is the same as in the classical case, and the action of $Q^\vee$ in $\hat{\mathfrak{h}}^*_K$ is given by*

$$\alpha^\vee:\widehat{\lambda}\mapsto\widehat{\lambda}+K\alpha^\vee-\big(\langle\widehat{\lambda},\alpha^\vee\rangle+\frac{1}{2}K(\alpha^\vee,\alpha^\vee)\big)\delta. \tag{6.1.2}$$

Now we can define the root lattice $\widehat{P}=P\oplus\mathbb{Z}\delta\oplus\mathbb{Z}\Lambda_0\subset\hat{\mathfrak{h}}^*$ and the cone of dominant weights $\widehat{P}^+=\{\widehat{\lambda}\in\hat{\mathfrak{h}}^*|\langle\widehat{\lambda},\alpha_i^\vee\rangle\in\mathbb{Z}_+, i=0,\dots,r\}$. We will also use the notation $\widehat{P}^+_K=\widehat{P}^+\cap\hat{\mathfrak{h}}^*_K=\{\lambda+n\delta+K\Lambda_0|\lambda\in P^+,(\lambda,\theta)\le K\}$. Note the cone of dominant weights is invariant with respect to the translations along δ direction, but if one factors this out then there is only a finite number of dominant weights for every level K. Abusing the notations, we will write $P^+_K=\{\lambda\in P^+|(\lambda,\theta)\le K\}$. We introduce an order on $\widehat{P}$ as usual: $\widehat{\lambda}\le\hat\mu\iff\hat\mu-\widehat{\lambda}\in\hat{Q}^+$.

We define the following affine analogue of ρ:

$$\hat\rho=\rho+h^\vee\Lambda_0; \tag{6.1.3}$$

then $\langle\hat\rho,\alpha_i^\vee\rangle=1, i=0,\dots,r$ and thus $\hat\rho\in\widehat{P}^+$.

Lemma 6.1.2.

(1) *$\widehat{W}$ preserves each $\widehat{P}_K=\widehat{P}\cap\hat{\mathfrak{h}}^*_K$.*

(2) *$\widehat{P}^+_K$ is a fundamental domain for the action of $\widehat{W}$ in $\widehat{P}_K$ for $K>0$.*

Proof. The proof is based on the fact that when we restrict the bilinear form $(\ ,\ )$ to $Q^\vee$ we get an integer even lattice – see [Lo].

Let us consider representations of $\hat{\mathfrak{g}}$. Unless otherwise specified, we only consider representations with weight decomposition. We say that a module V over $\hat{\mathfrak{g}}$ is of level K if $c|_V=K\,\mathrm{Id}_V$.

Since we have polarization of $\hat{\mathfrak{g}}$, we can define Verma modules $M_{\widehat{\lambda}}$ over $\hat{\mathfrak{g}}$ and their irreducible quotients $L_{\widehat{\lambda}}$. Note that $L_{\widehat{\lambda}}$ is infinite-dimensional unless $\widehat{\lambda} = a\delta$. Both of these modules admit weight decomposition.

As usual, we define the category $\mathcal{O}$ to be the category of finitely-generated $\hat{\mathfrak{g}}$-modules with weight decomposition such that for every $v \in V$ the space $U\hat{\mathfrak{n}}^+ v$ is finite-dimensional. Obviously, for every $\widehat{\lambda}$ the modules $M_{\widehat{\lambda}}, L_{\widehat{\lambda}}$ satisfy this condition.

We say that a module V from category $\mathcal{O}$ is integrable if for every simple root $\alpha_i, i = 0, \dots, r$ restriction of V to the Lie algebra $\mathfrak{sl}_2$ generated by e_i, f_i, h_i is a direct sum of finite dimensional modules. Integrable modules are natural analogues of finite-dimensional modules for $\mathfrak{g}$. It is known that $L_{\widehat{\lambda}}$ is integrable iff $\widehat{\lambda} \in \widehat{P}^+$, and that every irreducible integrable module has the form $L_{\widehat{\lambda}}$ for some $\widehat{\lambda} \in \widehat{P}^+$.

We will also need another type of modules, which are called evaluation representations. Let V be a finite-dimensional module over $\mathfrak{g}$ and let z be a non-zero complex number. Then we can construct an evaluation representation of $\tilde{\mathfrak{g}}$ (not $\hat{\mathfrak{g}}$!) in V by

$$\begin{gathered} \pi_{V(z)}(a[n]) = z^n \pi_V(a), \\ \pi_{V(z)}(c) = 0. \end{gathered} \tag{6.1.4}$$

Note that $V(z)$ has no $\widehat{P}$-grading but has a natural P-grading and $V(z)$ is a module of level zero.

We will be interested in intertwining operators of the following form, which are sometimes called vertex operators:

$$\Phi\colon L_{\widehat{\lambda}} \to \widehat{L_{\widehat{\lambda}}} \otimes V(z), \tag{6.1.5}$$

where $\widehat{L_{\widehat{\lambda}}}$ is the completion of the integrable highest-weight module $L_{\widehat{\lambda}}, \widehat{\lambda} \in \widehat{P}^+$ with respect to the d-grading. To prove the existence of such intertwiners, we use the following well-known result (see, for example, the arguments in [TK], which work for general Lie algebra in the same way as for $\mathfrak{sl}_2$):

Lemma 6.1.3. *The mapping* $\Phi \mapsto (v_{\widehat{\lambda}}, \Phi v_{\widehat{\lambda}})$ *establishes one-to-one correspondence between the space of all intertwiners of the form* (6.1.5) *and the subspace in* $V[0]$ *formed by the vectors* v *such that* $xv = 0$ *for every* $x \in U\hat{\mathfrak{n}}^-$ *such that* $xv_{\widehat{\lambda}} = 0$ *in* $L_{\widehat{\lambda}}$.

6.2 Group algebra of the weight lattice

One of the most important objects of study for us will be the algebra of $\widehat{W}$-invariants of the (suitably completed) group algebra of $\widehat{P}$. In the affine case its definition is much more subtle than in the finite-dimensional case; our exposition follows the paper of Looijenga ([Lo]).

Let us consider the group algebra $\mathbb{C}[\widehat{P}]$, i.e. the algebra spanned by the formal exponentials $e^{\widehat{\lambda}}, \widehat{\lambda} \in \widehat{P}$. It is naturally $\mathbb{Z}$-graded: $\mathbb{C}[\widehat{P}] = \bigoplus_{K \in \mathbb{Z}} \mathbb{C}[\widehat{P}_K]$. Consider the following completion:

(6.2.1)

$$\overline{\mathbb{C}[\widehat{P}_K]} = \left\{ \sum_{\widehat{\lambda} \in \widehat{P}_K} a_{\widehat{\lambda}} e^{\widehat{\lambda}} \,\middle|\, \text{for every } N \in \mathbb{Z}, \text{ there exists only finitely many } \widehat{\lambda} \in \widehat{P}_K \right.$$
$$\left. \text{such that } a_{\widehat{\lambda}} \neq 0 \text{ and } (\widehat{\lambda}, \hat{\rho}) \geq N \right\}.$$

Then $\overline{\mathbb{C}[\widehat{P}]} = \bigoplus_K \overline{\mathbb{C}[\widehat{P}_K]}$ is again a $\mathbb{Z}$-graded algebra. This completion is chosen so to include the characters of Verma modules (and more generally, modules from category $\mathcal{O}$) over $\hat{\mathfrak{g}}$. This algebra has a natural topology with the basis of neighborhoods of zero given by

$$X_N = \left\{ \sum_{\widehat{\lambda} \in \widehat{P}_K} a_{\widehat{\lambda}} e^{\widehat{\lambda}} \in \overline{\mathbb{C}[\widehat{P}_K]} \,\middle|\, (\widehat{\lambda}, \hat{\rho}) < -N \text{ if } a_{\widehat{\lambda}} \neq 0 \right\}.$$

For our purposes this algebra is too big. We will use a smaller algebra:

$$A = \bigcap_{w \in \widehat{W}} w\left(\overline{\mathbb{C}[\widehat{P}]}\right), \tag{6.2.2}$$

which is a natural analogue of the group algebra $\mathbb{C}[P]$ for finite-dimensional case. In particular, we have a natural action of $\widehat{W}$ in A. It is also $\mathbb{Z}$-graded: $A_K = \bigcap_{w \in \widehat{W}} w\left(\overline{\mathbb{C}[\widehat{P}_K]}\right)$. Again, this algebra has a topology, which we describe in terms of convergence: we say that $f_n \to 0$ in A if for every $w \in \widehat{W}$, $w(f_n) \to 0$ in $\overline{\mathbb{C}[\widehat{P}]}$.

One of the main objects of our study will be the algebra of $\widehat{W}$-invariants $A^{\widehat{W}} \subset A$. We will call elements of $A_K^{\widehat{W}}$ (formal) symmetric theta-functions of level K; indeed, we will show below that for $\mathfrak{g} = \mathfrak{sl}_2$ the condition of $\widehat{W}$-invariance coincides with definition of even theta-function.

For a module V with the weight decomposition define its character by the formula: $\operatorname{ch} V = \sum_{\widehat{\lambda} \in \widehat{P}} e^{\widehat{\lambda}} \dim V[\widehat{\lambda}]$ provided that this sum converges in the sense of the completion $\overline{\mathbb{C}[\widehat{P}]}$.

Example 1. For any $\widehat{\lambda} \in \widehat{P}_K, K \geq 0$ the orbitsum

$$m_{\widehat{\lambda}} = \sum_{\hat{\mu} \in \widehat{W}\widehat{\lambda}} e^{\hat{\mu}} \tag{6.2.3}$$

belongs to $A_K^{\widehat{W}}$.

Example 2. For any module V from the category $\mathcal{O}$ (in particular, for $M_{\widehat{\lambda}}, L_{\widehat{\lambda}}$) we have $\operatorname{ch} V \in \overline{\mathbb{C}[\widehat{P}]}$.

Example 3. If V is a module from the category $\mathcal{O}$ then $\operatorname{ch} V \in A$ iff V is integrable, in which case $\operatorname{ch} V \in A^W$. In particular, $\operatorname{ch} L_{\widehat{\lambda}} \in A^{\widehat{W}}$ iff $\widehat{\lambda} \in \widehat{P}^+$.

Example 4. Define the affine Weyl denominator $\hat{\delta}$ by

$$\hat{\delta} = e^{\hat{\rho}} \prod_{\hat{\alpha} \in \widehat{R}^+} (1 - e^{-\hat{\alpha}}) = \frac{1}{\operatorname{ch} M_{-\hat{\rho}}}. \tag{6.2.4}$$

Then $\hat{\delta} \in A$. It is easy to see that $\hat{\delta}$ is $\widehat{W}$-antiinvariant; moreover, it is known (see [Ka1, Lo]) that every $\widehat{W}$-antiinvariant element of A has the form $f\hat{\delta}, f \in A^{\widehat{W}}$.

Obviously, $A^{\widehat{W}}$ is $\mathbb{Z}$-graded. Moreover, the following is well-known:

Lemma 6.2.1. *$A_K^{\widehat{W}} = 0$ for $K < 0$, and*

$$A_0^{\widehat{W}} = \left\{ \sum_{n \leq n_0} a_n e^{n\delta}, a_n \in \mathbb{C} \right\}. \tag{6.2.5}$$

Theorem 6.2.2. (cf. [Lo]) *For every $K \in \mathbb{Z}_+$, the orbitsums $m_{\lambda + K\Lambda_0}, \lambda \in P_K^+$ form a basis of $A_K^{\widehat{W}}$ over the field $A_0^{\widehat{W}}$.*

This theorem follows from the fact that $\widehat{P}_K^+$ is a fundamental domain for the action of $\widehat{W}$ in $\widehat{P}_K$ for $K > 0$ and from Lemma 6.2.1.

It will be convenient to introduce formal variable $p = e^{-\delta}$; then every element of A can be written as a formal Laurent series in p with coefficients from $\mathbb{C}[P]$ (note that this is not so for $\overline{\mathbb{C}[\widehat{P}]}$). It is also compatible with convergence: if $f_n \to 0$ in A then $f_n \to 0$ in the usual p-adic topology in $\mathbb{C}((p))[P]$. In particular, in these notations $A_0^{\widehat{W}} \simeq \mathbb{C}((p))$. (Usually, $e^{-\delta}$ is denoted by q; we use the letter p to avoid confusion with the parameter q of the quantum group).

We will also need the analogues of the ring $\mathbb{C}[P](e^{\alpha} - 1)^{-1}$, considered in Section 2.2. First, note that if $\hat{\alpha} \in \widehat{R}^-$ then the series $(1 - e^{\hat{\alpha}})^{-1} = 1 + e^{\hat{\alpha}} + e^{2\hat{\alpha}} + \dots$ converges in $\overline{\mathbb{C}[\widehat{P}]}$, so there is no need to extend $\overline{\mathbb{C}[\widehat{P}]}$ any further.

As for the algebra A defined above, it does not contain $(1 - e^{\hat{\alpha}})^{-1}$. For this reason, we define the algebra $\hat{\mathcal{R}}$ as follows. Consider the algebra $\mathbb{C}[\widehat{P}](1 - e^{\hat{\alpha}})^{-1}$, obtained by adjoining to $\mathbb{C}[\widehat{P}]$ the inverses of $(1 - e^{\hat{\alpha}})$ (no completion so far). Then we have a morphism

$$\tau\colon \mathbb{C}[\widehat{P}](1-e^{\hat\alpha})^{-1} \to \overline{\mathbb{C}[\widehat{P}]}, \tag{6.2.6}$$

given by expanding $(1-e^{-\hat\alpha})^{-1} = 1 + e^{-\hat\alpha} + e^{-2\hat\alpha} + \dots$ for $\hat\alpha \in \widehat{R}^+$. Note that the image is not in A.

Similarly, for every $w \in \widehat{W}$ we have

$$\tau_w\colon \mathbb{C}[\widehat{P}](1-e^{\alpha})^{-1} \to w\Bigl(\overline{\mathbb{C}[\widehat{P}]}\Bigr),$$

given by expanding $(1-e^{-\alpha})^{-1} = 1 + e^{-\alpha} + \dots$ for $\alpha \in w\widehat{R}^+$.

Define

$$\begin{aligned}\hat{\mathcal{R}} = \Bigl\{ \sum a_n \mid {} & a_n \in \mathbb{C}[\widehat{P}](1-e^{\alpha})^{-1}, \\ & \sum \tau_w(a_n) \text{ converges in } w\Bigl(\overline{\mathbb{C}[\widehat{P}]}\Bigr) \text{ for every } w \in \widehat{W} \Bigr\}.\end{aligned} \tag{6.2.7}$$

This algebra obviously contains A. This is the right analogue of the ring of fractions $\mathbb{C}[P](e^{\alpha}-1)^{-1}$, introduced in Section 2.2; for example, $\sum_{\hat\alpha \in \widehat{R}^+} \frac{1}{1-e^{\hat\alpha}} \in \hat{\mathcal{R}}$. Note that there is a natural action of the Weyl group $\widehat{W}$ in $\hat{\mathcal{R}}$; also note that this algebra has a natural $\mathbb{Z}$-grading given by level.

6.3 Generalized characters for affine Lie algebras

In this section, we define and study some properties of generalized characters for affine Lie algebras. Results of this section are due to the author and P. Etingof ([E1, EK3, EK4]); in less general situation (and different language) some of the results were found earlier by Bernard ([Be]).

Definition 6.3.1. Let $\Phi : V \to V \otimes U$ be a $\tilde{\mathfrak{g}}$-intertwiner, where V is a module from the category $\mathcal{O}$ and U is a finite-dimensional $\tilde{\mathfrak{g}}$-module of level zero. Define the corresponding generalized character χ_Φ by

$$\chi_\Phi = \sum_{\widehat{\lambda} \in \widehat{P}} e^{\widehat{\lambda}} \operatorname{Tr}_{V[\widehat{\lambda}]} \Phi \in \overline{\mathbb{C}[\widehat{P}]} \otimes U[0]. \tag{6.3.1}$$

Remark. It is easy to see that if V is integrable then $\chi_\Phi \in A$.

As before, we can interpret generalized characters as functions of variables $h \in \mathfrak{h}, p \in \mathbb{C}$ by the rule

$$\chi_\Phi(h,p) = \mathrm{Tr}_V(\Phi p^{-d} e^{2\pi i h}), \tag{6.3.2}$$

provided that this sum converges. In the cases we will be interested in this sum does converge in a certain region (typically, $0 < |p| < 1$); however, this requires proof, which we will give later (see Section 8.1). For this reason, at the moment we consider generalized characters formally, i.e. as elements of the completed group algebra of the weight lattice.

Our next goal is to deduce some differential equations satisfied by these generalized characters. Unfortunately, we can not use central elements of $U\hat{\mathfrak{g}}$ for this purpose, since there are no non-trivial central elements in $U\hat{\mathfrak{g}}$ except c. However, we can introduce the following central element in the completion of $U\hat{\mathfrak{g}}$:

(6.3.3)

$$\widehat{C} = 2(c+h^\vee)d + \sum_{n\in\mathbb{Z}} \left(\sum_{\alpha\in R^+} (: e_\alpha[n] f_\alpha[-n] : + : f_\alpha[n] e_\alpha[-n] :) + \sum : x_l[n] x_l[-n] : \right),$$

where x_l is an orthonormal basis in $\mathfrak{h}$ with respect to $(\ ,\ )$ and the normal ordering is defined by

$$: x[n]y[m] : = \begin{cases} x[n]y[m] & m \geq n, \\ y[m]x[n] & m < n. \end{cases} \tag{6.3.4}$$

This is the affine analogue of the Casimir element. Though it is an infinite sum, it is well defined in every module from category $\mathcal{O}$, and $\hat{C}|_{M_{\widehat{\lambda}}} = (\widehat{\lambda}, \widehat{\lambda} + 2\hat{\rho})\,\mathrm{Id}_{M_{\widehat{\lambda}}}$.

Now we can formulate the main result of this section, which is the affine analogue of Theorem 2.2.4. Similarly to Section 2.2, let us introduce the following differential operators, which we consider formally, i.e. as derivations of the algebra $\mathbb{C}[\widehat{P}]$:

$$\begin{gathered} \hat{\Delta} e^{\widehat{\lambda}} = (\widehat{\lambda}, \widehat{\lambda}) e^{\widehat{\lambda}}, \\ \partial_{\hat{\alpha}} e^{\widehat{\lambda}} = (\hat{\alpha}, \widehat{\lambda}) e^{\widehat{\lambda}}, \quad \hat{\alpha} \in \hat{\mathfrak{h}}^*. \end{gathered} \tag{6.3.5}$$

If we use the notation $e^{-\delta} = p$ and write elements of $\mathbb{C}[\widehat{P}_K]$ as functions of p with coefficients from $\mathbb{C}[P]$: $\mathbb{C}[\widehat{P}_K] = e^{K\Lambda_0}\mathbb{C}[p, p^{-1}][P]$ then

$$\begin{gathered} \hat{\Delta} = -2Kp\frac{\partial}{\partial p} + \Delta_{\mathfrak{h}}, \\ \partial_{\alpha+n\delta} = \partial_\alpha + nK, \end{gathered} \tag{6.3.6}$$

where $\Delta_{\mathfrak{h}}$ is the Laplace operator in $\mathfrak{h}$ defined in Section 2.2.

Note also that both $\hat{\Delta}$ and $\partial_{\hat{\alpha}}$ can be extended to the completions $\overline{\mathbb{C}[\widehat{P}]}$ and A.

THEOREM 6.3.2. ([E1]) *Let $\widehat{\lambda} \in \widehat{P}^+$, and let $\Phi : L_{\widehat{\lambda}} \to L_{\widehat{\lambda}} \otimes U$ be a $\tilde{\mathfrak{g}}$-intertwiner, where U is as in Definition* 6.3.1. *Denote by $\chi_\Phi \in A \otimes U[0]$ the corresponding generalized character. Then χ_Φ satisfies the following differential equation:*

$$\left(\hat{\Delta} - \sum_{\hat{\alpha} \in \widehat{R}^+} \frac{e_{\hat{\alpha}} f_{\hat{\alpha}} + f_{\hat{\alpha}} e_{\hat{\alpha}}}{(e^{\hat{\alpha}/2} - e^{-\hat{\alpha}/2})^2} + 2\partial_{\hat{\rho}} - 2 \sum_{\hat{\alpha} \in \widehat{R}^+} \frac{1}{1 - e^{\hat{\alpha}}} \partial_{\hat{\alpha}}\right) \chi_\Phi = (\widehat{\lambda}, \widehat{\lambda} + 2\hat{\rho}) \chi_\Phi, \tag{6.3.7}$$

where we use the following convention: if $\alpha = n\delta \in \widehat{R}^+$ then $e_{\hat{\alpha}} f_{\hat{\alpha}} + f_{\hat{\alpha}} e_{\hat{\alpha}}$ should be replaced by $\sum_l x_l[n] x_l[-n]$, where x_l is an orthonormal basis in $\mathfrak{h}$. Similarly, in the second sum each root should be taken with multiplicity: for $\hat{\alpha} = n\delta$ the term $\frac{1}{1-e^{\hat{\alpha}}} \partial_{\hat{\alpha}}$ must be multiplied by r.

This equation can be rewritten in the following form: if χ_Φ has level K then

$$\begin{aligned} &\left(\hat{\Delta} - \sum_{\hat{\alpha} \in \widehat{R}^+} \frac{e_{\hat{\alpha}} f_{\hat{\alpha}} + f_{\hat{\alpha}} e_{\hat{\alpha}}}{(e^{\hat{\alpha}/2} - e^{-\hat{\alpha}/2})^2}\right) (\hat{\delta} \chi_\Phi) \\ &= \Bigg(\Delta_{\mathfrak{h}} - 2(K + h^\vee) p \frac{\partial}{\partial p} \\ &\qquad - \sum_{n \in \mathbb{Z}} \Bigg(\sum_{\alpha \in R^+} \frac{p^n e^\alpha}{(1 - p^n e^\alpha)^2} (f_\alpha[n] e_\alpha[-n] + e_\alpha[-n] f_\alpha[n]) \\ &\qquad\qquad + \sum_l \frac{p^n}{(1 - p^n)^2} x_l[n] x_l[-n] \Bigg)\Bigg) (\hat{\delta} \chi_\Phi) \\ &= (\widehat{\lambda} + \hat{\rho}, \widehat{\lambda} + \hat{\rho}) \hat{\delta} \chi_\Phi, \end{aligned} \tag{6.3.8}$$

where $\hat{\delta}$ is the affine Weyl denominator (6.2.4).

In less general form this equation has first appeared in the papers of Bernard ([Be]).

REMARK. It is easy to check that the differential operators written in the right-hand side of (6.3.7), (6.3.8) are well defined as operators in $\hat{\mathcal{R}}$.

PROOF. The proof is analogous to the finite-dimensional case (Proposition 2.2.4). It is based on the following identities, which can be easily proved by the same methods as their finite-dimensional analogues (2.2.5–2.2.7):

$$\text{(6.3.9)} \qquad \operatorname{Tr}(\Phi e_{\hat\alpha} f_{\hat\alpha} p^{-d} e^{2\pi \mathrm{i} h}) = \left(-\frac{e^{\hat\alpha}}{(1-e^{\hat\alpha})^2} e_{\hat\alpha} f_{\hat\alpha} - \frac{e_{\hat\alpha}}{1-e^{\hat\alpha}} \partial_{\hat\alpha} \right) \chi_\Phi,$$

$$\text{(6.3.10)} \qquad \operatorname{Tr}(\Phi f_{\hat\alpha} e_{\hat\alpha} p^{-d} e^{2\pi \mathrm{i} h}) = \left(-\frac{e^{\hat\alpha}}{(1-e^{\hat\alpha})^2} f_{\hat\alpha} e_{\hat\alpha} - \frac{1}{1-e^{\hat\alpha}} \partial_{\hat\alpha} \right) \chi_\Phi,$$

$$\text{(6.3.11)} \qquad \operatorname{Tr}(\Phi(2cd + \sum x_l^2) p^{-d} e^{2\pi \mathrm{i} h}) = \hat\Delta \chi_\Phi.$$

Since $\hat C|_{L_{\widehat\lambda}} = (\widehat\lambda, \widehat\lambda + 2\hat\rho)\,\mathrm{Id}_{L_{\widehat\lambda}}$, we have $\operatorname{Tr}(\Phi \hat C p^{-d} e^{2\pi \mathrm{i} h}) = (\widehat\lambda, \widehat\lambda + 2\hat\rho)\chi_\Phi$. Substituting in this formula the expression (6.3.3) for $\hat C$ and using identities (6.3.9–6.3.11), we obtain the desired equation (6.3.7). Formula (6.3.8) can be deduced straightforwardly; it also follows from more general statement, which we will prove in the next chapter (Theorem 7.1.2).

Example 6.3.3. Assume that $U = U(z)$ is an evaluation representation. Then $\chi_\Phi \in U[0]$, and thus, independently of z we have $x_l[n]\chi_\Phi = 0, e_{\hat\alpha} f_{\hat\alpha} \chi_\Phi = f_{\hat\alpha} e_{\hat\alpha} \chi_\Phi = e_\alpha f_\alpha \chi_\Phi$. Therefore, we can rewrite (6.3.8) as follows:

$$\text{(6.3.12)} \ \left(\Delta_h - 2(K + h^\vee) p \frac{\partial}{\partial p} - 2 \sum_{\substack{n \in \mathbb{Z} \\ \alpha \in R^+}} \frac{p^n e^\alpha}{(1 - p^n e^\alpha)^2} e_\alpha f_\alpha \right) (\hat\delta \chi_\Phi) = (\widehat\lambda + \hat\rho, \widehat\lambda + \hat\rho) \hat\delta \chi_\Phi.$$

Remark. In fact, both Theorem 6.3.2 and Example 6.3.3 are also true if one replaces the integrable module $L_{\widehat\lambda}$ by Verma module $M_{\widehat\lambda}$ or any its subfactor, and lets $\widehat\lambda$ be an arbitrary (not necessarily dominant) weight. The only difference is that in this case the generalized character should be understood as an element of $\overline{\mathbb{C}[\widehat P]}$ rather then A.

CHAPTER VII

AFFINE JACOBI POLYNOMIALS AND GENERALIZED CHARACTERS

In this chapter we define and study the affine analogue of Jacobi polynomials discussed in Chapter 3. This definitions is due to the author and Etingof [EK4]. (The name "polynomials" is not quite good, since they look rather like theta-functions; still, we use it to stress the analogy with finite-dimensional case).

We also prove that for the root system of type $\hat{A}_{n-1}$ these polynomials, which in this case we call affine Jack polynomials, can be obtained as ratio of generalized characters for the affine Lie algebra $\widehat{\mathfrak{sl}_n}$.

7.1 Definition of affine Jacobi polynomials.

As before, let us fix some positive integer k. Recall that we have defined in the previous chapter the algebra A, which is a certain completion of the group algebra $\mathbb{C}[\widehat{P}]$, and the ring of fractions $\hat{\mathcal{R}}$, which is a certain completion of the ring $\mathbb{C}[\widehat{P}](e^{\hat{\alpha}}-1)^{-1}$. Recall also the Laplace operator $\Delta_{\mathfrak{h}}$ defined in Section 2.2.

Definition 7.1.1. The Calogero-Sutherland operator for (affine) root system $\widehat{R}$ is the differential operator which acts in $\hat{\mathcal{R}}_K$ by the following formula (all the notations as before):

$$\hat{L}_k = \Delta_{\mathfrak{h}} - 2Kp\frac{\partial}{\partial p} - k(k-1)\sum_{\substack{\alpha\in R^+\\ n\in\mathbb{Z}}} \frac{p^n e^{\alpha}}{(1-p^n e^{\alpha})^2}(\alpha,\alpha). \tag{7.1.1}$$

Remarks.

(1) It is easy to see that the sum does belong to the ring $\hat{\mathcal{R}}$, so $\hat{L}_k$ is a well defined operator in $\hat{\mathcal{R}}$.

(2) This operator can be rewritten in the following form:

$$\hat{L}_k = \hat{\Delta} - k(k-1)\sum_{\hat{\alpha}\in\widehat{R}^+} \frac{(\hat{\alpha},\hat{\alpha})}{(e^{\hat{\alpha}/2}-e^{-\hat{\alpha}/2})^2}, \tag{7.1.2}$$

which shows that it is a complete analogue of the operator L_k introduced in Section 3.1. This also shows that $\hat{L}_k$ commutes with the action of $\widehat{W}$ in $\hat{\mathcal{R}}$.

(3) Note that for $K = 0$ this operator becomes so-called elliptic Calogero-Sutherland operator (it becomes clear if one rewrites $\hat{L}_k$ in terms of Weierstrass function – see formula (8.1.6) below).

Define

$$\widehat{M_k} = \hat{\delta}^{-k} \circ (\hat{L}_k - k^2(\hat{\rho}, \hat{\rho})) \circ \hat{\delta}^k, \tag{7.1.3}$$

where $\hat{\delta}$ is the affine Weyl denominator (6.2.4).

THEOREM 7.1.2.

(1) *$\widehat{M_k}$ is a well defined operator in $\hat{\mathcal{R}}$.*

(2)

$$\widehat{M_k} = \hat{\Delta} - 2k \sum_{\hat{\alpha} \in \widehat{R}^+} \frac{1}{1 - e^{\hat{\alpha}}} \partial_{\hat{\alpha}} + 2k\partial_{\hat{\rho}}. \tag{7.1.4}$$

(3) *$\widehat{M_k}$ commutes with the action of $\widehat{W}$.*

PROOF. It will be convenient to use vector fields in $\hat{\mathfrak{h}}$, i.e. the elements of $\hat{\mathcal{R}} \otimes \hat{\mathfrak{h}}$. If $v = \sum v_i \lambda_i, v_i \in \hat{\mathcal{R}}, \lambda_i \in \hat{\mathfrak{h}}$ then we will denote by $\partial_v = \sum v_i \partial_{\lambda_i}$ the corresponding differential operator. Also, for $f \in \hat{\mathcal{R}}$ denote by grad $f \in \hat{\mathcal{R}} \otimes \hat{\mathfrak{h}}$ its gradient, which is defined in the usual way, so that grad $e^{\widehat{\lambda}} = e^{\widehat{\lambda}} \cdot \widehat{\lambda}$ (as before, we identify $\hat{\mathfrak{h}}^*$ and $\hat{\mathfrak{h}}$). Then we have the following obvious formula:

$$\hat{\Delta}(fg) = (\hat{\Delta} f)g + f(\hat{\Delta} g) + 2(\text{grad } f, \text{grad } g) = (\hat{\Delta} f)g + f(\hat{\Delta} g) + 2\partial_{\text{grad } f} g,$$

where (,) is the inner product of the vector fields, i.e., the inner product on $\hat{\mathfrak{h}}$ extended by $\hat{\mathcal{R}}$-linearity to $\hat{\mathcal{R}} \otimes \hat{\mathfrak{h}}$ (it has nothing to do with the inner product on polynomials!). Thus,

$$\hat{\delta}^{-1} \circ \hat{\Delta} \circ \hat{\delta} = \hat{\Delta} + 2\partial_v + (\hat{\delta}^{-1} \hat{\Delta}(\hat{\delta})), \tag{7.1.5}$$

where

$$v = \hat{\delta}^{-1} \text{grad } \hat{\delta} = \hat{\rho} + \sum_{\hat{\alpha} \in \widehat{R}^+} \frac{e^{-\hat{\alpha}}}{1 - e^{-\hat{\alpha}}} \hat{\alpha}. \tag{7.1.6}$$

Note that $\widehat{W}$-antiinvariance of $\hat{\delta}$ implies $\widehat{W}$-invariance of v. Since $\hat{\Delta}(\hat{\delta}) = (\hat{\rho}, \hat{\rho})\hat{\delta}$, which follows from the denominator identity for affine root systems, this proves that $\hat{\delta}^{-1} \hat{\Delta} \hat{\delta}$ is a well defined operator in $\hat{\mathcal{R}}$.

It is easy to prove by induction that

$$\hat{\delta}^{-k} \circ \hat{\Delta} \circ \hat{\delta}^k = \hat{\Delta} + k(\hat{\rho}, \hat{\rho}) + 2k\partial_v + k(k-1)\hat{\delta}^{-1}(\partial_v \hat{\delta}).$$

Obviously, $\hat{\delta}^{-1}(\partial_v \hat{\delta}) = (v, v)$.

LEMMA 7.1.3.

$$(v,v) = (\hat{\rho},\hat{\rho}) + \sum_{\hat{\alpha}\in\widehat{R}^+} (\hat{\alpha},\hat{\alpha})\frac{1}{(e^{\hat{\alpha}/2}-e^{-\hat{\alpha}/2})^2}.$$

PROOF. Let us consider $X = (v,v) - (\hat{\rho},\hat{\rho}) - \sum_{\hat{\alpha}\in\widehat{R}^+}(\hat{\alpha},\hat{\alpha})\frac{1}{(e^{\hat{\alpha}/2}-e^{-\hat{\alpha}/2})^2} \in \hat{\mathcal{R}}$. Obviously, it is $\widehat{W}$-invariant. Also, from the explicit expression for v it follows that X has only simple poles. Consider $X\hat{\delta}$. It is an element of $\hat{\mathcal{R}}$ with no poles, thus it is an element of A. Also, it is antiinvariant. Thus, $X \in A$. Obviously, $X \in A_0$; since X is $\widehat{W}$-invariant, Lemma 6.2.1 implies that $X \in \mathbb{C}((p))$.

To complete the calculation, let us write the explicit expression for X; then, let us expand it in a series in $e^{-\hat{\alpha}}$, $\hat{\alpha} \in \widehat{R}^+$ (i.e., apply the map τ we used in Section 6.2 to define $\hat{\mathcal{R}}$) and keep only the terms of the form $e^{n\delta}$ in the expansion. This gives:

$$X = 2\left(rh^\vee - \sum_{\alpha\in R^+}(\alpha,\alpha)\right)\sum_{n=1}^{\infty}\frac{np^n}{1-p^n},$$

where r is the rank of $\mathfrak{g}$. On the other hand, for any simple Lie algebra $\mathfrak{g}$ we have

$$\sum_{\alpha\in R^+}(\alpha,\alpha) = rh^\vee.$$

The simplest way to prove it is to consider the action of the Casimir element $C \in U\mathfrak{g}$ in the adjoint representation. On one hand, $C|_{\mathfrak{g}} = 2h^\vee \operatorname{Id}_{\mathfrak{g}}$, and thus $\operatorname{Tr}|_{\mathfrak{h}}C = 2rh^\vee$. On the other hand, it is easy to deduce from the formula $C = \sum_{\alpha\in R^+} e_\alpha f_\alpha + f_\alpha e_\alpha + \sum x_l^2$ that $\operatorname{Tr}_{\mathfrak{h}} C = 2\sum_{\alpha\in R^+}(\alpha,\alpha)$. Thus, $X = 0$.

This lemma together with previous results immediately implies statements 1 and 2 of the theorem. Statement 3 follows from $\widehat{W}$-invariance of the operator L_k. □

Note that Theorem 7.1.2 is a complete analogue of Proposition 3.1.1.

REMARKS.

(1) This technique is borrowed from [Ma].
(2) In the simply-laced case the identity $\sum(\alpha,\alpha) = rh^\vee$ becomes $\dim\mathfrak{g} = r(h+1)$, where h is the Coxeter number for $\mathfrak{g}$. This latter identity is well known and has a beautiful interpretation in terms of the Coxeter automorphism (see [Kos]).

THEOREM 7.1.4. *$\widehat{M}$ preserves the algebra of $\widehat{W}$- invariant polynomials: $\widehat{M}A_K^{\widehat{W}} \subset A_K^{\widehat{W}}$. Moreover, in the basis of orbitsums its action is triangular:*

$$\widehat{M}m_{\widehat{\lambda}} = (\widehat{\lambda},\widehat{\lambda}+2k\hat{\rho})m_{\widehat{\lambda}} + \sum_{\substack{\hat{\mu}<\widehat{\lambda}\\ \hat{\mu}\in\hat{P}_K^+}} c_{\widehat{\lambda}\hat{\mu}}m_{\hat{\mu}}. \tag{7.1.7}$$

PROOF. Proof is based on the following lemma:

Lemma. *Let* $f \in A^{\widehat{W}}, f = e^{\widehat{\lambda}} +$ *lower terms*, $\hat{\alpha} \in \widehat{R}^+$. *Then* $\frac{1}{1-e^{\hat{\alpha}}}\partial_{\hat{\alpha}} f$ *is a well-defined element of* A *with highest term* $-(\widehat{\lambda}, \hat{\alpha})e^{\widehat{\lambda}-\hat{\alpha}}$.

Proof of the Lemma. The proof is based on the fact that due to $\widehat{W}$-invariance, f contains terms $e^{\hat{\mu}}$ and $e^{\hat{\mu}-\langle\hat{\mu},\hat{\alpha}^\vee\rangle\hat{\alpha}}$ with equal coefficients, and on explicit calculation.

Now the statement of the theorem follows from the explicit formula (7.1.4) for M_k. $\square$

Our main objective will be the study of the eigenfunctions of action of this operator in $A^{\widehat{W}}$, which we will call affine Jacobi polynomials. More precisely, let us consider the action of $\widehat{M}$ in the linear space spanned by $m_{\hat{\mu}}$ with $\hat{\mu} \le \widehat{\lambda}$. This space is not finite-dimensional; however, one can still check that $\widehat{M}$ has a unique eigenvector with eigenvalue $(\widehat{\lambda}, \widehat{\lambda} + 2k\hat{\rho})$ in this space (this is based on the affine analogue of Lemma 3.1.3, and on some standard arguments on convergence). Thus, we adopt the following definition:

Definition 7.1.5. Affine Jacobi polynomials $\hat{J}_{\widehat{\lambda}}, \widehat{\lambda} \in \widehat{P}^+$ are the elements of $A^{\widehat{W}}$ defined by the following conditions:

1. $\hat{J}_{\widehat{\lambda}} = m_{\widehat{\lambda}} + \sum\limits_{\hat{\mu}<\widehat{\lambda}} c_{\widehat{\lambda}\hat{\mu}} m_{\hat{\mu}}$.
2. $\widehat{M}_k \hat{J}_{\widehat{\lambda}} = (\widehat{\lambda}, \widehat{\lambda} + 2k\hat{\rho})\hat{J}_{\widehat{\lambda}}$.

As was said above, these conditions determine $\hat{J}_{\widehat{\lambda}}$ uniquely. Note that if $\hat{\mu} = \widehat{\lambda} + n\delta$ then $\hat{J}_{\hat{\mu}} = p^{-n}\hat{J}_{\widehat{\lambda}}$. Thus, it suffices to consider only the polynomials $J_{\widehat{\lambda}}$ for $\widehat{\lambda} = \lambda + K\Lambda_0, \lambda \in P^+_K$.

Examples.

1. Let $k = 0$; then $\hat{J}_{\widehat{\lambda}} = m_{\widehat{\lambda}}$.
2. Let $k = 1$; then $\hat{J}_{\widehat{\lambda}} = \operatorname{ch} L_{\widehat{\lambda}}$.

Theorem 7.1.6. *For every* $K \in \mathbb{Z}_+$, *affine Jacobi polynomials* $\{\hat{J}_{\lambda+K\Lambda_0}\}_{\lambda\in P^+_K}$ *form a basis over* $\mathbb{C}((p))$ *in* $A^{\widehat{W}}_K$.

Proof. It follows from the fact that the orbitsums form a basis of $A^{\widehat{W}}_K$ (Theorem 6.2.2) and the following elementary lemma the proof of which is left to the reader:

Lemma 7.1.7. *Fix* $K \in \mathbb{Z}_+$. *Let* $A = \{a_{\widehat{\lambda}\hat{\mu}}\}_{\widehat{\lambda},\hat{\mu}\in\widehat{P}^+_K}$ *be a matrix with complex entries satisfying the following conditions:*

1. $a_{\widehat{\lambda}\hat{\mu}} = 0$ *unless* $\hat{\mu} \le \widehat{\lambda}$.
2. $a_{\widehat{\lambda}\widehat{\lambda}} = 1$.

Then this matrix has an inverse: there exists a unique matrix $B = \{b_{\widehat{\lambda}\hat{\mu}}\}$ *satisfying the same conditions such that* $AB = \delta_{\widehat{\lambda}\hat{\mu}}$.

7.2 Affine Jack polynomials as generalized characters

In this section we consider affine Jacobi polynomials (see Definition 7.1.5) for the root system $\hat{A}_{n-1}$, which we call affine Jack polynomials, and show that they can be obtained from the generalized characters for the Lie algebra $\widehat{\mathfrak{sl}_n}$. In this section, we only consider $\hat{\mathfrak{g}} = \widehat{\mathfrak{sl}_n}$.

As before, we fix a positive integer k. As in Section 3.1, let U be the finite-dimensional irreducible representation of $\mathfrak{sl}_n$ with highest weight $n(k-1)\omega_1$. Recall that $U[0]$ is one dimensional, and we have fixed an element $u_0 \in U[0]$, thus identifying $U[0] \simeq \mathbb{C} : u_0 \mapsto 1$. Let $U(z)$ be the corresponding evaluation representation of $\widehat{\mathfrak{sl}_n}$.

Proposition 7.2.1. *Let* $\mu \in \widehat{P}^+$. *A non-zero intertwiner*

$$\Phi\colon L_{\hat{\mu}} \to \widehat{L_{\hat{\mu}}} \otimes U(z)$$

exists iff $\hat{\mu} = (k-1)\hat{\rho} + \widehat{\lambda}, \widehat{\lambda} \in \widehat{P}^+$; *if it exists, it is unique up to a scalar. We will denote such an intertwiner by* $\Phi_{\widehat{\lambda}}$.

Proof. The proof is based on Lemma 6.1.3.

Let us consider the corresponding generalized characters

$$\varphi_{\widehat{\lambda}} = \chi_{\Phi_{\widehat{\lambda}}}, \quad \widehat{\lambda} \in \widehat{P}^+. \tag{7.2.1}$$

They take values in $U[0]$, which is one-dimensional, and thus can be considered as scalar-valued so that

$$\varphi_{\widehat{\lambda}} = e^{\widehat{\lambda}+(k-1)\hat{\rho}} + \textit{lower order terms.}$$

Proposition 7.2.2. *For every* $\hat{\alpha} \in \widehat{R}^+$, $\varphi_{\widehat{\lambda}}$ *is divisible by* $(1-e^{\hat{\alpha}})^{k-1}$ (*divisibility is to be understood in the algebra* A).

Proof. This statement is trivially true for $\hat{\alpha} = n\delta$, since $(1 - p^{-n})$ is invertible in $\mathbb{C}((p))$. Therefore, we can assume that $\hat{\alpha} = \alpha + n\delta, \alpha \in R$, in which case we can repeat the arguments used in finite-dimensional situation (see proof of Proposition 4.2.2). Let us consider the traces of the form

$$\varphi^F_{\widehat{\lambda}} = \sum_{\hat{\mu} \in \widehat{P}} e^{\hat{\mu}} \operatorname{Tr}|_{L_{\widehat{\lambda}}[\hat{\mu}]}(\Phi_{\widehat{\lambda}} F), \tag{7.2.2}$$

where F is an arbitrary element of $U\widehat{\mathfrak{sl}_n}$. Let us take $F = f_{\hat{\alpha}}^{k-1}$. Then, using the intertwining property of $\Phi_{\widehat{\lambda}}$ and the identity $\Delta(f_{\hat{\alpha}}) = f_{\hat{\alpha}} \otimes 1 + 1 \otimes f_{\hat{\alpha}}$, we can prove by induction that

$$\varphi_{\widehat{\lambda}}^F = \frac{f_{\hat{\alpha}}^{k-1} \varphi_{\widehat{\lambda}}}{(1 - e^{-\hat{\alpha}})^{k-1}}.$$

Since $U[0], U[-(k-1)\alpha]$ are one-dimensional, we can identify both of them with $\mathbb{C}$; then $f_{\hat{\alpha}}^{k-1} \colon U[0] \to U[-(k-1)\alpha]$ becomes a non-zero constant. On the other hand, it is easy to see that $\varphi_{\widehat{\lambda}}^F \in A$, which proves the proposition. □

THEOREM 7.2.3. *Let* $\varphi_{\widehat{\lambda}}$ *be the generalized character for* $\widehat{\mathfrak{sl}_n}$*, defined by* (7.2.1)*. Then*

$$\varphi_0 = \hat{\delta}^{k-1}, \tag{7.2.3}$$

where $\hat{\delta}$ *is the affine Weyl denominator* (6.2.4)*.*

PROOF. Let us consider the ratio $f = \varphi_0 / \hat{\delta}^{k-1}$. It follows from Proposition 7.2.2 that $f \in A$. It has level zero and highest term 1. Moreover, similar arguments show that if we twist the order on $\widehat{P}$ by the action of the Weyl group: $\widehat{\lambda} \geq_w \hat{\mu}$ if $\widehat{\lambda} - \hat{\mu} \in w(\hat{Q}^+), w \in \widehat{W}$ then the highest term of f with respect to any such twisted ordering is still 1. This is only possible if $f \in \mathbb{C}((p))$. To complete the proof, we have to use the differential equation for the characters. Indeed, Example 6.3.3 implies that $\varphi_{\widehat{\lambda}}$ satisfies the following equation:

$$\hat{L}(\varphi_{\widehat{\lambda}} \hat{\delta}) = (\widehat{\lambda} + k\hat{\rho}, \widehat{\lambda} + k\hat{\rho})(\varphi_{\widehat{\lambda}} \hat{\delta}),$$

since $e_\alpha f_\alpha|_{U[0]} = k(k-1)\,\mathrm{Id}_{U[0]}$.

Substituting in this equation $\varphi_0 = f(p)\hat{\delta}^{k-1}$, we see that f satisfies $\widehat{M} f = 0$. Using formula (7.1.4) for $\widehat{M}$ we get $2kh^\vee p \frac{\partial}{\partial p} f = 0$, which is possible only if f is a constant. Comparing the highest terms of φ_0 and $\hat{\delta}^{k-1}$, we get the statement of the theorem. □

Now we can prove the main theorem of this section:

THEOREM 7.2.4. *Let* $\varphi_{\widehat{\lambda}}, \widehat{\lambda} \in \widehat{P}^+$ *be the generalized character for* $\widehat{\mathfrak{sl}_n}$ *defined by formula* (7.2.1)*, and let* $\hat{J}_{\widehat{\lambda}}$ *be Jack* (*Jacobi*) *polynomial for the root system* $\hat{A}_{n-1}$*. Then*

$$\frac{\varphi_{\widehat{\lambda}}}{\varphi_0} = \hat{J}_{\widehat{\lambda}}.$$

PROOF. The proof is again quite similar to the finite-dimensional case (Theorem 3.2.3). First, we prove that $\varphi_{\widehat{\lambda}} / \varphi_0 \in A^{\widehat{W}}$. Consider the module $L =$

$L_{\widehat{\lambda}} \otimes L_{(k-1)\hat{\rho}}$. This module is unitary (since both factors are unitary); thus, it is completely reducible and can be decomposed in a direct sum of the modules $L_{\hat{\mu}}$:

$$L = L_{(k-1)\hat{\rho}+\widehat{\lambda}} + \sum_{\substack{\hat{\mu}\in \widehat{P}^+ \\ \hat{\mu}<\widehat{\lambda}+(k-1)\hat{\rho}}} N_{\hat{\mu}} L_{\hat{\mu}}.$$

This sum is, of course, infinite; however, all multiplicities $N_{\hat{\mu}}$ are finite. In particular, this implies that the character of this module belongs to the algebra $A^{\widehat{W}}$ (that is, $\widehat{W}$-invariants of completed group algebra of $\widehat{P}$), so in a certain sense this sum converges.

Let us construct an intertwiner $\Psi\colon L \to L \otimes U(z)$ as $\Psi = \mathrm{Id}_{L_{\widehat{\lambda}}} \otimes \Phi_0$. Consider the corresponding generalized character χ_Ψ. Then it follows from the decomposition of L that

$$\chi_\Psi = \varphi_{\widehat{\lambda}} + \sum_{\substack{\hat{\mu}\in \widehat{P}^+ \\ \hat{\mu}<\widehat{\lambda}}} a_{\widehat{\lambda}\hat{\mu}} \varphi_{\hat{\mu}}.$$

On the other hand, $\chi_\Psi = \varphi_0 \operatorname{ch} L_{\widehat{\lambda}}$. Dividing both sides by φ_0 we get

$$\operatorname{ch} L_{\widehat{\lambda}} = \sum_{\hat{\mu}\in \widehat{P}^+} a_{\widehat{\lambda}\hat{\mu}} \frac{\varphi_{\hat{\mu}}}{\varphi_0},$$

where $a_{\widehat{\lambda}\hat{\mu}} \neq 0$ only if $\hat{\mu} \leq \widehat{\lambda}$, and $a_{\widehat{\lambda}\widehat{\lambda}} = 1$. It follows from Lemma 7.1.7 that we can invert this matrix, writing

$$\frac{\varphi_{\widehat{\lambda}}}{\varphi_0} = \sum_{\hat{\mu}\in \widehat{P}^+} b_{\widehat{\lambda}\hat{\mu}} \operatorname{ch} L_{\hat{\mu}},$$

and the coefficients $b_{\widehat{\lambda}\hat{\mu}}$ satisfy the same conditions as $a_{\widehat{\lambda}\hat{\mu}}$. Thus, $\varphi_{\widehat{\lambda}}/\varphi_0 \in A^{\widehat{W}}$ and has highest term $e^{\widehat{\lambda}}$.

We have proved that $\varphi_{\widehat{\lambda}}/\varphi_0$ satisfies the first condition in the definition of Jack polynomials. Now, using the differential equation for generalized characters (Example 6.3.3) along with the fact that $e_\alpha f_\alpha|_{U[0]} = k(k-1)\,\mathrm{Id}_{U[0]}$, we get

$$\hat{L}(\varphi_{\widehat{\lambda}}\hat{\delta}) = (\widehat{\lambda} + k\hat{\rho}, \widehat{\lambda} + k\hat{\rho})(\varphi_{\widehat{\lambda}}\hat{\delta}).$$

Since $\varphi_0 = \hat{\delta}^{k-1}$ we can rewrite this as follows:

$$\hat{L}\left(\frac{\varphi_{\widehat{\lambda}}}{\varphi_0}\hat{\delta}^k\right) = (\widehat{\lambda} + k\hat{\rho}, \widehat{\lambda} + k\hat{\rho})\left(\frac{\varphi_{\widehat{\lambda}}}{\varphi_0}\hat{\delta}^k\right),$$

which is precisely the definition of affine Jack's polynomials. $\square$

CHAPTER VIII

MODULAR PROPERTIES OF AFFINE JACOBI POLYNOMIALS

In this chapter we consider the affine Jacobi polynomials defined in the previous chapter from the analytical point of view. We prove that they define an analytic function in a certain domain, show their connection with theta-functions and study their modular properties, following the paper [EK4].

All the constructions of this section are valid for arbitrary Lie algebra $\mathfrak{g}$.

8.1 Functional interpretation of $\mathbb{C}[\widehat{P}]$.

So far, we have considered elements of $\mathbb{C}[\widehat{P}]$ and its various completions and extensions formally. In this section we discuss the analytic approach.

Let us define the following domain

$$Y = \mathfrak{h} \times \mathbb{C} \times \mathcal{H}, \tag{8.1.1}$$

where $\mathcal{H}$ is the upper half-plane: $\mathcal{H} = \{\tau \in \mathbb{C} | \operatorname{Im} \tau > 0\}$. Then every element $e^{\widehat{\lambda}} \in \mathbb{C}[\widehat{P}]$ can be considered as a function on Y as follows: if $\widehat{\lambda} = \lambda + a\delta + K\Lambda_0$ then put

$$e^{\widehat{\lambda}}(h, u, \tau) = e^{2\pi \mathrm{i}[\langle \lambda, h\rangle + Ku - a\tau]}. \tag{8.1.2}$$

Note that this in particular implies that $p = e^{-\delta}$ is given by

$$p = e^{2\pi \mathrm{i}\tau}. \tag{8.1.3}$$

It is easy to see that if $f \in \mathbb{C}[\widehat{P}]$ then $f(h, u, \tau + 1) = f(h, u, \tau)$, so we can as well consider f as a function of h, u, p, where $p \in \mathbb{C}$ is such that $0 < |p| < 1$ (the reason for this choice of possible values for p will be clear later). Note that $f \in \mathbb{C}[\widehat{P}_K] \iff f(h, u, \tau) = e^{2\pi \mathrm{i} Ku} f(h, 0, \tau)$; in this case we say that f is a function of level K.

Of course, we can't extend this rule to the completion A. However, it turns out that we can extend it to certain elements of A.

DEFINITION 8.1.1. Let $f = \sum a_{\widehat{\lambda}} e^{\widehat{\lambda}} \in \overline{\mathbb{C}[\widehat{P}]}$, and let $U \subset Y$. We say that f is an analytic function on U (or that f converges on U) if this sum, considered as a sum of functions on Y by the above rule absolutely converges in U, and this convergence is uniform on compact subsets in U. Similarly, if $f \in \hat{\mathcal{R}}$ then we say that f is an analytic function on U if $\tau(f)$ converges on U, where $\tau : \hat{\mathcal{R}} \to \overline{\mathbb{C}[\widehat{P}]}$ is the homomorphism obtained by expanding each $(1-e^{\hat{\alpha}})^{-1} = 1 + e^{\hat{\alpha}} + e^{2\hat{\alpha}} + \dots$, $\hat{\alpha} \in \widehat{R}^-$ (see the definition of $\hat{\mathcal{R}}$ in Section 6.2).

The following theorem is well known (see, for example, [Ka1, Chapters 11,13])

THEOREM 8.1.2.

(1) *For every $\widehat{\lambda} \in \widehat{P}^+$, the orbitsum $m_{\widehat{\lambda}}$ defined by* (6.2.3) *is an analytic function on Y.*

(2) *The affine Weyl denominator $\hat{\delta}$ defined by* (6.2.4) *is an analytic function in Y, and*

$$\hat{\delta}(h,u,\tau) = e^{2\pi \mathrm{i} h^\vee u} \mathrm{i}^{|R^+|} p^{-\frac{\dim \mathfrak{g}}{24}} \eta(p)^{r-|R^+|} \prod_{\alpha \in R^+} \theta_1(\langle \alpha, h\rangle), \tag{8.1.4}$$

where the theta (θ_1) and eta (η) functions are defined in the Appendix.

It is easy to rewrite the definition of $\widehat{W}$-invariance in analytic terms. Indeed, define the action of the affine Weyl group $\widehat{W}$ on Y so that $e^{w\widehat{\lambda}}(h,u,\tau) = e^{\widehat{\lambda}}(w^{-1}(h,u,\tau))$. One easily checks that when restricted to the finite Weyl group W this action coincides with the usual action of W on $\mathfrak{h}$ (leaving u, τ invariant), and the action of $\alpha^\vee \in Q^\vee$ is given by

$$\alpha^\vee(h,u,\tau) = (h - \alpha^\vee \tau, u + \frac{1}{2}(\alpha^\vee, \alpha^\vee)\tau - \langle \alpha^\vee, h\rangle, \tau).$$

This immediately implies the following proposition:

THEOREM 8.1.3. *Let $f \in A_K$ be such that it converges in Y. Define $\tilde{f}(h,\tau) = e^{-2\pi \mathrm{i} K u} f(h,u,\tau)$. Let us consider the ring $A^{Q^\vee}$ of elements from A invariant with respect to the action of $Q^\vee \subset \widehat{W}$. Then $f \in A^{Q^\vee}$ iff $\tilde{f}$ satisfies the following conditions: for every $\alpha^\vee \in Q^\vee$,*

$$\begin{aligned} \tilde{f}(h + \alpha^\vee, \tau) &= \tilde{f}(h,\tau), \\ \tilde{f}(h + \alpha^\vee \tau, \tau) &= e^{-2\pi \mathrm{i} K (\frac{1}{2}(\alpha^\vee, \alpha^\vee)\tau + \langle \alpha^\vee, h\rangle)} \tilde{f}(h,\tau), \end{aligned} \tag{8.1.5}$$

and $f \in A^{\widehat{W}}$ iff $\tilde{f}$ satisfies (8.1.5) *and is W-invariant.*

Functions on $\mathfrak{h} \times \mathcal{H}$ satisfying (8.1.5) are usually called theta-functions of level K; thus we see that $A^{\widehat{W}}$ is the formal analogue of the ring of W-symmetric theta-functions (cf. [Lo]).

Finally, let us consider the questions of convergence of affine Jacobi polynomials and generalized characters. Recall (see Section 2.2) that we have introduced the differential operators $\partial_x, x \in \mathfrak{h}$ and $\Delta_{\mathfrak{h}}$ acting in functions on $\mathfrak{h}$ so that

$$\begin{aligned}\partial_x e^{2\pi\mathrm{i}\langle\alpha,h\rangle} &= \langle\alpha,x\rangle e^{2\pi\mathrm{i}\langle\alpha,h\rangle},\\ \Delta_{\mathfrak{h}} e^{2\pi\mathrm{i}\langle\alpha,h\rangle} &= (\alpha,\alpha) e^{2\pi\mathrm{i}\langle\alpha,h\rangle}.\end{aligned}$$

If we choose an orthonormal basis $x_l, l = 1,\dots,r$ in $\mathfrak{h}$ and denote by c_l the corresponding coordinates in $\mathfrak{h}$: $h = \sum c_l x_l, c_l \in \mathbb{C}$ then these operators can be rewritten as follows:

$$\begin{aligned}\partial_{x_l} &= \frac{1}{2\pi\mathrm{i}}\frac{\partial}{\partial c_l}\\ \Delta_{\mathfrak{h}} &= -\frac{1}{4\pi^2}\sum\frac{\partial^2}{\partial c_l^2}.\end{aligned} \tag{8.1.6}$$

Now we can write the differential operators $\hat{L}, \widehat{M}$ defined in Section 7.1 as usual differential operators in Y with analytic coefficients:

Proposition 8.1.4. *The coefficients of the differential operators* $\hat{L}, \widehat{M}$ *defined by* (7.1.1), (7.1.5) *respectively are analytic functions in the region* $\{0 < \operatorname{Im}\langle\alpha,h\rangle < \operatorname{Im}\tau$ *for all* $\alpha \in R^+\}$, *and in this region the following formulas hold: if we consider them as operators acting on functions of level* K *then*

$$\begin{aligned}\hat{L}_k &= \Delta_{\mathfrak{h}} - 2Kp\frac{\partial}{\partial p} + \frac{k(k-1)}{4\pi^2}\sum_{\alpha\in R^+}(\wp(\langle\alpha,h\rangle) + c(\tau)),\\ \widehat{M} &= \Delta_{\mathfrak{h}} - 2(K + kh^\vee)p\frac{\partial}{\partial p} - 2k\sum_{\alpha\in R^+}\sigma(\langle\alpha,h\rangle)\partial_\alpha\\ &\quad -2kK\sum_{n\geq1}\left(r\frac{1}{1-p^{-n}}n + \sum_{\alpha\in R}\frac{1}{1-p^{-n}e^\alpha}n\right),\end{aligned} \tag{8.1.7}$$

where the functions $\theta_1(x), \wp(x), \sigma(x)$ *are the usual theta-function, Weierstrass p-function and sigma-function, respectively* (*see Appendix*), *and the constant* $c(\tau)$ *is defined in the Appendix* (*formula* A4).

Proof. Explicit calculation.

Theorem 8.1.5. *For every* $\widehat{\lambda} \in \widehat{P}^+$, *the affine Jacobi polynomial* $\hat{J}_{\widehat{\lambda}}$ *converges in* Y.

Proof. Assume $\widehat{\lambda} \in \widehat{P}_K$. Due to Theorem 6.2.2, we can write $\hat{J}_{\widehat{\lambda}}$ as linear combination of orbitsums: $\hat{J}_{\widehat{\lambda}} = \sum_{\lambda\in P_K^+} f_\lambda(p) m_{\lambda+K\Lambda_0}$ for some $f_\lambda \in \mathbb{C}((p))$. Let us substitute this in the defining equation

$$\widehat{M}\hat{J}_{\widehat{\lambda}} = (\widehat{\lambda}, \widehat{\lambda} + 2k\hat{\rho})\hat{J}_{\widehat{\lambda}}.$$

Since $\hat{\Delta}(m_{\widehat{\lambda}}) = (\widehat{\lambda}, \widehat{\lambda})m_{\widehat{\lambda}}$, it is easy to deduce that $f_\lambda(p)$ satisfy the following system of first-order differential equations:

$$-2(K + kh^\vee)p\frac{\partial}{\partial p}f_\lambda = \sum_{\mu \in P_K^+} f_\mu a_{\mu\lambda},$$

where the coefficients $a_{\lambda\mu} \in \mathbb{C}((p))$ are defined by

$$\widehat{M}m_{\lambda+K\Lambda_0} = \sum_{\mu \in P_K^+} a_{\lambda\mu}m_{\mu+K\Lambda_0}.$$

It follows from Proposition 8.1.4 that $\widehat{M}m_{\lambda+K\Lambda_0}$ converges in the region $0 < \mathrm{Im}\,\langle\alpha, h\rangle < \mathrm{Im}\,\tau$; this is only possible if each $a_{\lambda\mu}$ converges in $0 < |p| < 1$. It is well known that this implies that $f_\lambda(p)$ absolutely converges in the same region. Since $m_{\widehat{\lambda}}$ converges in Y (Theorem 8.1.2), we get the statement of the theorem. $\square$

COROLLARY 8.1.6. *For $\mathfrak{g} = \mathfrak{sl}_n$, the generalized characters $\varphi_{\widehat{\lambda}}$ defined by* (6.2.1)*, converge in Y.*

In fact, it is a general result, which we prove later (see Chapter 9): if L is an integrable module over $\hat{\mathfrak{g}}$, U – finite-dimensional representation of $\mathfrak{g}$, and $\Phi : L \to L \otimes U(z)$ is the $\tilde{\mathfrak{g}}$-intertwiner then χ_Φ converges in Y.

8.2 Normalized characters and modular invariance

In this section we study modular properties of affine Jacobi polynomials, considered – in accordance with the previous section – as functions on $\mathfrak{h} \times \mathbb{C} \times \mathcal{H}$ (see 8.1.1). In this section we fix level $K \in \mathbb{Z}_+$ and assume that $\varkappa = K + kh^\vee \neq 0$, thus excluding the trivial case $K = k = 0$. Then $f \in A_K^{\widehat{W}}$ can be written as $f(h, u, \tau) = e^{2\pi \mathrm{i} Ku}f(h, 0, \tau)$. For this reason, we can consider f as function of only h, τ without losing information.

To make our functions modular invariant we need to introduce some factors of the form $p^t, t \in \mathbb{Q}$. Thus, we need to consider slightly more general setting than in the previous sections. Namely, instead of the weight lattice $\widehat{P}$ we consider a bigger abelian group $\widehat{P}' = P \oplus \mathbb{C}\delta \oplus \mathbb{Z}\Lambda_0$. Also, we can consider the algebra A' formed by finite sums of the form $\sum p^{a_i}f_i, a_i \in \mathbb{C}, f_i \in A$, and the subalgebra of $\widehat{W}$-invariants in A' in a manner quite similar to that of the previous section. All the results of Chapter 6 hold with obvious changes. Again, we can consider elements of A' as functions of h, τ, repeating all the arguments of Section 8.1; the only difference is that elements of A' define multivalued functions of p; of course, this ambiguity vanishes if we consider them as functions of τ.

Define the normalized analogues of $\hat\rho$ and $\hat\delta$ as follows:

$$\begin{aligned}\hat\rho' &= \hat\rho - \frac{(\rho,\rho)}{2h^\vee}\delta = \rho - \frac{(\rho,\rho)}{2h^\vee}\delta + h^\vee\Lambda_0,\\ \hat\delta' &= e^{\hat\rho'}\prod_{\hat\alpha\in\widehat{R}^+}(1-e^{-\hat\alpha}).\end{aligned} \tag{8.2.1}$$

This renormalization is chosen so that $\hat\Delta\hat\delta' = 0$; another reason for this renormalization is that so defined $\hat\delta'$ possesses nice modular properties. Indeed, if we consider $\hat\delta'$ as a function then (8.1.4) implies

$$\hat\delta'(h,u,\tau) = e^{2\pi \mathrm{i} h^\vee u}\mathrm{i}^{|R^+|}\eta(p)^{r-|R^+|}\prod_{\alpha\in R^+}\theta_1(\langle\alpha,h\rangle), \tag{8.2.2}$$

since it is known that

$$\frac{(\rho,\rho)}{2h^\vee} = \frac{\dim \mathfrak{g}}{24}$$

("strange formula" of Freudenthal–de Vries, see [Ka1, Chapter 12]).

Now, let us define the renormalized operator

$$\widehat{M}' = \hat\delta'^{-k}\hat L\hat\delta'^{k} = \widehat{M} - \frac{Kk(\rho,\rho)}{h^\vee}. \tag{8.2.3}$$

It is easy to see that the definition of affine Jacobi polynomials $\hat J_{\widehat\lambda}$ which was given for $\widehat\lambda \in \widehat{P}^+$ can be easily extended to $\widehat\lambda = \lambda + K\Lambda_0 + a\delta \in \widehat{P}', K\in\mathbb{Z}_+, \lambda\in P^+_K, a$– arbitrary. In particular, the following choice of a is of special interest for us:

Definition 8.2.1. Let $K\in\mathbb{Z}_+, \lambda\in P^+_K$. Define the normalized affine Jacobi polynomial $J_{\lambda,K}$ by

$$J_{\lambda,K} = \hat J_{\lambda+\alpha\delta+K\Lambda_0},$$

where

$$a = \frac{k(\rho,\rho)}{2h^\vee} - \frac{(\lambda+k\rho,\lambda+k\rho)}{2(K+kh^\vee)}. \tag{8.2.4}$$

Note that for $k=1$ they are precisely the (usual) characters of integrable highest-weight modules, and the normalization coincides with that in [Ka1, Chapter 13].

Note that in the case $\mathfrak{g} = \mathfrak{sl}_n$ it follows from Theorem 7.2.4 that these polynomials can be defined as follows:

$$J_{\lambda,K}(h,\tau) = \frac{1}{\hat{\delta}'^{k-1}} \operatorname{Tr}_{L_{\lambda^k,K'}} \left(\Phi p^{-d-\frac{c}{24}} e^{2\pi i h} \right), \tag{8.2.5}$$

where, as before, $\lambda^k = \lambda + (k-1)\rho, K' = K + (k-1)h^\vee$, $L_{\lambda^k,K'} = L_{\lambda^k + K'\Lambda_0 + a\delta}$, where

$$a = -\frac{(\lambda^k, \lambda^k + 2\rho)}{2(K' + h^\vee)},$$

and c is the central charge for the action of Virasoro algebra in $L_{\lambda^k,K'}$: $c = \frac{K' \dim \mathfrak{g}}{K'+h^\vee}$ (see [Ka1]). This form is quite usual in the conformal field theory (WZW model on the torus).

It follows from the definition of affine Jacobi polynomials (see Definition 7.1.5) that the normalized affine Jacobi polynomials satisfy the following differential equation:

$$\widehat{M}' J_{\lambda,K} = 0. \tag{8.2.6}$$

Theorem 8.2.2. *The space of solutions of the equation $\widehat{M}' f = 0$ in $A'^{\widehat{W}}_K$ is finite-dimensional, and the basis over $\mathbb{C}$ in the space of solutions is given by the normalized affine Jacobi polynomials $J_{\lambda,K}$.*

Proof. Let $f \in A'^{\widehat{W}}_K$ be such that $\widehat{M}' f = 0$. Due to Theorem 7.1.6, we can write $f = \sum_{\lambda \in P^+_K} f_\lambda(p) J_{\lambda,K}$. Substituting it in $\widehat{M}' f = 0$, we get $p\frac{\partial}{\partial p} f_\lambda = 0$, which proves the theorem. □

It follows from Theorem 8.1.5 that $J_{\lambda,K}$ converges on $\mathfrak{h} \times \mathcal{H}$. Thus, we can formulate the analytic version of the theorem above:

Theorem 8.2.3. *For any $K \in \mathbb{Z}_+$, the normalized affine Jacobi polynomials $\{J_{\lambda,K}\}_{\lambda \in P^+_K}$ form a basis over $\mathbb{C}$ of solutions of the equation $\widehat{M}' f = 0$ in the space of W-symmetric theta-functions of level K (see conditions* (8.1.5)*).*

Now let us recall some facts about the modular group and its action. Recall that the modular group $\Gamma = SL_2(\mathbb{Z})$ is generated by the elements

$$S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

satisfying the defining relations $(ST)^3 = S^2, S^2 T = T S^2, S^4 = 1$. This group acts in a natural way on Y as follows:

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} (h, u, \tau) = \left(\frac{h}{c\tau + d}, u - \frac{c(h,h)}{2(c\tau+d)}, \frac{a\tau + b}{c\tau + d} \right). \tag{8.2.7}$$

In particular,

$$\begin{aligned} T(h,u,\tau) &= (h,u,\tau+1),\\ S(h,u,\tau) &= \left(\frac{h}{\tau}, u-\frac{(h,h)}{2\tau}, -\frac{1}{\tau}\right). \end{aligned}$$

Also, for any $j \in \mathbb{C}$ we will define a right action of Γ on functions on Y as follows: if $\alpha = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ then let

$$(f[\alpha]_j)(h,u,\tau) = (c\tau+d)^{-j} f(\alpha(h,u,\tau)).$$

In fact, this is a projective action, which is related to the ambiguity in the choice of $(c\tau+d)^{-j}$ for non-integer j; to make it a true action one must consider a central extension of $SL_2(\mathbb{Z})$; we are not going into details here, only mentioning that the corresponding cocycle takes values in the unit circle in $\mathbb{C}$. We will call this action "an action of weight j". This action obviously commutes with the action of W. Moreover, it is well known (and is easy to check) that for every $K \in \mathbb{Z}_+$, this action preserves the space of theta-functions of level K.

Our main goal will be to find the behavior of the (normalized) affine Jacobi polynomials under modular transformations. The first result in this direction is

THEOREM 8.2.4. *Fix $\varkappa \in \mathbb{N}$. Then the space of all solutions of the equation $\hat{L}f = 0$ in the space of functions of level $\varkappa$ on Y (i.e., functions f satisfying $f(h,u,\tau) = e^{2\pi i \varkappa u} f(h,0,\tau)$) is invariant under the action of Γ of weight $j = \frac{r}{2}\left(1+\frac{k(k-1)h^\vee}{\varkappa}\right)$.*

PROOF. It is easy to see that the operator $\hat{L}$ is invariant under the action of $T \in \Gamma$. Thus, to prove the theorem, it suffices to prove the following formula:

$$\hat{L}(f[S]_j) = \tau^{-2}((\hat{L}g)[S]_j) - \frac{1}{2\pi i \tau}(\varkappa(r-2j) + k(k-1)rh^\vee) f[S]_j.$$

This is based on formula (8.1.7) for $\hat{L}$, which we rewrite in the following form:

$$\hat{L}_k = \Delta_{\mathfrak{h}} - 2\varkappa p \frac{\partial}{\partial p} - k(k-1)\sum_{\alpha\in R^+} (\alpha,\alpha)\varphi(\langle\alpha,h\rangle,\tau),$$

where

$$\begin{aligned} \varphi(x,\tau) = \sum_{n\in\mathbb{Z}} \frac{p^n e^{2\pi i x}}{(1-p^n e^{2\pi i x})^2} &= \frac{1}{4\pi^2}\partial_x^2 \log\theta_1(x)\\ &= -\frac{1}{4\pi^2}\wp(x) + c(\tau) \text{ for some constant } c(\tau). \end{aligned}$$

(see formula (A4) in the Appendix).

Using modular properties of the theta-function (see, for example, [Mu]), we can show that $\varphi(\frac{x}{\tau}, -\frac{1}{\tau}) = \tau^2\varphi(x,\tau) + \frac{\mathrm{i}}{2\pi}\tau$.

Also, it is not too difficult to check that

$$\hat{\Delta}\,(f[S]_j) = \tau^{-2}((\hat{\Delta}f)[S]_j) - \frac{\varkappa}{2\pi\mathrm{i}\tau}(r-2j)(f[S]_j),$$

using the expression for $\hat{\Delta}$ given in Section 8.1

Since $\sum_{\alpha\in R^+}(\alpha,\alpha) = rh^\vee$ (see proof of Lemma 7.1.3), we get the desired formula. □

Corollary 8.2.5. *The space of solutions of the equation $\widehat{M}'f = 0$, where f is a theta-function of level K, is invariant under the action of modular group with the weight $j = -\frac{K(k-1)r}{2(K+kh^\vee)}$.*

Proof. First recall that the space of W-invariant theta-functions of level K is preserved by this action. Now the statement of the theorem follows from the previous theorem and the following well known fact (see [Ka1, Chapter 13]):

$$\hat{\delta}'[\alpha]_{r/2} = l(\alpha)\hat{\delta}',$$

for some function (not a character) $l : \Gamma \to \mathbb{C}^\times$ such that $l^{24} = 1$.

Since we know that the basis of W-invariant solutions is given by the normalized affine Jacobi polynomials, we can rewrite Corollary 8.2.5 in the following form:

Theorem 8.2.6. *Let $K \in \mathbb{Z}_+$. Denote*

$$V_K = \bigoplus_{\lambda\in P_K^+} \mathbb{C}J_{\lambda,K}. \tag{8.2.8}$$

We consider elements of V_K as functions on Y. Then V_K is preserved by the action of Γ with weight $j = -\frac{K(k-1)r}{2(K+kh^\vee)}$. In particular, this means that V_K is naturally endowed with a structure of a projective representation of Γ, and the corresponding cocycle takes values in the unit circle $S^1 = \{z \in \mathbb{C} \mid |z| = 1\}$.

CHAPTER IX

CORRELATION FUNCTIONS ON THE TORUS AND ELLIPTIC KZ EQUATIONS

In this chapter we consider generalized characters for affine Lie algebras with values in a tensor product of evaluation representations. These characters have appeared in the (twisted) Wess-Zumino-Witten (WZW) model of conformal field theory as correlation functions on the torus, and the differential equation for the generalized characters deduced in Chapter 6 can be interpreted as describing their dependence on the modular parameter τ of the torus. It turns out that their dependence on the parameters z_i of evaluation representations can also be described by a simple differential equation, which was first derived by Bernard ([Be]). We derive rigorously these equations and study their monodromies, which has not been done before. We call them elliptic Knizhnik-Zamolodchikov (KZ) equations, the reason being that in the limit $\tau \to i\infty$ these generalized characters become the usual correlation functions on the sphere, i.e. certain matrix coefficients of product of intertwining operators, and the equations reduce to well known Knizhnik-Zamolodchikov equations (see [KZ, TK, FR]).

9.1 Intertwiners and currents

In this section we briefly review the facts about intertwining operators, commutation relations and (usual) KZ equation we are going to use. We refer the reader to [TK, FR] for the proofs. We keep all the notations of Chapters 6,7. Recall that we have defined for every $\widehat{\lambda} \in \hat{\mathfrak{h}}^*$ Verma module $M_{\widehat{\lambda}}$ and irreducible highest-weight module $L_{\widehat{\lambda}}$. In this chapter we always assume that we have fixed level K so that we only consider weights of the form $\widehat{\lambda} = \lambda + a\delta + K\Lambda_0$. Moreover, we will always assume that λ, K satisfy the following condition:

$$\lambda \in P^+, \qquad K \notin \mathbb{Q}. \tag{9.1.1}$$

This condition ensures that $L_{\widehat{\lambda}}$ is equal to the module over $\hat{\mathfrak{g}}$ obtained by induction from the module L_λ over $\mathfrak{g}$. Note that this condition implies that $K + h^\vee \neq 0$. We discuss possible generalizations later.

It is known that the structure of the module $L_{\lambda+a\delta+K\Lambda_0}$ does not depend on the choice of a. It will be convenient for us to fix a as follows: we let $a = -\Delta(\lambda)$, where

$$\Delta(\lambda) = \frac{(\lambda, \lambda + 2\rho)}{2(K + h^\vee)}. \tag{9.1.2}$$

We denote the corresponding irreducible module by $L_{\lambda,K}$ and Verma module by $M_{\lambda,K}$. The reason for such a choice is that in this case the action of d can be written in terms of action of $\tilde{\mathfrak{g}}$ as follows (see [Ka1]):

(9.1.3)
$$d = -\frac{1}{2(K + h^\vee)} \sum_{n\in\mathbb{Z}} \left(\sum_{\alpha\in R^+} (: e_\alpha[n] f_\alpha[-n] : + : f_\alpha[n] e_\alpha[-n] :) + \sum : x_l[n] x_l[-n] : \right),$$

(compare with (6.3.3)), where normal ordered product is defined by (6.3.4).

Also, we have defined evaluation representations $V(z)$. It will be convenient to change this definition, considering z as a formal variable rather than a complex number, as follows:

Definition 9.1.1. Let V be a module over $\mathfrak{g}$. The evaluation module $V(z)$ is the representation of $\hat{\mathfrak{g}}$ in the space $V \otimes z^{-\Delta}\mathbb{C}[z, z^{-1}]$, where Δ is an arbitrary complex number, with the action defined as follows:

$$\begin{gathered} \pi_{V(z)}(x[n]) = z^n \pi_V(x), \\ \pi_{V(z)}(c) = 0, \pi_{V(z)}(d) = z\frac{d}{dz}. \end{gathered} \tag{9.1.4}$$

Note that so defined evaluation representations are indeed modules over the full affine Lie algebra $\hat{\mathfrak{g}}$, not only $\tilde{\mathfrak{g}}$, and they have weight decomposition: $V(z) = \bigoplus_{\widehat{\lambda}\in\hat{\mathfrak{h}}^*} V(z)[\widehat{\lambda}]$ if V has a weight decomposition as $\mathfrak{g}$-module.

We will be interested in the intertwining operators of the form

$$\Phi : L_{\lambda,K} \to L_{\nu,K} \hat{\otimes} L_\mu(z). \tag{9.1.5}$$

where the completed tensor product is defined as follows: if V, W are $\hat{\mathfrak{g}}$-modules then

(9.1.6)
$$(V \hat{\otimes} W)[\widehat{\lambda}] = \left\{ \sum a_i \otimes b_i, a_i \in V[\widehat{\lambda}_i], b_i \in W[\widehat{\lambda}'_i] \,\middle|\, \widehat{\lambda}_i + \widehat{\lambda}'_i = \widehat{\lambda}, (\widehat{\lambda}_i, \hat{\rho}) \to -\infty \right\}.$$

Lemma 9.1.2.

(1) *Let* Φ *be a* $\tilde{\mathfrak{g}}$ *intertwining operator of the form* (9.1.5). *Then it can be uniquely written in the form*

$$\Phi = \sum_{n\in\mathbb{Z}} \Phi[n] z^{-n-\Delta}, \tag{9.1.7}$$

where Δ *is the complex number used in the definition of evaluation representation, and* $\Phi[n]$ *are* $\mathfrak{g}$*-intertwiners* $L_{\lambda,K} \to L_{\nu,K} \otimes L_\mu$ *such that*

$$\Phi[0](L_\lambda) \subset L_\nu \otimes L_\mu,$$

where L_λ *is the* $\mathfrak{g}$*-submodule in* $L_{\lambda,K}$ *generated by the highest-weight vector, and similarly for* ν*.*

(2) *The map* $\Phi \mapsto \Phi[0]$ *establishes isomorphism between the space of* $\tilde{\mathfrak{g}}$ (*not* $\hat{\mathfrak{g}}$!) *intertwiners of the form* (9.1.5) *and the space of* $\mathfrak{g}$*-intertwiners* $L_\lambda \to L_\nu \otimes L_\mu$*.*

(3) *A* $\tilde{\mathfrak{g}}$*-intertwiner* Φ *defined above is a* $\hat{\mathfrak{g}}$ *intertwiner iff* $\Delta = \Delta(\lambda) - \Delta(\nu)$, *where* $\Delta(\lambda)$ *is defined by* (9.1.2)*.*

From now on, whenever we consider an intertwining operator of the form (9.1.5) we assume that Δ is chosen to be $\Delta(\lambda) - \Delta(\nu)$, so that Φ is a $\hat{\mathfrak{g}}$-intertwiner. We will treat Φ as a formal power series in z. Note that the condition that Φ commutes with the action of d can be rewritten as follows:

$$\Phi d - (d \otimes 1)\Phi = z\frac{d}{dz}\Phi. \tag{9.1.8}$$

We introduce the following series, quite usual in physical literature. Let $x \in \mathfrak{g}$ be a homogeneous element with respect to the root decomposition, and let z be a formal variable. Define

$$\tilde{J}_x(z) = \sum_{n\in\mathbb{Z}} x[n] z^{-n}. \tag{9.1.9}$$

Similarly, we define

$$\begin{aligned} \tilde{J}^+_x(z) &= \sum_{n<0} x[n] z^{-n} + \begin{cases} 0, & x \in \mathfrak{n}^+, \\ \frac{1}{2}x, & x \in \mathfrak{h}, \\ x, & x \in \mathfrak{n}^-, \end{cases} \\ \tilde{J}^-_x(z) &= -\sum_{n>0} x[n] z^{-n} - \begin{cases} 0, & x \in \mathfrak{n}^-, \\ \frac{1}{2}x, & x \in \mathfrak{h}, \\ x, & x \in \mathfrak{n}^+. \end{cases} \end{aligned} \tag{9.1.10}$$

Thus, $\tilde{J}_x(z) = \tilde{J}_x^+(z) - \tilde{J}_x^-(z)$. We extend these definitions from homogeneous x to arbitrary $x \in \mathfrak{g}$ by linearity. We consider $\tilde{J}(z)$ as a series in z with coefficients from $U\hat{\mathfrak{g}}$; similar agreement will be applied to other series we introduce.

Usually in the literature a slightly different form of these currents and polarization is used, namely:

$$J_x(z) = \sum x[n]z^{-n-1} = J_x^+(z) - J_x^-(z),$$
$$J_x^+(z) = \sum_{n<0} x[n]z^{-n-1}.$$

It is easy to express $\tilde{J}$ in terms of J and vice versa; however, the form we use will be more convenient in our calculations.

Note that for every $\hat{\mathfrak{g}}$-module V from category $\mathcal{O}$, $\tilde{J}_x^\pm(z)$ can be considered as an operator $V \to \overline{V} \otimes \mathbb{C}[z, z^{-1}]$, where $\overline{V}$ is the completion defined similarly to (9.1.6). Moreover, $\tilde{J}_x^-(z)$ is in fact a well-defined operator $V \to V \otimes \mathbb{C}[z, z^{-1}]$.

We can rewrite the definition of intertwining operator in terms of commutation relations with currents as follows. Let $\Phi(z)$ be an intertwiner of the form (9.1.5), and $u \in L_\mu^*$. Define $\Phi_u(z) : L_{\lambda,K} \to L_{\nu,K}$ by $\Phi_u(z)v = \langle \Phi(z)v, u\rangle$. Then the definition of intertwiner operator takes the form

$$[x[n], \Phi_u(z)] = z^n \Phi_{xu}(z), \tag{9.1.11}$$

where xu denotes the usual action of $x \in \mathfrak{g}$ in the dual representation L_μ^*. Note that

$$\Phi_{xu}(z) = -\langle (1 \otimes x)\Phi(z), u\rangle.$$

Lemma 9.1.3. *Let Φ be a $\hat{\mathfrak{g}}$-intertwining operator of the form* (9.1.5)*. Then we have the following identities of power series in z, w:*

$$[\tilde{J}_x^\pm(w), \Phi_u(z)] = \begin{cases} \frac{w}{z-w}\Phi_{xu}(z), x \in \mathfrak{n}^+, \\ \frac{1}{2}\frac{z+w}{z-w}\Phi_{xu}(z), x \in \mathfrak{h}, \\ \frac{z}{z-w}\Phi_{xu}(z), x \in \mathfrak{n}^-. \end{cases} \tag{9.1.12}$$

Here by definition, we let

$$\frac{1}{z-w} = z^{-1}\sum_{n\geq 0}\left(\frac{w}{z}\right)^n \tag{9.1.13}$$

in the formula involving $\tilde{J}^+$ and

$$\frac{1}{z-w} = -w^{-1}\sum_{n\geq 0}\left(\frac{z}{w}\right)^n \tag{9.1.14}$$

in the formula involving $\tilde{J}^-$*.*

Often we will write $\frac{1}{z-w}, |z| > |w|$ *for the expansion* (9.1.13)*, meaning by this that we are expanding in positive powers of* w/z*; similarly, we write* $\frac{1}{z-w}, |w| > |z|$ *for* (9.1.14)*.*

The proof is straightforward use of (9.1.11) and can be found, for example, in [FR].

Theorem 9.1.4. ([FR]) *In the notations of Lemma* 9.1.3*, we have*

$$(K+h^\vee)z\frac{d}{dz}\Phi_u(z) = \sum_a \vdots\tilde{J}_a(z)\Phi_{au}(z)\vdots + \Phi_{h_\rho u}(z), \tag{9.1.15}$$

where a *runs over an orthonormal basis in* $\mathfrak{g}$*, and*

$$\vdots\tilde{J}_a(z)\Phi_{au}(z)\vdots = \tilde{J}^+_a(z)\Phi_{au}(z) - \Phi_{au}(z)\tilde{J}^-_a(z). \tag{9.1.16}$$

Proof. The simplest way to prove it is to substitute in the commutation relation (9.1.8) between d and Φ_u the expression (9.1.3) for d. After some trivial though boring computations, we get the desired formula (9.1.15).

9.2 The correlation functions on the torus and elliptic KZ equations.

In this section we define the correlation function on the torus. We keep all notations and conventions of the previous sections. Let $L_{\lambda_i,K}, i = 0\dots n$ be a collection of irreducible highest-weight modules such that $\lambda_0 = \lambda_n = \lambda$, and let $\Phi^i(z_i) : L_{\lambda_i,K} \to L_{\lambda_{i-1},K}\otimes L_{\mu_i}(z_i)$ be intertwining operators. Then we can consider the following "correlation function on the torus":

$$\chi_{\lambda,K}(z_1\dots z_n, p, h) = \mathrm{Tr}\,|_{L_{\lambda,K}}\left(\Phi^1(z_1)\dots\Phi^n(z_n)p^{-d}e^{2\pi i h}\right), \tag{9.2.1}$$

where $h\in\mathfrak{h}$. This is nothing but the generalized character χ_Ψ, where

$$\begin{gathered}\Psi = \Phi^1(z_1)\dots\Phi^n(z_n) : L_{\lambda,K} \to L_{\lambda,K}\otimes V,\\ V = L_{\mu_1}\otimes\dots\otimes L_{\mu_n}.\end{gathered} \tag{9.2.2}$$

So far, we consider χ as a formal series in p, z_i with values in the finite-dimensional space V; in fact, it follows from weight considerations that χ takes values in the zero-weight subspace $V[0]$. Later we will study the questions of its convergence. Note also that since Ψ commutes with d we have

$$\sum_i z_i \frac{\partial}{\partial z_i} \chi = 0,$$

thus χ only depends on the ratios z_i/z_j. We will write $\pi_i(x)$ for action of $x \in \mathfrak{g}$ in the i-th factor L_{μ_i} in the product $V = L_{\mu_1} \otimes \ldots \otimes L_{\mu_n}$.

These functions are the main objects of study in this chapter. We call them "correlation functions on the torus": readers familiar with Wess-Zumino-Witten model of conformal field theory immediately recognize that if we let $h = 0$ in (9.2.1) then we get what is known as n-point correlation function on the torus (up to a factor of $p^{-c/24}$, where c is the central charge of Virasoro).

The following theorem follows immediately from Lemma 9.1.2(2).

Theorem 9.2.1. *The space of $\hat{\mathfrak{g}}$-intertwining operators Ψ of the form* (9.2.2) *is isomorphic to the space of all $\mathfrak{g}$-intertwiners $L_\lambda \to L_\lambda \otimes V$.*

Let us consider the limit of χ as $p \to 0$ of this correlation function.

Lemma 9.2.2.

(1) *In the limit $p \to 0$, the trace χ defined by* (9.2.1) *has the following asymptotic:*

$$\chi \sim p^{\Delta(\lambda)} \operatorname{Tr}_{L_\lambda} \left(\Phi^1(z_1) \ldots \Phi^n(z_n) e^{2\pi \mathrm{i} h} \right). \tag{9.2.3}$$

Note that the last trace is taken over the finite-dimensional module $L_\lambda \subset L_{\lambda,K}$, not over the whole module $L_{\lambda,K}$, and is independent of p.

(2) *In the limit $p \to 0, h \to +\mathrm{i}\rho\infty$ so that $pe^{2\pi \mathrm{i}\langle h,\alpha\rangle} \to 0$ for all $\alpha \in R$, $e^{-2\pi \mathrm{i}\langle h,\alpha\rangle} \to 0$ for all $\alpha \in R^+$, the trace χ has the following asymptotics:*

$$\chi \sim p^{\Delta(\lambda)} e^{2\pi \mathrm{i}\langle h,\lambda\rangle} f(z_1, \ldots, z_n), \tag{9.2.4}$$

where

$$f(z_1, \ldots, z_n) = \langle v^*_{\lambda,K}, \Phi^1(z_1) \ldots \Phi^n(z_n) v_{\lambda,K} \rangle \tag{9.2.5}$$

is a function of $z_1, \ldots, z_n$ with values in V.

The limits should be understood in the formal sense (*as limits of formal power series*).

(3) *The function f defined by* (9.2.5) *converges to an analytic function in the region $|z_1| > |z_2| > \cdots > |z_n|$ and satisfies there the following system of*

differential equations (trigonometric form of Knizhnik-Zamolodchikov equations):

$$(K+h^\vee)z_i\frac{\partial}{\partial z_i}f=\Big(\sum_{j\ne i}\frac{z_j\Omega^+_{ij}+z_i\Omega^-_{ij}}{z_i-z_j}-\pi_i(h_{\lambda+\rho})\Big)f, \tag{9.2.6}$$

where

$$\begin{aligned}\Omega^+&=\sum_{\alpha\in R^+}e_\alpha\otimes f_\alpha+\frac{1}{2}\sum_l x_l\otimes x_l,\\ \Omega^-&=\sum_{\alpha\in R^+}f_\alpha\otimes e_\alpha+\frac{1}{2}\sum_l x_l\otimes x_l\end{aligned}$$

and $\Omega^\pm_{ij}=\pi_i\otimes\pi_j(\Omega^\pm)$.

The proof of the first two statements is quite trivial; as for the last one, we refer the reader to [FR].

So far, we considered $\chi_{\lambda,K}$ as a formal series in z_i, p. Let us prove that in fact it defines an analytic function in a certain region. Let $\mathcal{D}$ be the set of all $(z_1,\dots,z_n,h,p), z_i,p\in\mathbb{C},h\in\mathfrak{h}$ satisfying the following conditions:

$$\begin{aligned}&0<|p|<1\\ &|pe^{2\pi\mathrm{i}\langle h,\alpha\rangle}|<1\quad\text{for all }\alpha\in R\\ &|e^{-2\pi\mathrm{i}\langle h,\alpha\rangle}|<1\quad\text{for all }\alpha\in R^+\\ &|z_1|>|z_2|>\ldots>|z_n|>|pz_1|\end{aligned} \tag{9.2.7}$$

Recall the notations $\partial_x, x\in\mathfrak{h},\Delta_\mathfrak{h}$, introduced in Section 2.2 (see also formula (8.1.6)).

THEOREM 9.2.3.

1. *The trace $\chi_{\lambda,K}(z_i,h,p)$ defined by (9.2.1) converges to an analytical function in $\mathcal{D}$.*
2. *This trace satisfies the following differential equation in $\mathcal{D}$:*

$$(\Delta_\mathfrak{h}-2(K+h^\vee)p\frac{\partial}{\partial p}+A)(\chi\hat{\delta})=(\rho,\rho)\chi\hat{\delta}, \tag{9.2.8}$$

where

(9.2.9)

$$A = -\sum_{i,j=1}^{n} \pi_i \otimes \pi_j \Bigg[2 \sum_{\substack{\alpha \in R^+ \\ m \in \mathbb{Z}}} f_\alpha \otimes e_\alpha (z_i/z_j)^m \frac{e^{2\pi \mathrm{i}\langle \alpha, h\rangle} p^m}{(1 - e^{2\pi \mathrm{i}\langle \alpha, h\rangle} p^m)^2} + \sum_{\substack{l=1\dots r \\ m \neq 0}} x_l \otimes x_l (z_i/z_j)^m \frac{p^m}{(1-p^m)^2} \Bigg]$$

$$= -\sum_{i,j=1}^{n} \pi_i \otimes \pi_j \Bigg[2 \sum_{\alpha \in R^+} \phi(-\langle \alpha, h\rangle, \zeta_i - \zeta_j) f_\alpha \otimes e_\alpha + \sum_l \phi_0(\zeta_i - \zeta_j) x_l \otimes x_l \Bigg],$$

where ζ_i *are defined by* $z_i = e^{2\pi \mathrm{i}\zeta_i}$*, the functions* ϕ, ϕ_0 *are defined in the Appendix and* $\hat{\delta}$ *is the affine Weyl denominator given by* (6.2.4) (*see also* (8.1.4))*. We use the following convention:* $\pi_i \otimes \pi_i (a \otimes b) = \pi_i(ab)$*.*

Proof. The proof goes as follows. First, formula (9.2.8), interpreted as a an equality of formal series, follows from more general formula (6.3.8). Next, it is easy to see that in fact all coefficients of the differential equation (9.2.8) converge in $\mathcal{D}$. It follows from the Lemma 9.2.2 and well-known facts about convergence of matrix elements of products of intertwining operators (see [TK]) that in the limit $p \to 0$ the trace χ converges for $|z_1| > \cdots > |z_n|$. These observations along with the standard fact from the theory of ordinary differential equations (namely, that if a formal series satisfies a differential equation with analytic coefficients then this series converges to an analytic solution) prove that $\chi\hat{\delta}$ is an analytic function in $\mathcal{D}$; slight modification of the above arguments, using (6.3.7), allows to prove that in fact χ converges in $\mathcal{D}$.

As we have mentioned before, in the case $n = 1$ the trace χ is independent of z. However, in general case it is not so, and we can study the dependence of this correlation function on z_i, which should generalize the Knizhnik-Zamolodchikov equation (9.2.6).

Theorem 9.2.4. *The product* $\chi\hat{\delta}$*, where* χ *is given by* (9.2.1) *and* $\hat{\delta}$ *is the affine Weyl denominator* (6.2.4)*, satisfies the following system of equations:*

$$(K + h^\vee) \frac{\partial}{\partial \zeta_i} \chi\hat{\delta} = \left(\sum_{j \neq i} r_{ij}(\zeta_i - \zeta_j) - 2\pi \mathrm{i} \sum_l \pi_i(x_l) \partial_{x_l} \right) \chi\hat{\delta}, \tag{9.2.10}$$

where

(9.2.11)

$$r(\zeta) = 2\pi \mathrm{i} \left[\sum_{\alpha \in R^+} \big(e_\alpha \otimes f_\alpha \cdot g(\zeta, \langle \alpha, h\rangle) + f_\alpha \otimes e_\alpha \cdot g(\zeta, -\langle \alpha, h\rangle) \big) - \sum x_l \otimes x_l \cdot \sigma(\zeta) \right],$$

with the elliptic functions g, σ defined in the Appendix. Here the parameters ζ_i are related with z_i by $z_i = e^{2\pi \mathrm{i} \zeta_i}$. As usual, we let $r_{ij} = \pi_i \otimes \pi_j(r)$.

We will call the equations (9.2.10) *elliptic KZ equations*; they are also known under the name Knizhnik-Zamolodchikov-Bernard equations, since they first appeared in the papers of Bernard (see [Be]).

The remaining part of this section is devoted to the proof of this theorem.

PROOF. The idea of proof is quite simple, though the computations are a bit boring. Let us fix $u_1 \in L^*_{\mu_1}, \dots, u_n \in L^*_{\mu_n}$ and consider $\chi(u) = \langle \chi, u_1 \otimes u_2 \otimes \dots \otimes u_n \rangle \in \mathbb{C}$. Then, using the expression for the derivative of Φ, given in Theorem 9.1.4, we can write:

$$(K+h^\vee) z_i \frac{\partial}{\partial z_i} \chi(u) = \sum_a \operatorname{Tr} \left(\Phi^1 \dots \vdots \tilde{J}_a(z_i) \Phi_{a u_i}(z_i) \vdots \dots \Phi^n e^{2\pi \mathrm{i} h} p^{-d} \right) - (\pi_i(h_\rho)\chi)(u),$$

where we write for brevity Φ^k instead of $\Phi^k_{u_k}(z_k)$, and a is any orthonormal basis in $\mathfrak{g}$. Since $\sum a \otimes a = \sum_{\alpha \in R^+} (e_\alpha \otimes f_\alpha + f_\alpha \otimes e_\alpha) + \sum_l x_l \otimes x_l$, where x_l is an orthonormal basis in $\mathfrak{h}$, we get the following expression:

(9.2.11)
$$(K+h^\vee) z_i \frac{\partial}{\partial z_i} \chi(u) = \sum_{\alpha \in R^+} (a^+_\alpha + a^-_\alpha + b^+_\alpha + b^-_\alpha) + \sum_l (c^+_l + c^-_l) - (\pi_i(h_\rho)\chi)(u),$$

where

$$\begin{aligned}
a^+_\alpha &= \operatorname{Tr} \left(\Phi^1 \dots \tilde{J}^+_{e_\alpha}(z_i) \Phi^i_{f_\alpha u_i}(z_i) \dots \Phi^n e^{2\pi \mathrm{i} h} p^{-d} \right), \\
a^-_\alpha &= -\operatorname{Tr} \left(\Phi^1 \dots \Phi^i_{f_\alpha u_i}(z_i) \tilde{J}^-_{e_\alpha}(z_i) \dots \Phi^n e^{2\pi \mathrm{i} h} p^{-d} \right), \\
b^+_\alpha &= \operatorname{Tr} \left(\Phi^1 \dots \tilde{J}^+_{f_\alpha}(z_i) \Phi^i_{e_\alpha u_i}(z_i) \dots \Phi^n e^{2\pi \mathrm{i} h} p^{-d} \right), \\
b^-_\alpha &= -\operatorname{Tr} \left(\Phi^1 \dots \Phi^i_{e_\alpha u_i}(z_i) \tilde{J}^-_{f_\alpha}(z_i) \dots \Phi^n e^{2\pi \mathrm{i} h} p^{-d} \right), \\
c^+_l &= \operatorname{Tr} \left(\Phi^1 \dots \tilde{J}^+_{x_l}(z_i) \Phi^i_{x_l u_i}(z_i) \dots \Phi^n e^{2\pi \mathrm{i} h} p^{-d} \right), \\
c^-_l &= -\operatorname{Tr} \left(\Phi^1 \dots \Phi^i_{x_l u_i}(z_i) \tilde{J}^-_{x_l}(z_i) \dots \Phi^n e^{2\pi \mathrm{i} h} p^{-d} \right).
\end{aligned} \tag{9.2.12}$$

Let us calculate a^+_α. Using the commutation relation

$$\Phi_u(z_j) \tilde{J}^+_{e_\alpha}(z_i) = \tilde{J}^+_{e_\alpha}(z_i) \Phi_u(z_j) - \frac{z_i}{z_j - z_i} \Phi_{e_\alpha u}(z_j)$$

for $|z_j| > |z_i|$ (see Lemma 9.1.3), we get

$$a^+_\alpha = \sum_{j<i} -\frac{z_i}{z_j - z_i}\chi(u_1, \dots, e_\alpha u_j, \dots, f_\alpha u_i, \dots) \\ + \operatorname{Tr}\left(\tilde{J}^+_{e_\alpha}(z_i)\Phi^1 \dots \Phi^i_{f_\alpha u_i}(z_i)\dots\Phi^n e^{2\pi \mathrm{i} h} p^{-d}\right).$$

Using the cyclic property of the trace and the obvious relation

$$e^{2\pi \mathrm{i} h} p^{-d} \tilde{J}^+_{e_\alpha}(z_i) = e^{2\pi \mathrm{i}\langle h,\alpha\rangle} \tilde{J}^+_{e_\alpha}(p z_i) e^{2\pi \mathrm{i} h} p^{-d},$$

we can rewrite it as follows:

$$a^+_\alpha = \sum_{j<i} -\frac{z_i}{z_j - z_i}\chi(u_1, \dots, e_\alpha u_j, \dots, f_\alpha u_i, \dots) \\ + e^{2\pi \mathrm{i}\langle h,\alpha\rangle} \operatorname{Tr}\left(\Phi^1 \dots \Phi^i_{f_\alpha u_i}(z_i)\dots\Phi^n \tilde{J}^+_{e_\alpha}(p z_i) e^{2\pi \mathrm{i} h} p^{-d}\right).$$

Now we again can commute $\tilde{J}^+_{e_\alpha}$ with all Φ^j, and so on. It follows from the definition of the domain $\mathcal{D}$, see (9.2.7), that $\lim_{m\to\infty} e^{2\pi \mathrm{i} m\langle h,\alpha\rangle} \tilde{J}^+_{e_\alpha}(p^m z_i) = 0$. Thus, we get the following expression:

$$a^+_\alpha = \left(\sum_{m\geq 0}\sum_{j<i} \frac{p^m e^{2\pi \mathrm{i} m\langle h,\alpha\rangle} z_i}{p^m z_i - z_j}\pi_i \otimes \pi_j(f_\alpha \otimes e_\alpha) \\ + \sum_{m>0}\sum_{j\geq i} \frac{p^m e^{2\pi \mathrm{i} m\langle h,\alpha\rangle} z_i}{p^m z_i - z_j}\pi_i \otimes \pi_j(f_\alpha \otimes e_\alpha)\right)\chi(u), \tag{9.2.13}$$

where for $j = i$ we let, by definition, $\pi_i \otimes \pi_j(f_\alpha \otimes e_\alpha) = \pi_i(f_\alpha e_\alpha)$. Repeating similar arguments for a^-_α (but moving $\tilde{J}^-$ in the opposite direction), we get the following result:

$$a^+_\alpha + a^-_\alpha = \left(\sum_{m,j} \frac{p^m e^{2\pi \mathrm{i} m\langle h,\alpha\rangle} z_i}{p^m z_i - z_j}\pi_i \otimes \pi_j(f_\alpha \otimes e_\alpha)\right)\chi(u), \tag{9.2.14}$$

where the sum is taken over all pairs $m \in \mathbb{Z}, j = 1, \dots, n$ except the pair $(m = 0, j = i)$, and we have the same convention as before for the term with $j = i$.

In the same way we get the following expression:

$$b^+_\alpha + b^-_\alpha = \left(\sum_{m,j} \frac{z_j e^{-2\pi \mathrm{i} m\langle h,\alpha\rangle}}{p^m z_i - z_j}\pi_i \otimes \pi_j(e_\alpha \otimes f_\alpha)\right)\chi(u), \tag{9.2.15}$$

with the same conventions as before.

As for the terms $c_l^{\pm}$, we must be more careful, since $\lim_{m\to\infty}\tilde{J}^{\pm}_{x_l}(p^{\pm m}z_i) = \pm\frac{1}{2}x_l$. This gives us

$$\begin{aligned} c_l^+ + c_l^- =&\Big(\sum_{m,j}\frac{1}{2}\frac{p^m z_i + z_j}{p^m z_i - z_j}\pi_i\otimes\pi_j(x_l\otimes x_l)\Big)\chi(u) \\ &\qquad + \mathrm{Tr}\big(\Phi^1\dots\Phi^i_{x_l u_i}(z_i)\dots\Phi^n x_l e^{2\pi \mathrm{i} h}p^{-d}\big) \\ =&\Big(\sum_{m,j}\frac{1}{2}\frac{p^m z_i + z_j}{p^m z_i - z_j}\pi_i\otimes\pi_j(x_l\otimes x_l) - \pi_i(x_l)\partial_{x_l}\Big)\chi(u), \end{aligned} \tag{9.2.16}$$

where in addition to the previous conventions we must also specify that the summation is done in the following order: $\sum_{m,j} = \sum_m\sum_j$. This can be replaced by $\sum_{m,j} = \sum_j \text{V.P.}\sum_m$, where the principal value of a series is given by

$$\text{V.P.}\sum_{m\in\mathbb{Z}} a_m = \lim_{N\to+\infty}\sum_{m=-N}^{N} a_m. \tag{9.2.17}$$

Summarizing the previous calculations, we get the following answer:

$$(K+h^\vee)z_i\frac{\partial}{\partial z_i}\chi = \Big(\sum_{j,m} r^m_{ij}(z_i/z_j) - \sum \pi_i(x_l)\partial_{x_l} - \pi_i(h_\rho)\Big)\chi, \tag{9.2.18}$$

where the sum is taken over all pairs $m\in\mathbb{Z}, j = 1,\dots,n$ except the pair $(m = 0, j = i)$, the order of summation is given by $\sum_{m,j} = \sum_j \text{V.P.}\sum_m$, and

$$\begin{aligned} r^m_{ij}(z_i/z_j) = \sum_{\alpha\in R^+}\Big(&\frac{p^m e^{2\pi \mathrm{i} m\langle h,\alpha\rangle}z_i}{p^m z_i - z_j}\pi_i\otimes\pi_j(f_\alpha\otimes e_\alpha) \\ &\qquad + \frac{z_j e^{-2\pi \mathrm{i} m\langle h,\alpha\rangle}}{p^m z_i - z_j}\pi_i\otimes\pi_j(e_\alpha\otimes f_\alpha)\Big) \\ &+ \frac{1}{2}\sum_l\frac{p^m z_i + z_j}{p^m z_i - z_j}\pi_i\otimes\pi_j(x_l\otimes x_l). \end{aligned} \tag{9.2.19}$$

Recall that by definition for $j = i$ we let $\pi_i\otimes\pi_j(a\otimes b) = \pi_i(ab)$.

We can simplify the expression above. Namely, note that the terms with $j = i$ in (9.2.18) have the form:

$$
\begin{aligned}
\sum_{m\neq 0} r^m_{ii}(1) =&\pi_i\left(\sum_{\alpha\in R^+}\sum_{m\neq 0}\Big(\frac{p^m e^{2\pi \mathrm{i} m\langle h,\alpha\rangle}}{p^m-1} f_\alpha e_\alpha + \frac{e^{-2\pi \mathrm{i} m\langle h,\alpha\rangle}}{p^m-1} e_\alpha f_\alpha\Big)\right)\\
&+ \pi_i\Big(\text{V.P.}\sum_l \frac{1}{2}\frac{p^m+1}{p^m-1}x_l^2\Big)\\
=&\pi_i\Big(\sum_{\alpha\in R^+}\sum_{m\neq 0}\frac{e^{-2\pi \mathrm{i} m\langle h,\alpha\rangle}}{p^m-1}[e_\alpha, f_\alpha]\Big)\\
=&\pi_i\Big(\sum_{\alpha\in R^+} -\Big(\sigma(-\langle\alpha,h\rangle)-\frac{1}{2}\Big)h_\alpha\Big)\\
=&\pi_i\Big(h_\rho + \sum_{\alpha\in R^+}\sigma(\langle\alpha,h\rangle)h_\alpha\Big),
\end{aligned}
\tag{9.2.20}
$$

where $\sigma(x)$ is the logarithmic derivative of theta-function defined in the Appendix (see formulas (A2), (A7)).

It turns out that if we write the differential equation for the product $\chi\hat{\delta}$ then these terms cancel. Indeed, since $\hat{\delta}$ does not depend on z, we have

$$
\begin{aligned}
(K+h^\vee)z_i\frac{\partial}{\partial z_i}(\chi\hat{\delta}) &= \Big((K+h^\vee)z_i\frac{\partial}{\partial z_i}\chi\Big)\hat{\delta}\\
&= \hat{\delta}\Big(\sum r^m_{ij}(z_i/z_j) - \sum \pi_i(x_l)\partial_{x_l} - \pi_i(h_\rho)\Big)\chi\\
&= \Big(\sum r^m_{ij}(z_i/z_j) - \sum \pi_i(x_l)\partial_{x_l} - \pi_i(h_\rho)\Big)(\chi\hat{\delta}) + \sum_l \frac{\partial_{x_l}\hat{\delta}}{\hat{\delta}}\pi_i(x_l)(\chi\hat{\delta}).
\end{aligned}
\tag{9.2.21}
$$

Since $\hat{\delta} = a(\tau)\prod_{\alpha\in R^+}\theta_1(\langle\alpha,h\rangle)$ (see (8.1.4)), we have:

$$
\frac{\partial_{x_l}\hat{\delta}}{\hat{\delta}} = \frac{1}{2\pi\mathrm{i}}\sum_{\alpha\in R^+}\frac{\theta_1'(\langle\alpha,h\rangle)}{\theta_1(\langle\alpha,h\rangle)}\langle\alpha,x_l\rangle = -\sum_{\alpha\in R^+}\sigma(\langle\alpha,h\rangle)\langle\alpha,x_l\rangle.
$$

Therefore,

$$
\sum_l \frac{\partial_{x_l}\hat{\delta}}{\hat{\delta}}\pi_i(x_l)(\chi\hat{\delta}) = -\sum_{\alpha\in R^+}\sigma(\langle\alpha,h\rangle)\pi_i(h_\alpha)(\chi\hat{\delta}).
$$

Substituting this in (9.2.21) we see that this term cancels the terms r^m_{ii}. Introducing ζ_i such that $z_i = e^{2\pi\mathrm{i}\zeta_i}$ and using the formulas from the Appendix, we can rewrite $r_{ij}, j \neq i$ in terms of elliptic functions g, σ, which gives us the statement of the Theorem. $\square$

Remark. In fact, all the constructions of this section can be generalized to the case when λ is not necessarily dominant; we could replace the conditions $\lambda \in P^+, K \notin \mathbb{Q}$ by the following conditions:

$$(\widehat{\lambda} + \hat{\rho}, \hat{\alpha}) - \frac{N}{2}(\hat{\alpha}, \hat{\alpha}) = 0 \text{ with } N \in \mathbb{N}, \hat{\alpha} \in \widehat{R}^+$$
$$\text{is only possible if } \hat{\alpha} \in R^+.$$

These conditions ensure that $L_{\lambda,K}$ is induced from the module L_λ over $\mathfrak{g}$, and all the arguments of Sections 9.1, 9.2 can be repeated literally.

We can also include in the consideration integrable modules; in this case, of course, the differential equations are the same, but it is not true that the space of intertwiners Φ of form (9.1.5) is isomorphic to the space of $\mathfrak{g}$-intertwiners (see, for example, [TK], where this is discussed in detail for $\mathfrak{g} = \mathfrak{sl}_2$).

9.3 Monodromies of elliptic KZ equations

In this section we study some properties of the elliptic KZ equations (9.2.10). Let us consider the trace $\chi = \chi(z_1, \dots, z_n, h, p)$ defined by (9.2.1) as a function of $z_1, \dots, z_n, p$ with values in the space

$$\mathbf{V} = (L_{\mu_1} \otimes \dots \otimes L_{\mu_n})[0] \otimes C^\infty(\mathfrak{h}). \tag{9.3.1}$$

Then the elliptic KZ equations can be rewritten in the following form:

$$(K + h^\vee) \frac{\partial}{\partial \zeta_i} \varphi = A_i \varphi, \tag{9.3.2}$$

where $\varphi = \chi \hat{\delta}$, and the operators $A_i : \mathbf{V} \to \mathbf{V}$ are defined by

$$A_i(\zeta_1, \dots, \zeta_n) = \sum_{j \neq i} r_{ij}(\zeta_i - \zeta_j) - 2\pi \mathrm{i} \sum_l \pi_i(x_l) \partial_{x_l}, \tag{9.3.3}$$

where all the notations are as in Theorem 9.2.5; in particular, $r(\zeta)$ is defined by (9.2.11).

Lemma 9.3.1. *The elliptic KZ system is consistent, i.e.,*

$$[(K + h^\vee) \frac{\partial}{\partial \zeta_i} - A_i, (K + h^\vee) \frac{\partial}{\partial \zeta_j} - A_j] = 0 \tag{9.3.4}$$

as operators in $\mathbf{V}$.

PROOF. The proof is based on the fact that we have sufficiently many solutions of this system given by the correlation functions. Indeed, let us consider the commutator (9.3.4); denote it by D_{ij}. It is easy to see that it does not contain derivatives in ζ_i and thus is some operator in $\mathbf{V}$. More precisely, it is some first order differential operator in $\mathfrak{h}$ tensored by some operator in $(L_{\mu_1} \otimes \ldots \otimes L_{\mu_n})[0]$. It follows from Theorem 9.2.4 that it annihilates all the correlation functions $\varphi = \chi\hat{\delta}$.

Now, let us take λ such that $\langle \lambda, \alpha_i^\vee \rangle$ are large enough. It is known that we can do it in such a way that the space of all $\mathfrak{g}$-intertwiners $\Psi : L_\lambda \to L_\lambda \otimes V$ (recall that $V = L_{\mu_1} \otimes \ldots \otimes L_{\mu_n}$) is isomorphic to the zero-weight subspace $V[0]$.

In this case it is easy to show, using Theorem 9.2.1 and some properties of correlation functions on the sphere, that Lemma 9.2.2 can be rewritten as follows:

LEMMA 9.3.2. *Let us fix* $z_1, \ldots, z_n$ *such that* $|z_1| > \cdots > |z_n|$. *Then in the limit* $\tau \to +\mathrm{i}\infty$ *the trace* χ *has the following asymptotics:*

$$\chi(z_1, \ldots, z_n, p, h - \frac{\tau\rho}{h^\vee}) \simeq p^{\Delta(\lambda) - \frac{\langle \lambda, \rho \rangle}{h^\vee}} e^{2\pi \mathrm{i} \langle h, \lambda \rangle} v_\chi$$

for some vector $v_\chi \in V[0]$. *Moreover, for generic* z_i *the mapping* $\chi \mapsto v_\chi$ *is an isomorphism of the space of all traces of the form* (9.2.1) *for all possible choices of* λ_i *and the space* $V[0]$.

Therefore, we see that since the commutator D_{ij} annihilates every trace $\chi\delta$, it also annihilates every function of the form $\hat{\delta} e^{2\pi \mathrm{i} \langle h, \lambda \rangle} v_\chi$ for every sufficiently large λ and small enough p. But this is only possible if it is identically zero. □

Note that straightforward proof of the consistency would require use of Riemann identities for theta-functions, and would necessarily be very tiresome.

Now, let us study some properties of the elliptic KZ system. First of all, we want to know the elliptic properties, i.e. the behavior of the coefficients under the translations $\zeta_i \mapsto \zeta_i + 1, \zeta_i \mapsto \zeta_i + \tau$. Explicit calculation gives the following result:

THEOREM 9.3.3. *Let* $r(\zeta)$ *be defined by* (9.2.11). *Then*

(1) $r_{12}(\zeta) + r_{21}(-\zeta) = 0$, *where* $r_{12} = r, r_{21} = P(r), P(a \otimes b) = b \otimes a$.

(2) r *has a pole of first order at* $\zeta = 0$, *and* $Res_{\zeta=0} r(\zeta) = \Omega$, *where* Ω *is the canonical* $\mathfrak{g}$*-invariant element in* $\mathfrak{g} \otimes \mathfrak{g}$ *defined by* (1.1.3).

(3)

$$\begin{aligned} r(\zeta + 1) &= r(\zeta), \\ r(\zeta + \tau) &= (e^{2\pi \mathrm{i} h} \otimes 1) \circ r(\zeta) \circ (e^{-2\pi \mathrm{i} h} \otimes 1) - 2\pi \mathrm{i} \sum_l x_l \otimes x_l \\ &= (1 \otimes e^{-2\pi \mathrm{i} h}) \circ r(\zeta) \circ (1 \otimes e^{2\pi \mathrm{i} h}) - 2\pi \mathrm{i} \sum x_l \otimes x_l. \end{aligned} \tag{9.3.5}$$

Corollary 9.3.4. *Let A_i be the operator in $\mathbf{V}$ defined by* (9.3.3)*. Then for any $j = 1, \dots, n$ (including $j = i$), we have*

$$\begin{gathered}A_i(\zeta_1, \dots, \zeta_j + 1, \dots \zeta_n) = A_i(\zeta_1, \dots, \zeta_n),\\ A_i(\zeta_1, \dots, \zeta_j + \tau, \dots \zeta_n) = \pi_j(e^{2\pi \mathrm{i} h}) A_i(\zeta_1, \dots, \zeta_n) \pi_j(e^{-2\pi \mathrm{i} h}).\end{gathered} \tag{9.3.6}$$

Thus, the elliptic KZ system is "almost" invariant under the elliptic transformations. Let us modify it so that it becomes really invariant.

Introduce the following operator in $\mathbf{V}$:

$$\psi(\zeta_1, \dots, \zeta_n, h) = \prod_{i<j} \frac{\theta_1\Big(\zeta_i - \zeta_j + \frac{1}{n}(\pi_i(h) - \pi_j(h))\Big)}{\theta_1(\zeta_i - \zeta_j)}; \tag{9.3.7}$$

then

$$\begin{gathered}\psi(\zeta_1, \dots \zeta_i + 1, \dots, \zeta_n) = \psi(\zeta_1, \dots, \zeta_n),\\ \psi(\zeta_1, \dots \zeta_i + \tau, \dots, \zeta_n) = \pi_i(e^{-2\pi \mathrm{i} h}) \psi(\zeta_1, \dots, \zeta_n).\end{gathered} \tag{9.3.8}$$

Then we have the following theorem:

Theorem 9.3.5.

(1) *If φ is a solution of the elliptic KZ system* (9.3.2) *then $\tilde{\varphi} = \psi\varphi$, where ψ is given by* (9.3.7)*, is a solution of the system*

$$(K + h^\vee) \frac{\partial}{\partial \zeta_i} \tilde{\varphi} = \tilde{A}_i(\zeta_1, \dots, \zeta_n) \tilde{\varphi}, \tag{9.3.9}$$

where

$$\tilde{A}_i = \psi A_i \psi^{-1} + (K + h^\vee) \frac{\partial}{\partial \zeta_i} \log \psi. \tag{9.3.10}$$

(2) *This new system is invariant under the translations by $1, \tau$: for every $j \in 1, \dots, n$*

$$\begin{gathered}\tilde{A}_i(\zeta_1, \dots \zeta_j + 1, \dots, \zeta_n) = \tilde{A}_i(\zeta_1, \dots, \zeta_n),\\ \tilde{A}_i(\zeta_1, \dots \zeta_j + \tau, \dots, \zeta_n) = \tilde{A}_i(\zeta_1, \dots, \zeta_n).\end{gathered}$$

This means that we can consider (9.3.9) as a local system on the torus $X = E^n \setminus \{\zeta_i = \zeta_j\}$, where $E = \mathbb{C}/(\mathbb{Z} + \mathbb{Z}\tau)$. Therefore, it makes sense to consider monodromies of this local system. It will give us some representation of the fundamental group of X in the space $\mathbf{V}$. This fundamental group, called the pure braid group of the torus, is well known and can be described in terms of generators and relations. It is convenient to consider a larger group: the braid group of the torus $BT_n = \pi_1(X/S_n)$, where the symmetric group S_n acts in X by permutation of coordinates. Then $\pi_1(X) = \mathrm{Ker}\ \sigma$, where σ is the natural homomorphism $\sigma : BT_n \to S_n$. The local system above defines a representation of BT_n in the following sense: to each $\gamma \in BT_n$ we assign a holonomy operator

$$M(\gamma) : \big(L_{\mu_1} \otimes \ldots \otimes L_{\mu_n}\big)[0] \otimes C^\infty(\mathfrak{h}) \to \big(L_{\mu_{\sigma(1)}} \otimes \ldots \otimes L_{\mu_{\sigma(n)}}\big)[0] \otimes C^\infty(\mathfrak{h}),$$

where $\sigma = \sigma(\gamma)$.

To describe this representation, recall that the braid group BT_n is generated by the elements $T_i, 1 \le i \le n-1, X_j, Y_j, 1 \le j \le n$ (in fact, it suffices to take only T_i, X_1, Y_1). Their geometric meaning is as follows: if we take a base point $\zeta = (\zeta_1, \ldots, \zeta_n)$ such that $\zeta_i = x_i + y_i\tau$, $x_i, y_i \in \mathbb{R}, 0 < x_1 < \ldots < x_n < 1, 0 < y_1 < \ldots < y_n < 1$ then T_i corresponds to a transposition of ζ_i and ζ_{i+1} such that ζ_i passes ζ_{i+1} from the left and X_i, Y_i correspond to ζ_i going around the 1 and τ cycles in negative direction:



T_i



X_i



Y_i

To calculate the monodromies of the local system above we use the fact that we have a large number of solutions given by the functions $\tilde{\varphi} = \psi\hat{\delta}\chi$, where χ is the correlation function on the torus (9.2.1).

Recall the known results about "commutation" of vertex operators. Denote by $H_\lambda^{\nu\mu}$ the space of $\mathfrak{g}$-intertwiners $L_\lambda \to L_\nu \otimes L_\mu$:

$$H_\lambda^{\nu\mu} = \mathrm{Hom}_{\mathfrak{g}}(L_\lambda, L_\nu \otimes L_\mu).$$

Then, due to Lemma 9.1.2, for every $g \in H_\lambda^{\nu\mu}$ we have a unique $\hat{\mathfrak{g}}$-intertwiner $\Phi^g(z) : L_{\lambda,K} \to L_{\nu,K} \otimes L_\mu(z)$. Similarly, if we have weights $\lambda_0, \lambda_1, \lambda_2$ and μ_1, μ_2 and $g_i \in H_{\lambda_i}^{\lambda_{i-1}\mu_i}$ then we can define a $\hat{\mathfrak{g}}$-intertwiner

$$\Phi^{g_1}(z_1)\Phi^{g_2}(z_2) : L_{\lambda_2,K} \to L_{\lambda_0,K} \otimes L_{\mu_1}(z_1) \otimes L_{\mu_2}(z_2). \tag{9.3.11}$$

Summing over all possible λ_1, we can assign to every intertwiner g from

$$\bigoplus_{\lambda_1} H_{\lambda_1}^{\lambda_0\mu_1} \otimes H_{\lambda_2}^{\lambda_1\mu_2} = \mathrm{Hom}_{\mathfrak{g}}(L_{\lambda_2}, L_{\lambda_0} \otimes L_{\mu_1} \otimes L_{\mu_2})$$

(note that this sum is in fact finite) a corresponding $\hat{\mathfrak{g}}$-intertwiner $\Phi^{g^{(1)}}(z_1)\Phi^{g^{(2)}}(z_2) = \sum \Phi^{g_i^{(1)}}(z_1)\Phi^{g_i^{(2)}}(z_2)$ if $g = \sum g_i^{(1)} \otimes g_i^{(2)}$.

Note that the product $\Phi^{g^{(1)}}(z_1)\Phi^{g^{(2)}}(z_2)$ is a well-defined operator for $|z_1| > |z_2|, z_i \notin \mathbb{R}_-$ in the sense that all its matrix coefficients are well-defined analytic functions in this region. It turns out that we can analytically continue this product as a multivalued function in the whole region $z_1 \neq z_2, z_i \neq 0$. Denote by A^+ the analytic continuation along the path when z_1 passes z_2 from the left:



. The following theorem is well-known (see, for example, [MS]).

THEOREM 9.3.6. *As before, let us fix the weights $\lambda_0, \lambda_2, \mu_1, \mu_2$ and define*

$$H^{\mu_1\mu_2} = \bigoplus_{\lambda_1} H^{\lambda_0\mu_1}_{\lambda_1} \otimes H^{\lambda_1\mu_2}_{\lambda_2} = \mathrm{Hom}_{\mathfrak{g}}(L_{\lambda_2}, L_{\lambda_0} \otimes L_{\mu_1} \otimes L_{\mu_2}). \tag{9.3.12}$$

Then there exists some linear operator $R : H^{\mu_1\mu_2} \to H^{\mu_2\mu_1}$ such that for every $g \in H^{\mu_1\mu_2}$ we have the following identity in the region $|z_2| > |z_1|$:

$$A^+(\Phi^{g^{(1)}}(z_1)\Phi^{g^{(2)}}(z_2)) = \Phi^{(Rg)^{(1)}}(z_2)\Phi^{(Rg)^{(2)}}(z_1). \tag{9.3.13}$$

REMARK. In fact, this matrix R coincides (in a suitably chosen basis) with the matrix $\check{R} = PR$, where R is the R-matrix for the corresponding quantum group. This is probably one of the deepest results in conformal field theory; see its discussion, for example, in the book of Kassel [Kas] or in [KL]. However, we won't need it.

This statement allows us to find the monodromies of the elliptic KZ system. Let us recall (see Lemma 9.1.2) that for every collection of $\mathfrak{g}$-intertwiners $g^i : L_{\lambda_i} \to L_{\lambda_{i-1}} \otimes L_{\mu_i}$ we can define the corresponding $\hat{\mathfrak{g}}$-intertwiners $\Phi^{g^i}(z_i) : L_{\lambda_i,K} \to L_{\lambda_{i-1},K} \otimes L_{\mu_i}(z_i)$, and the corresponding trace $\chi^{g^1\otimes\cdots g^n}$, defined by (9.2.1). We can extend this by linearity to define χ^g for arbitrary intertwiner

$$\begin{aligned} g \in H^{\mu_1\cdots\mu_n} &= \mathrm{Hom}_{\mathfrak{g}}\big(L_\lambda, L_\lambda \otimes L_{\mu_1} \otimes \ldots \otimes L_{\mu_n}\big) \\ &= \bigoplus_{\lambda_1,\ldots,\lambda_{n-1}} H^{\lambda\mu_1}_{\lambda_1} \otimes H^{\lambda_1\mu_2}_{\lambda_2} \otimes \ldots \otimes H^{\lambda_{n-1}\mu_n}_{\lambda}. \end{aligned} \tag{9.3.14}$$

Let us fix the base point ζ^0 as before and define the the subspace $\mathbf{V}^0 \subset \mathbf{V}$ which is spanned by the expressions of the form

$$\tilde{\varphi}^g = \chi^g_{\lambda,K}(\zeta^0, p, h)\hat{\delta}(h)\psi(\zeta^0, p, h), \tag{9.3.15}$$

for all possible choices of λ, g. In some sense, we could say that such expressions form a basis in $\mathbf{V}$; however, since this space is infinite-dimensional this requires more accurate approach, and we prefer just to speak of the subspace $\mathbf{V}^0$ in $\mathbf{V}$ formed by finite linear combinations of the functions of the form (9.3.15).

THEOREM 9.3.7. *The representation* $M : BT_n \to \operatorname{End}\mathbf{V}$*, given by the monodromies of the local system* (9.3.9)*, preserves the subspace* $\mathbf{V}^0$ *and in this subspace is given by*

$$\begin{aligned}
&T_i : \tilde{\varphi}^g \mapsto \tilde{\varphi}^{R_{i,i+1}(g)},\\
&X_i : \tilde{\varphi}^g \mapsto e^{2\pi \mathrm{i}(\Delta_i - \Delta_{i-1})}\tilde{\varphi}^g,\\
&Y_i : \tilde{\varphi}^g \mapsto \tilde{\varphi}^{g'},\\
&\qquad g' = R_{i+1,i}R_{i+2,i}\dots R_{n,i}R_{1,i}\dots R_{i-1,i}g.
\end{aligned} \tag{9.3.16}$$

Here R_{ii+1} *is defined as follows:*

$$R_{ii+1} = 1 \otimes \dots R \otimes \dots \otimes 1 : H^{\mu_1\cdots\mu_n} \to H^{\mu_1\cdots\mu_{i+1}\mu_i\cdots\mu_n}, \tag{9.3.17}$$

where $H^{\mu_1\cdots\mu_n}$ *is defined by* (9.3.14) *and the operator* R *was defined in Theorem* 9.3.6.

The numbers Δ_i *are defined as follows: if* $g \in H^{\mu_1\cdots\mu_n}$ *lies in the component* $H^{\lambda\mu_1}_{\lambda_1} \otimes \dots \otimes H^{\lambda_{n-1}\mu_n}_{\lambda}$ *for some* $\lambda_1, \dots, \lambda_{n-1}$ *then* $\Delta_i = \Delta(\lambda_i)$ (*see* (9.1.2)).

PROOF. Let us assume that g is homogeneous, i.e. $g \in H^{\lambda\mu_1}_{\lambda_1} \otimes \dots \otimes H^{\lambda_{n-1}\mu_n}_{\lambda}$ for some $\lambda_1, \dots, \lambda_{n-1}$. Let $\tilde{\varphi}^g(\zeta^0, h, p) \in \mathbf{V}^0$ be a function of the form (9.3.15). Consider the corresponding function $\tilde{\varphi}^g(\zeta, h, p)$, for arbitrary ζ. Then we know that $\tilde{\varphi}^g(\zeta, h, p)$ is a solution of the elliptic KZ system (9.3.9) (see Theorem 9.3.5). Thus, calculation, for example, of $X_i\tilde{\varphi}^g$ is equivalent to finding the result of the analytic continuation of $\tilde{\varphi}^g$ along the path X_i, i.e. the path which connects $\zeta^0 = (\zeta^0_1, \dots, \zeta^0_n)$ and $(\zeta^0_1, \dots, \zeta^0_i - 1, \dots, \zeta^0_n)$. Since the dependence of $\tilde{\varphi}$ on ζ_i has the form

$$\tilde{\varphi} = e^{-2\pi \mathrm{i}\zeta_i(\Delta(\lambda_i) - \Delta(\lambda_{i-1}))} \cdot \textit{some periodic function of } \zeta_i,$$

we get the formula for action of X_i.

The action of T_i can be calculated quite similarly; it is essentially the statement on the commutation of intertwiners (Theorem 9.3.6)

To prove the statement about Y_i, we should commute the operator $\Phi^{g^{(i)}}(z_i)$ with all $\Phi^{g^{(k)}}(z_k), 1 \le k \le i-1$; then use the cyclic property of trace and the relation $p^{-d}e^{2\pi \mathrm{i}h}\Phi(z) = \pi(e^{2\pi \mathrm{i}h})\Phi(pz)p^{-d}e^{2\pi \mathrm{i}h}$. The factor $\pi(e^{2\pi \mathrm{i}h})$ is canceled by the factor appearing in the transformation law for ψ (see (9.3.8)). □

CHAPTER X

PERSPECTIVES AND OPEN QUESTIONS

In this chapter we briefly review the results and list some open questions. So far, we have defined generalized characters for simple finite-dimensional Lie algebras, corresponding quantum groups and affine Lie algebras. We also have derived a number of interesting properties of these characters and have shown that for $\mathfrak{g} = \mathfrak{sl}_n$ and some special choice of U they coincide (after renormalization) with well-known special functions – Jack polynomials; similarly, for $U_q\mathfrak{sl}_n$ we get q-analogue of Jack polynomials – Macdonald's polynomials of type A_{n-1}. This construction allowed us to reprove certain identities for Macdonald's polynomials. We also generalized a large part of this theory to the affine case.

However, a number of interesting questions remain open. Here are some of them.

1. Relation with affine Hecke algebras. In a recent series of papers ([C1, C2]), Ivan Cherednik has shown that Macdonald's polynomials naturally appear in representations of double affine Hecke algebras. Using this technique, he proved Macdonald's inner product conjectures for arbitrary root systems. It would be very interesting to establish a relation of our construction with his work. Morally, it should be something like the duality between representations of GL_n and S_m; so far, no results in this direction are known.

2. Critical level limit. So far, we have assumed in the affine case that the level K of highest-weight modules is not equal to the dual Coxeter number $h^\vee$. However, from many points of view the case $K = -h^\vee$ (critical level) is very interesting. For example, it was shown by E. Frenkel and B. Feigin that (a certain completion of) the algebra $U\hat{\mathfrak{g}}$ has a large center at the critical level. Using this result, Etingof has shown that the elliptic Calogero-Sutherland operator can be included in a large commutative family of differential operators (see [E1]).

However, it is easy to check that one can not define the generalized characters at the critical level in the same manner as we have done before, since in this case there are no intertwining operators of the form we need. One way to study the critical level is to study the asymptotics of generalized characters as $K \to -h^\vee$. For $\mathfrak{g} = \mathfrak{sl}_2$ it was done in the paper [EK3], using explicit integral formulas for the generalized characters. This allowed us to recover the classical formulas for Lamé functions. In general case, no integral formulas are known so far; even if one writes such formulas (which can be done), finding their asymptotics is technically rather difficult.

It seems plausible that there is a direct way to get the generalized characters at the critical level, using the language of coinvariants rather than intertwining operators. Such a construction should be closely related with geometric Langlands correspondence as defined in recent papers of Beilinson and Drinfeld.

3. Inner product in the affine case. In the finite-dimensional case, we had two ways to characterize generalized characters: in terms of differential equations or in terms of inner product (due to the orthogonality theorem). In the affine case, the differential equations work just fine, but defining inner product is quite complicated: any straightforward approach fails (the formulas have singularities). However, some arguments from conformal field theory suggest that there exists a natural inner product on the generalized characters such that we have an analogue of the orthogonality theorem. Also, this inner product should be modular invariant. This question is the subject of the author's present work ([K2]).

Appendix

LIST OF FORMULAS RELATED TO ELLIPTIC FUNCTIONS

In this Appendix, we briefly list the main definitions and formulas for elliptic (theta-, sigma-) functions we are using. Many of them can be found in [WW, Mu]; others can be proved in the usual way.

Throughout this section, we assume that we are given a number $p = e^{2\pi\mathrm{i}\tau} \in \mathbb{C}, |p| < 1$, and all our functions depend on it.

In some formulas, we use the notion of the principal values of a series, which is defined as follows:

$$\text{V.P.} \sum_{n\in\mathbb{Z}} a_n = \lim_{N\to\infty} \sum_{n=-N}^{N} a_n.$$

Functions and identities

$$\theta_1(x) = 2p^{1/8} \sin \pi x \prod_{n\geq 1} (1 - e^{2\pi\mathrm{i}x} p^n)(1 - e^{-2\pi\mathrm{i}x} p^n)(1 - p^n) \tag{A1}$$

$$\sigma(x) = -\frac{1}{2\pi\mathrm{i}} \frac{\theta_1'(x)}{\theta_1(x)} = \sum_{n\geq 0} \frac{p^n e^{2\pi\mathrm{i}x}}{1 - p^n e^{2\pi\mathrm{i}x}} + \sum_{n<0} \frac{1}{1 - p^n e^{2\pi\mathrm{i}x}} + \frac{1}{2} \tag{A2}$$

$$\wp(x) = \frac{1}{x^2} + \sum_{(m,n)\in\mathbb{Z}^2\setminus\{(0,0)\}} \left(\frac{1}{(x - m - n\tau)^2} - \frac{1}{(m + n\tau)^2} \right) \tag{A3}$$

$$\begin{aligned} \sum_{n\in\mathbb{Z}} \frac{p^n e^{2\pi\mathrm{i}x}}{(1 - p^n e^{2\pi\mathrm{i}x})^2} &= \frac{1}{4\pi^2} \partial_x^2 \log \theta_1(x) = \frac{1}{2\pi\mathrm{i}} \sigma'(x) \\ &= -\frac{1}{4\pi^2} \wp(x) + c(\tau) \text{ for some constant } c(\tau) \end{aligned} \tag{A4}$$

$$\eta(\tau) = p^{\frac{1}{24}} \prod_{n\geq 1} (1 - p^n) \tag{A5}$$

$$g(x,\zeta)=\sum_{m\in\mathbb{Z}}\frac{e^{2\pi\mathrm{i}m\zeta}}{1-p^{m}e^{-2\pi\mathrm{i}x}}=-\frac{1}{2\pi\mathrm{i}}\frac{\theta_1(x-\zeta)\theta_1'(0)}{\theta_1(x)\theta_1(\zeta)} \tag{A6}$$

$$\sum_{m\neq 0}\frac{e^{2\pi\mathrm{i}m\zeta}}{1-p^{m}}=\sigma(\zeta)-\frac{1}{2} \tag{A7}$$

$$\frac{1}{2}\mathrm{V.P.}\sum_{m\in\mathbb{Z}}\frac{1+p^{m}e^{2\pi\mathrm{i}\zeta}}{1-p^{m}e^{2\pi\mathrm{i}\zeta}}=\sigma(\zeta) \tag{A8}$$

$$\phi(x,\zeta)=\sum_{m\in\mathbb{Z}}\frac{e^{-2\pi\mathrm{i}x}p^{m}e^{2\pi\mathrm{i}m\zeta}}{(1-p^{m}e^{-2\pi\mathrm{i}x})^{2}}=-\frac{1}{2\pi\mathrm{i}}\frac{\partial}{\partial x}g(x,\zeta) \tag{A9}$$

$$\phi_0(\zeta)=\sum_{m\neq 0}\frac{p^{m}e^{2\pi\mathrm{i}m\zeta}}{(1-p^{m})^{2}}=\frac{\theta_1''(\zeta)}{\pi^{2}\theta_1(\zeta)}-\frac{\theta_1'''(0)}{24\pi^{3}\theta_1(0)}+\frac{1}{12} \tag{A10}$$

Symmetries and elliptic properties

$$\begin{gathered}\theta_1(-x)=-\theta_1(x),\\ \theta_1(x+1)=-\theta_1(x),\qquad \theta_1(x+\tau)=-p^{-1/2}e^{-2\pi\mathrm{i}x}\theta_1(x),\end{gathered} \tag{A11}$$

$$\begin{gathered}\wp(-x)=\wp(x),\\ \wp(x+1)=\wp(x+\tau)=\wp(x),\end{gathered} \tag{A12}$$

$$\begin{gathered}\sigma(-\zeta)=-\sigma(\zeta),\\ \sigma(\zeta+1)=\sigma(\zeta),\qquad \sigma(\zeta+\tau)=\sigma(\zeta)+1,\end{gathered} \tag{A13}$$

$$\begin{gathered}g(x,\zeta)=-g(\zeta,x)=-g(-x,-\zeta),\\ g(x+1,\zeta)=g(x,\zeta),\qquad g(x,\zeta+1)=g(x,\zeta),\\ g(x+\tau,\zeta)=e^{2\pi\mathrm{i}\zeta}g(x,\zeta),\qquad g(x,\zeta+\tau)=e^{2\pi\mathrm{i}x}g(x,\zeta).\end{gathered} \tag{A14}$$

References

[B] Bourbaki, N., *Groupes et algèbres de Lie*, Hermann, Paris, 1969.

[Be] Bernard, D., *On the Wess-Zumino-Witten models on the torus*, Nucl. Phys. **B303** (1988), 77–93.

[BF] Bernard, D., Felder, G., *Fock representations and BRST cohomology in SL(2) current algebra*, Commun. Math. Phys **127** (1990), 145-168.

[C1] Cherednik, I., *The Macdonald constant term conjecture*, IMRN **6** (1993), 165–177.

[C2] ———, *Double affine Hecke algebras and Macdonald's conjectures*, Annals of Math. **141** (1995), 191–216.

[C3] ———, *Macdonald's evaluation conjectures and difference Fourier transform*, preprint, December 1994, q-alg/9412016.

[CK] De Concini, C. and Kac, V.G., *Representations of quantum groups at roots of 1*, Operator algebras, Unitary Representations, Enveloping Algebras and Invariant Theory (A. Connes et al, eds.), Birkhäuser, 1990, pp. 471–506.

[D] van Diejen, J.F., *Commuting difference operators with polynomial eigenfunctions*, to appear in Compos. Math..

[Dr1] Drinfeld, V.G., *Quantum groups*, Proc. Int. Congr. Math., Berkeley, 1986, pp. 798–820.

[Dr2] ———, *On almost cocommutative Hopf algebras*, Leningrad Math.J. **1** (1990), no. 2, 321–342.

[E1] Etingof, P.I., *Quantum integrable systems and representations of Lie algebras, hep-th 9311132*, J. of Math. Physics (1995) (to appear).

[E2] ———, *Central elements for quantum affine algebras and affine Macdonald's operators*, hep-th 9412184 (1994).

[EFK] Etingof, P.I., Frenkel, I.B. and Kirillov, A.A., Jr, *Spherical functions on affine Lie groups*, hep-th/9403168 (to appear in Duke Math. J) (1994).

[EK1] Etingof, P.I. and Kirillov, A.A., Jr, *A unified representation-theoretic approach to special functions, hep-th 9312101*, Functional Anal. and its Applic. **28** (1994), no. 1, 91–94.

[EK2] ———, *Macdonald's polynomials and representations of quantum groups*, Math. Res. Let. **1** (1994), 279–296.

[EK3] ———, *Representations of affine Lie algebras, parabolic differential equations and Láme functions*, Duke Math. J. **74** (1994), 585–614.

[EK4] ———, *On the affine analogue of Jack's and Macdonald's polynomials*, to appear in Duke Math. J. (1994).

[EK5] ———, *Representation-theoretic proof of inner product and symmetry identities for Macdonald's polynomials*, hep-th 9410169, submitted to Comp. Math. (1994).

[ES] Etingof, P.I. and Styrkas, K., *Algebraic integrability of Schrödinger operators and representations of Lie algebras*, preprint, hep-th/9403135 (1994), to appear in Compositio Math..

[FGK] Falceto, F., Gawędzki, K., Kupiainen, A., *Scalar product of current blocks in WZW theory*, Phys. Let. **B260** (1991), 101–108.

[FRT] Reshetikhin, N.Yu., Takhtadzhyan, L.A. and Faddeev, L.D., *Quantization of Lie groups and Lie algebras*, Leningrad Math. J. **1** (1990), no. 1, 193–225.

[HC] Harish-Chandra, *Spherical functions on a semisimple Lie group I*, Amer J. Math. **80** (1958), 241–310.

[HO] Heckman, G.J., Opdam, E.M., *Root systems and hypergeometric functions I*, Compos. Math. **64** (1987), 329–352.

[H1] Heckman, G.J., *Root systems and hypergeometric functions II*, Compos. Math. **64** (1987), 353–373.

[Hu] Humphreys, J.E., *Introduction to Lie algebras and representation theory*, Springer-Verlag, New York, 1972.

[J1] Jimbo, M.A., *A q-difference analogue of $U\mathfrak{g}$ and the Yang-Baxter equation*, Lett. Math. Phys. **10** (1985), 62–69.

[J2] ———, *A q-analogue of $U(\mathfrak{gl}(N+1))$, Hecke algebra and the Yang-Baxter equation*, Lett. Math. Phys. **11** (1986), 247–252.

[K1] Kirillov, A.A., Jr., *On the monodromies of elliptic Knizhnik-Zamolodchikov equations*, preprint (1994).

[K2] ———, *Inner product on conformal blocks and Macdonald's polynomials at roots of unity*, in preparation.

[Ka1] Kac, V.G., *Infinite-dimensional Lie algebras*, 3rd ed., Cambridge Univ. Press, 1990.

[Ka2] ———, *Laplace operators of infinite-dimensional Lie algebras and theta-functions*, Proc. Natl. Acad. Sci. USA **81** (1984), 645-647.

[Kas] Kassel, C., *Quantum groups*, Springer, New York, 1995.

[KL] Kazhdan, D. and Lusztig, G., *Tensor structures arising from affine Lie algebras.* III, J. of AMS **7** (1994), 335–381.

[Koh] Kohno, T., *Monodromy representations of braid groups and Yang-Baxter equations*, Ann. Inst. Fourier **37** (1987), 139-160.

[Kos] Kostant, B., *The principal three-dimensional subgroup and the Betti numbers of a complex simple Lie group*, Amer. J. Math. **81** (1959), 973–1032.

[KZ] Knizhnik, V.G., and Zamolodchikov, A.B., *Current algebra and Wess-Zumino model in two dimensions*, Nucl. Phys. **B247** (1984), 83-103.

[L1] Lusztig, G., *Quantum deformations of certain simple modules over enveloping algebras*, Adv. Math. **70** (1988), 237–249.

[L2] ———, *Introduction to quantum groups*, Birkhäuser, Boston, 1993.

[Lo] Looijenga, E., *Root systems and elliptic curves*, Inv. Math. **33** (1976), 17–32.

[M1] Macdonald, I.G., *A new class of symmetric functions*, Publ. I.R.M.A. Strasbourg, 372/S-20, Actes 20 Séminaire Lotharingien (1988), 131-171.

[M2] ———, *Orthogonal polynomials associated with root systems*, preprint (1988).

[Ma] Matsuo, A., *Integrable connections related to zonal spherical functions*, Inv. Math. **110** (1992), 95–121.

[Mi] Mimachi, K., *Macdonald's operators from the center of the quantized universal enveloping algebra $U_q(gl(N))$*, preprint (September 1994).

[MS] Moore, G., Seiberg, N., *Classical and quantum conformal field theory*, Com. Math. Phys. **123** (1989), 177–254.

[Mu] Mumford, D., *Tata Lectures on Theta I*, Birkhäuser, 1983.

[N] Noumi, M., *Macdonald's symmetric polynomials as zonal spherical functions on some quantum homogeneous spaces*, preprint, Dept. of Math. Sciences, Univ. of Tokyo, Japan, Oct. 1993, (to appear in Adv. in Math.).

[O1] Opdam, E.M., *Root systems and hypergeometric functions III*, Compos. Math. **67** (1988), 21–49.

[O2] ———, *Root systems and hypergeometric functions IV*, Compos. Math. **67** (1988), 191–207.

[OOS] Ochiai, H., Oshima, T., and Sekiguchi, H., *Commuting families of symmetric differential operators*, Proc. of the Japan Acad. **70, Ser. A** (1994), no. 2, 62–68.

[OP] Olshanetsky, M.A., and Perelomov, A.M., *Quantum integrable systems related to Lie algebras*, Phys. Rep. **94** (1983), 313-404.

[R] Reshetikhin, N.Yu., *Quasitriangle Hopf algebras and invariants of tangles*, Leningrad Math J. **1** (1990), no. 2, 491-513.

[Su] Sutherland, B., *Exact results for quantum many-body problem in one dimension*, Phys. Rep. **A5** (1972), 1375–1376.

[T] Tanisaki, T., *Killing forms, Harish-Chandra isomorphisms and universal R-matrices for quantum algebras*, Infinite Analysis, part A and part B (Kyoto, 1991), Adv. Ser. Math. Phys. 17, World Scientific, pp. 941–961.

[TK] Tsuchiya, A., Kanie, Y., *Vertex operators in conformal field theory on P^1 and monodromy representations of braid group*, Adv. Stud. Pure Math. **16** (1988), 297-372.

[V] Vilenkin, N. Ja., *Special functions and the theory of group representations*, Amer. Math. Soc., Providence, 1968.

[VK] Vilenkin, N. Ja., and Klimyk, A.U., *Representations of Lie groups and special functions*, Kluwer Academic Publishers, 1991.

[W] Warner, G., *Harmonic analysis on semi-simple Lie groups II*, Springer-Verlag, 1972.

[WW] Whittaker, E.T., Watson, G.N., *Course of modern analysis*, 4th ed., Cambridge Univ. Press, 1958.

arXiv:q-alg/9503012v1 21 Mar 1995

Abstract

Traces of intertwining operators and Macdonald's polynomials

Alexander A. Kirillov, Jr.

May 1995

Let $\Phi : V \to V \otimes U$ be an intertwining operator between representations of a simple Lie algebra (quantum group, affine Lie algebra). We define its generalized character to be the following function on the Cartan subalgebra with values in U: $\chi_\Phi(h) = \mathrm{Tr}_V(\Phi e^h)$. This is a generalization of usual characters. These generalized characters are a rich source of special functions, possessing many interesting properties; for example, they are common eigenfunctions of a family of commuting differential (difference) operators. We show that the special functions that can be obtained this way include Macdonald's polynomials of type A, and this technique allows to prove inner product and symmetry identities for these polynomials (though proved earlier by other methods).

Generalized characters for affine Lie algebras are closely related with so-called correlation functions on the torus in the Wess-Zumino-Witten (WZW) model of conformal field theory. We derive differential equations satisfied by these correlation functions (elliptic Knizhnik-Zamolodchikov, or Knizhnik-Zamolodchikov-Bernard equations) and study their monodromies.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$-TEX

Traces of intertwining operators and Macdonald's polynomials

A Dissertation
Presented to the Faculty of the Graduate School
of
Yale University
in Candidacy for the Degree of
Doctor of Philosophy

by
Alexander A. Kirillov, Jr.

Dissertation Director: Professor Igor B. Frenkel

May 1995

©1995 by Alexander A. Kirillov

All rights reserved

CONTENTS

Acknowledgements **1**

Introduction **2**

Chapter I. Basic definitions

1.1 Simple Lie algebras and their representations 5

1.2 Quantum groups 9

Chapter II. Generalized characters

2.1 Weighted traces of intertwiners and orthogonality theorem 13

2.2 Center of $U\mathfrak{g}$ and commuting differential operators 15

2.3 Center of $U_q\mathfrak{g}$ and difference operators 18

2.4 Generalized characters as spherical functions 20

Chapter III. Jacobi polynomials

3.1 Jacobi polynomials and corresponding differential operators 24

3.2 Jack polynomials as generalized characters 27

Chapter IV. Macdonald's polynomials as generalized characters

4.1 Definition of Macdonald's polynomials 30

4.2 Macdonald's polynomials of type A as generalized characters 32

4.3 The center of $U_q\mathfrak{sl}_n$ and Macdonald's operators 35

Chapter V. Inner product and symmetry identities for Macdonald's polynomials

5.1 Inner product identities 38
5.2 Algebra of intertwiners 41
5.3 Symmetry identities 48

Chapter VI. Generalized characters for affine Lie algebras

6.1 Affine Lie algebras 51
6.2 Group algebra of the weight lattice 53
6.3 Generalized characters for affine Lie algebras 56

Chapter VII. Affine Jacobi polynomials

7.1 Definition of affine Jacobi polynomials 60
7.2 Affine Jack polynomials as generalized characters 64

Chapter VIII. Modular properties of affine Jacobi polynomials

8.1 Functional interpretation of $\mathbb{C}[\widehat{P}]$ 67
8.2 Normalized characters and modular invariance 70

Chapter IX. Correlation functions on the torus and elliptic KZ equations

9.1 Intertwiners and currents 75
9.2 Correlation functions on the torus and differential equations 79
9.3 Monodromies of elliptic KZ equations 86
Chapter X. Perspectives and open questions 93
Appendix: List of formulas related to elliptic functions 95
References 97